



**Physics 132**

**2<sup>nd</sup> Hour Exam**  
**Time: 85:00 min**

**2<sup>nd</sup> Semester 2017/2018**  
**Date: 13 /5/2018**

**Student Name:----- Student NO-----**

✓	Sec	Instructor Name	Classes Time	✓	Sec	Instructor Name	Classes Time
○	1	Areej Abdel Rahman	S 9:00-9:50	○	8	Hazem Abu Sara	W 12:00-12:50
○	2	Hazem Abu Sara	M 12:00-12:50	○	9	Wael Karain	W 9:00-9:50
○	3	Areej Abdel Rahman	M 14:00-14:50	○	10	Abdallah Sayyed	W 14:00-14:50
○	4	Abdallah Sayyed	S 14:00-14:50	○	11	Abdallah Sayyed	W 11:00-11:50
○	5	Dua' Abu Mura	S 14:00-14:50	○	12	Areej Abdel Rahman	W 12:00-12:50
○	6	ghassan abbas	W 13:00-13:50	○			
○	7	Areej Abdel Rahman	M 15:00-15:50				

**Answer Sheet:**

Q#	a	b	c	d	e
1					
2					
3					
4					
5					
6					
7					
8					
9					
10					
11					
12					
13					
14					
15					
16					

## Useful Formulae and Constants

1.  $\epsilon_0 = 8.85 \times 10^{-12} \text{C}^2/\text{N}\cdot\text{m}^2$
2.  $e = 1.6 \times 10^{-19} \text{C}$
3.  $\mu_0 = 4\pi \times 10^{-7} \text{T}\cdot\text{m}/\text{A}$
4.  $m_e = 9.11 \times 10^{-31} \text{Kg}$
5.  $m_p = 1.67 \times 10^{-27} \text{Kg}$
6.  $g = 10 \text{m}/\text{s}^2$
7.  $\text{eV} = 1.6 \times 10^{-19} \text{J}$

### Capacitance

1.  $q = CV$
2.  $C = \frac{\epsilon_0 A}{d}$
3.  $\frac{1}{C_{eq}} = \sum \frac{1}{C_i}$  (Series connection)
4.  $C_{eq} = \sum C_i$  (Parallel connection)
5.  $U = \frac{q^2}{2C}$
6.  $u = \frac{1}{2} \epsilon_0 E^2$
7.  $\epsilon_0 \oint \kappa \vec{E} \cdot d\vec{A} = q$

### Current and Resistance

8.  $\vec{J} = ne\vec{v}_d$
9.  $\vec{E} = \rho\vec{J}$
10.  $R = \frac{\rho L}{A}$
11.  $P = i^2 R$

### Circuits

12.  $\epsilon = \frac{dW}{dq}$
13.  $V = \epsilon(1 - e^{-\frac{t}{RC}})$


### Magnetic Fields

14.  $\vec{F} = q\vec{v} \times \vec{B}$
15.  $qvB = \frac{mv^2}{r}$
16.  $f = \frac{qB}{2\pi m}$
17.  $\vec{F} = i\vec{l} \times \vec{B}$
18.  $\vec{\tau} = \vec{\mu} \times \vec{B}$
19.  $U = -\vec{\mu} \cdot \vec{B}$
20.  $W_a = U_f - U_i$

- 1- A potential difference of  $200V$  is applied to a series connection of two capacitors of capacitance  $C_1 = 3 \mu F$  and  $C_2 = 6 \mu F$ . What is the charge on  $C_1$ ?
- a)  $1200 \mu C$
  - b)  $5400 \mu C$
  - c)  $400 \mu C$  \*
  - d)  $200 \mu C$
  - e)  $600 \mu C$
- 2- Two capacitors,  $C_1 = 4 \mu F$  is charged so its charge is  $q_1 = 100 \mu C$  and  $C_2 = 6 \mu F$  is uncharged. The two capacitors are then connected in parallel. Find the charge on  $C_1$ ?
- a)  $100 \mu C$
  - b)  $32 \mu C$
  - c)  $48 \mu C$
  - d)  $40 \mu C$  \*
  - e)  $60 \mu C$
- 3- A certain capacitor has a capacitance of  $C$ . After it is charged to a charge  $q$  and isolated, the two plates are pull apart so its capacitance becomes  $C/3$ . Find the work done by the agent?
- a)  $-\frac{q^2}{2C}$
  - b)  $\frac{q^2}{2C}$
  - c)  $\frac{q^2}{C}$  \*
  - d)  $-\frac{q^2}{C}$
  - e) *zero*

- 4- A slab of copper of thickness ( $b$ ) is thrust into a parallel plate capacitor of plate area ( $A$ ) and plate separation ( $d$ ) as shown in the figure. What is the capacitance after the slab is introduced?

- a)  $\frac{\epsilon_0 A}{d}$   
 b)  $\frac{\epsilon_0 A}{b}$   
 c)  $\frac{\epsilon_0 A}{d+b}$   
 d)  $\frac{\epsilon_0 A}{d-b}$ \*  
 e) 0



- 5- A isolated conducting sphere whose radius is  $R$  and has a charge  $Q$ . How much potential energy is stored in the electric field of this charged conductor?

- a)  $\frac{kQ^2}{R}$   
 b)  $\frac{kQ^2}{2R}$ \*  
 c)  $\frac{kQ}{R}$   
 d)  $\frac{kQ}{2R}$   
 e)  $\frac{kQ}{R^2}$

- 6- A certain wire has a resistance  $R$ . What is the resistance of a second wire, made of the same material, that is quarter (¼) as long and has quarter the radius

- a)  $2R$   
 b)  $4R$ \*  
 c)  $R/2$   
 d)  $R/4$   
 e)  $R$

7- What is the current in a wire of radius ( $a$ ) if the magnitude of the current density is variable and given by  $J = br^2$ , in which  $b$  is constant and  $r$  is the radial distance?


- a)  $\pi ba^4/2^*$
- b)  $\pi ba^3/3$
- c)  $\pi ba^3/4$
- d)  $\pi ba^4/4$
- e)  $\pi ba^4$

8- What is the value of 1 *Ampere.hour*?

- a)  $1.6 \times 10^{-19} C$
- b) 3600 J
- c) 3600 Watt
- d) 3600 *electrons*
- e) 3600 C\*


9- In the circuit shown, the ideal batteries have *emfs*,  $\varepsilon_1 = 5 V$ ,  $\varepsilon_2 = 20 V$  and  $R_1 = 3 \Omega$ ,  $R_2 = 2 \Omega$ . If the potential at  $a$  is 10 V, what is the potential at  $b$ ?

- a)  $-1 V^*$
- b)  $+1 V$
- c)  $-11 V$
- d)  $+11 V$
- e)  $-21 V$



10- In the circuit shown find  $I$ ?

- a) zero
- b) 3.5 A
- c) 7 A\*
- d) 2 A
- e) 5 A



11- A capacitor with initial charge  $q_0$  is discharged through a resistor. What is the time taken by the capacitor to lose one-quarter of its charge?

- a)  $0.693\tau$
- b)  $0.40\tau$
- c)  $0.29\tau$ \*
- d)  $0.71\tau$
- e)  $0.60\tau$

12- In an RC series circuit,  $\epsilon = 10 V$ , resistance  $R = 2 M\Omega$ , and capacitance  $c = 2.5 \mu F$ , the circuit is closed at  $t = 0$  to begin charging. Find the voltage across the capacitor at  $t = 15 s$ ?

- a) 3.7 V
- b) 6.3 V
- c) 1.4 V
- d) 8.6 V
- e) 9.5 V\*

13- At one instant,  $\vec{v} = (200\hat{i} + 500\hat{k}) \text{ m/s}$  is the velocity of a proton in a uniform magnetic field  $\vec{B} = (0.05\hat{i} - 0.15\hat{k}) \text{ T}$ . At that instant find the force on the proton?


- a)  $+72 \times 10^{-19}\hat{k} \text{ N}$
- b)  $-72 \times 10^{-19}\hat{k} \text{ N}$
- c)  $-45 \times 10^{-19}\hat{k} \text{ N}$
- d)  $+88 \times 10^{-19}\hat{j} \text{ N}^*$
- e)  $+88 \times 10^{-19}\hat{i} \text{ N}$

14- An electron moves with speed  $v$  into a region of uniform magnetic field  $B$ . The angle between them is  $= 85^\circ$ . Describe the motion of the electron and find the periodic time?

- a) Uniform circular motion,  $T = \frac{2\pi m}{eB}$
- b) Nonuniform circular motion,  $T = \frac{2\pi m}{eB} \cos 85$
- c) Helical motion,  $T = \frac{2\pi m}{eB} \cos 85$
- d) Helical motion,  $T = \frac{2\pi m}{eB}^*$
- e) Helical motion,  $T = \frac{2\pi m}{eB} \sin 85$

15- The coil in the figure, which is parallel to the  $xz$  - plane, carries current  $I = 2 \text{ A}$  in the direction indicated, has 3 turns and area of  $4 \times 10^{-3} \text{ m}^2$ , and lies in a uniform magnetic field  $\vec{B} = (-3\hat{i} + 2\hat{j} - 4\hat{k}) \text{ mT}$ . What is the potential energy of the coil in the magnetic field?

- a)  $-48 \mu\text{J}^*$
- b)  $+24 \mu\text{J}$
- c)  $+96 \mu\text{J}$
- d)  $+72 \mu\text{J}$
- e) zero



16- A magnetic field CANNOT:

- a) Exert a force on a charge
- b) Accelerate a charge
- c) Change the momentum of a charge
- d) Change the kinetic energy of a charge\*
- e) exist