

## Chapter 38: PHOTONS AND MATTER WAVES

- The units of the Planck constant  $h$  are those of:
  - energy
  - power
  - momentum
  - angular momentum
  - frequencyans: D
- If  $h$  is the Planck constant, then  $\hbar$  is:
  - $2\pi h$
  - $2h$
  - $h/2$
  - $h/2\pi$
  - $2h/\pi$ans: D
- The quantization of energy,  $E = nhf$ , is not important for an ordinary pendulum because:
  - the formula applies only to mass-spring oscillators
  - the allowed energy levels are too closely spaced
  - the allowed energy levels are too widely spaced
  - the formula applies only to atoms
  - the value of  $h$  for a pendulum is too largeans: B
- The frequency of light beam A is twice that of light beam B. The ratio  $E_A/E_B$  of photon energies is:
  - 1/2
  - 1/4
  - 1
  - 2
  - 4ans: D
- The wavelength of light beam B is twice the wavelength of light beam A. The energy of a photon in beam A is:
  - half the energy of a photon in beam B
  - one-fourth the energy of a photon in beam B
  - equal to the energy of a photon in beam B
  - twice the energy of a photon in beam B
  - four times the energy of a photon in beam Bans: A

6. A photon in light beam A has twice the energy of a photon in light beam B. The ratio  $p_A/p_B$  of their momenta is:
- A.  $1/2$
  - B.  $1/4$
  - C. 1
  - D. 2
  - E. 4

ans: D

7. Which of the following electromagnetic radiations has photons with the greatest energy?
- A. blue light
  - B. yellow light
  - C. x rays
  - D. radio waves
  - E. microwaves

ans: C

8. Which of the following electromagnetic radiations has photons with the greatest momentum?
- A. blue light
  - B. yellow light
  - C. x rays
  - D. radio waves
  - E. microwaves

ans: C

9. Rank following electromagnetic radiations according to the energies of their photons, from least to greatest.
- 1. blue light
  - 2. yellow light
  - 3. x rays
  - 4. radio waves

- A. 1, 2, 3, 4
- B. 4, 2, 1, 3
- C. 4, 1, 2, 3
- D. 3, 2, 1, 4
- E. 3, 1, 2, 4

ans: B

10. The intensity of a uniform light beam with a wavelength of 500 nm is  $2000 \text{ W/m}^2$ . The photon flux (in number/ $\text{m}^2 \cdot \text{s}$ ) is about:
- A.  $5 \times 10^{17}$
  - B.  $5 \times 10^{19}$
  - C.  $5 \times 10^{21}$
  - D.  $5 \times 10^{23}$
  - E.  $5 \times 10^{25}$


ans: C

11. The concentration of photons in a uniform light beam with a wavelength of 500 nm is  $1.7 \times 10^{13} \text{ m}^{-3}$ . The intensity of the beam is:
- A.  $6.7 \times 10^{-6} \text{ W/m}^2$
  - B.  $1.0 \times 10^3 \text{ W/m}^2$
  - C.  $2.0 \times 10^3 \text{ W/m}^2$
  - D.  $4.0 \times 10^3 \text{ W/m}^2$
  - E.  $3.2 \times 10^2 \text{ W/m}^2$
- ans: C
12. Light beams A and B have the same intensity but the wavelength associated with beam A is longer than that associated with beam B. The photon flux (number crossing a unit area per unit time) is:
- A. greater for A than for B
  - B. greater for B than for A
  - C. the same for A and B
  - D. greater for A than for B only if both have short wavelengths
  - E. greater for B than for A only if both have short wavelengths
- ans: A
13. In a photoelectric effect experiment the stopping potential is:
- A. the energy required to remove an electron from the sample
  - B. the kinetic energy of the most energetic electron ejected
  - C. the potential energy of the most energetic electron ejected
  - D. the photon energy
  - E. the electric potential that causes the electron current to vanish
- ans: E
14. In a photoelectric effect experiment at a frequency above cut off, the stopping potential is proportional to:
- A. the energy of the least energetic electron before it is ejected
  - B. the energy of the least energetic electron after it is ejected
  - C. the energy of the most energetic electron before it is ejected
  - D. the energy of the most energetic electron after it is ejected
  - E. the electron potential energy at the surface of the sample
- ans: D
15. In a photoelectric effect experiment at a frequency above cut off, the number of electrons ejected is proportional to:
- A. their kinetic energy
  - B. their potential energy
  - C. the work function
  - D. the frequency of the incident light
  - E. the number of photons that hit the sample
- ans: E

16. In a photoelectric effect experiment no electrons are ejected if the frequency of the incident light is less than  $A/h$ , where  $h$  is the Planck constant and  $A$  is:
- the maximum energy needed to eject the least energetic electron
  - the minimum energy needed to eject the least energetic electron
  - the maximum energy needed to eject the most energetic electron
  - the minimum energy needed to eject the most energetic electron
  - the intensity of the incident light

ans: D

17. The diagram shows the graphs of the stopping potential as a function of the frequency of the incident light for photoelectric experiments performed on three different materials. Rank the materials according to the values of their work functions, from least to greatest.



- 1, 2, 3
- 3, 2, 1
- 2, 3, 1
- 2, 1, 3
- 1, 3, 2

ans: A

18. The work function for a certain sample is 2.3 eV. The stopping potential for electrons ejected from the sample by  $7.0 \times 10^{14}$ -Hz electromagnetic radiation is:
- 0
  - 0.60 V
  - 2.3 V
  - 2.9 V
  - 5.2 V

ans: B

19. The stopping potential for electrons ejected by  $6.8 \times 10^{14}$ -Hz electromagnetic radiation incident on a certain sample is 1.8 V. The kinetic energy of the most energetic electrons ejected and the work function of the sample, respectively, are:
- 1.8 eV, 2.8 eV
  - 1.8 eV, 1.0 eV
  - 1.8 eV, 4.6 eV
  - 2.8 eV, 1.0 eV
  - 1.0 eV, 4.6 eV

ans: B

20. Separate Compton effect experiments are carried out using visible light and x rays. The scattered radiation is observed at the same scattering angle. For these experiments:
- A. the x rays have the greater shift in wavelength and the greater change in photon energy
  - B. the two radiations have the same shift in wavelength and the x rays have the greater change in photon energy
  - C. the two radiations have the same shift in wavelength and the visible light has the greater change in photon energy
  - D. the two radiations have the same shift in wavelength and the same change in photon energy
  - E. the visible light has the greater shift in wavelength and the greater shift in photon energy
- ans: B
21. In Compton scattering from stationary particles the maximum change in wavelength can be made smaller by using:
- A. higher frequency radiation
  - B. lower frequency radiation
  - C. more massive particles
  - D. less massive particles
  - E. particles with greater charge
- ans: C
22. Of the following, Compton scattering from electrons is most easily observed for:
- A. microwaves
  - B. infrared light
  - C. visible light
  - D. ultraviolet light
  - E. x rays
- ans: E
23. In Compton scattering from stationary electrons the largest change in wavelength occurs when the photon is scattered through:
- A.  $0^\circ$
  - B.  $22.5^\circ$
  - C.  $45^\circ$
  - D.  $90^\circ$
  - E.  $180^\circ$
- ans: E
24. In Compton scattering from stationary electrons the frequency of the emitted light is independent of:
- A. the frequency of the incident light
  - B. the speed of the electron
  - C. the scattering angle
  - D. the electron recoil energy
  - E. none of the above
- ans: E

25. In Compton scattering from stationary electrons the largest change in wavelength that can occur is:
- A.  $2.43 \times 10^{-15}$  m
  - B.  $2.43 \times 10^{-12}$  m
  - C.  $2.43 \times 10^{-9}$  m
  - D. dependent on the frequency of the incident light
  - E. dependent on the work function
- ans: B

26. Electromagnetic radiation with a wavelength of  $5.7 \times 10^{-12}$  m is incident on stationary electrons. Radiation that has a wavelength of  $6.57 \times 10^{-12}$  m is detected at a scattering angle of:
- A.  $10^\circ$
  - B.  $121^\circ$
  - C.  $40^\circ$
  - D.  $50^\circ$
  - E.  $69^\circ$
- ans: D

27. Electromagnetic radiation with a wavelength of  $3.5 \times 10^{-12}$  m is scattered from stationary electrons and photons that have been scattered through  $50^\circ$  are detected. An electron from which one of these photons was scattered receives an energy of:
- A. 0
  - B.  $1.1 \times 10^{-14}$  J
  - C.  $1.9 \times 10^{-14}$  J
  - D.  $2.3 \times 10^{-14}$  J
  - E.  $1.3 \times 10^{-13}$  J
- ans: B

28. Electromagnetic radiation with a wavelength of  $3.5 \times 10^{-12}$  m is scattered from stationary electrons and photons that have been scattered through  $50^\circ$  are detected. After a scattering event the magnitude of the electron's momentum is:
- A. 0
  - B.  $1.5 \times 10^{-22}$  kg · m/s
  - C.  $2.0 \times 10^{-22}$  kg · m/s
  - D.  $2.2 \times 10^{-22}$  kg · m/s
  - E.  $8.7 \times 10^{-23}$  kg · m/s
- ans: B

29. Consider the following:
1. a photoelectric process in which some emitted electrons have kinetic energy greater than  $hf$ , where  $f$  is the frequency of the incident light.
  2. a photoelectric process in which all emitted electrons have energy less than  $hf$ .
  3. Compton scattering from stationary electrons for which the emitted light has a wavelength that is greater than that of the incident light.
  4. Compton scattering from stationary electrons for which the emitted light has a wavelength that is less than that of the incident light.

The only possible processes are:

- A. 1
- B. 3
- C. 1 and 3
- D. 2 and 3
- E. 2 and 4

ans: D

30. J. J. Thompson's measurement of  $e/m$  for electrons provides evidence of the:
- A. wave nature of matter
  - B. particle nature of matter
  - C. wave nature of radiation
  - D. particle nature of radiation
  - E. transverse wave nature of light

ans: B

31. Evidence for the wave nature of matter is:
- A. electron diffraction experiments of Davisson and Germer
  - B. Thompson's measurement of  $e/m$
  - C. Young's double slit experiment
  - D. the Compton effect
  - E. Lenz's law

ans: A

32. Which of the following is NOT evidence for the wave nature of matter?
- A. The photoelectric effect
  - B. The diffraction pattern obtained when electrons pass through a slit
  - C. Electron tunneling
  - D. The validity of the Heisenberg uncertainty principle
  - E. The interference pattern obtained when electrons pass through a two-slit system

ans: A

33. Of the following which is the best evidence for the wave nature of matter?
- A. The photoelectric effect
  - B. The Compton effect
  - C. The spectral radiance of cavity radiation
  - D. The relationship between momentum and energy for an electron
  - E. The reflection of electrons by crystals

ans: E

34. Monoenergetic electrons are incident on a single slit barrier. If the energy of each incident electron is increased the central maximum of the diffraction pattern:
- A. widens
  - B. narrows
  - C. stays the same width
  - D. widens for slow electrons and narrows for fast electrons
  - E. narrows for slow electrons and widens for fast electrons
- ans: B

35. A free electron and a free proton have the same kinetic energy. This means that, compared to the matter wave associated with the proton, the matter wave associated with the electron has:
- A. a shorter wavelength and a greater frequency
  - B. a longer wavelength and a greater frequency
  - C. a shorter wavelength and the same frequency
  - D. a longer wavelength and the same frequency
  - E. a shorter wavelength and a smaller frequency
- ans: D

36. A free electron and a free proton have the same momentum. This means that, compared to the matter wave associated with the proton, the matter wave associated with the electron:
- A. has a shorter wavelength and a greater frequency
  - B. has a longer wavelength and a greater frequency
  - C. has the same wavelength and the same frequency
  - D. has the same wavelength and a greater frequency
  - E. has the same wavelength and a smaller frequency
- ans: D

37. A free electron and a free proton have the same speed. This means that, compared to the matter wave associated with the proton, the matter wave associated with the electron:
- A. has a shorter wavelength and a greater frequency
  - B. has a longer wavelength and a greater frequency
  - C. has the same wavelength and the same frequency
  - D. has the same wavelength and a greater frequency
  - E. has a longer wavelength and a smaller frequency
- ans: E

38. Consider the following three particles:
1. a free electron with speed  $v_0$
  2. a free proton with speed  $v_0$
  3. a free proton with speed  $2v_0$

Rank them according to the wavelengths of their matter waves, least to greatest.

- A. 1, 2, 3
  - B. 3, 2, 1
  - C. 2, 3, 1
  - D. 1, 3, 2
  - E. 1, then 2 and 3 tied
- ans: B



39. Consider the following three particles:

1. a free electron with kinetic energy  $K_0$
2. a free proton with kinetic energy  $K_0$
3. a free proton with kinetic energy  $2K_0$

Rank them according to the wavelengths of their matter waves, least to greatest.

- A. 1, 2, 3
- B. 3, 2, 1
- C. 2, 3, 1
- D. 1, 3, 2
- E. 1, then 2 and 3 tied

ans: B

40. A free electron has a momentum of  $5.0 \times 10^{-24} \text{ kg} \cdot \text{m/s}$ . The wavelength of its wave function is:

- A.  $1.3 \times 10^{-8} \text{ m}$
- B.  $1.3 \times 10^{-10} \text{ m}$
- C.  $2.1 \times 10^{-11} \text{ m}$
- D.  $2.1 \times 10^{-13} \text{ m}$
- E. none of these

ans: B

41. The frequency and wavelength of the matter wave associated with a 10-eV free electron are:

- A.  $1.5 \times 10^{34} \text{ Hz}$ ,  $3.9 \times 10^{-10} \text{ m}$
- B.  $1.5 \times 10^{34} \text{ Hz}$ ,  $1.3 \times 10^{-34} \text{ m}$
- C.  $2.4 \times 10^{15} \text{ Hz}$ ,  $1.2 \times 10^{-9} \text{ m}$
- D.  $2.4 \times 10^{15} \text{ Hz}$ ,  $3.9 \times 10^{-10} \text{ m}$
- E.  $4.8 \times 10^{15} \text{ Hz}$ ,  $1.9 \times 10^{-10} \text{ m}$

ans: D

42. If the kinetic energy of a non-relativistic free electron doubles, the frequency of its wave function changes by the factor:

- A.  $1/\sqrt{2}$
- B.  $1/2$
- C.  $1/4$
- D.  $\sqrt{2}$
- E. 2

ans: E

43. A non-relativistic free electron has kinetic energy  $K$ . If its wavelength doubles, its kinetic energy is:

- A.  $4K$
- B.  $2K$
- C. still  $K$
- D.  $K/2$
- E.  $K/4$

ans: E

44. The probability that a particle is in a given small region of space is proportional to:
- its energy
  - its momentum
  - the frequency of its wave function
  - the wavelength of its wave function
  - the square of the magnitude of its wave function
- ans: E
45.  $\psi(x)$  is the wave function for a particle moving along the  $x$  axis. The probability that the particle is in the interval from  $x = a$  to  $x = b$  is given by:
- $\psi(b) - \psi(a)$
  - $|\psi(b)|/|\psi(a)|$
  - $|\psi(b)|^2/|\psi(a)|^2$
  - $\int_a^b \psi(x) dx$
  - $\int_a^b |\psi(x)|^2 dx$
- ans: E
46. The significance of  $|\psi|^2$  is:
- probability
  - energy
  - probability density
  - energy density
  - wavelength
- ans: C
47. Maxwell's equations are to electric and magnetic fields as \_\_\_\_\_ equation is to the wave function for a particle.
- Einstein's
  - Fermi's
  - Newton's
  - Schrödinger's
  - Bohr's
- ans: D
48. A free electron in motion along the  $x$  axis has a localized wave function. The uncertainty in its momentum is decreased if:
- the wave function is made more narrow
  - the wave function is made less narrow
  - the wave function remains the same but the energy of the electron is increased
  - the wave function remains the same but the energy of the electron is decreased
  - none of the above
- ans: B

49. The uncertainty in position of an electron in a certain state is  $5 \times 10^{-10}$  m. The uncertainty in its momentum might be:
- A.  $5.0 \times 10^{-24}$  kg · m/s
  - B.  $4.0 \times 10^{-24}$  kg · m/s
  - C.  $3.0 \times 10^{-24}$  kg · m/s
  - D. all of the above
  - E. none of the above
- ans: D
50. The reflection coefficient  $R$  for a certain barrier tunneling problem is 0.80. The corresponding transmission coefficient  $T$  is:
- A. 0.80
  - B. 0.60
  - C. 0.50
  - D. 0.20
  - E. 0
- ans: D
51. An electron with energy  $E$  is incident upon a potential energy barrier of height  $E_{\text{pot}} > E$  and thickness  $L$ . The transmission coefficient  $T$ :
- A. is zero
  - B. decreases exponentially with  $L$
  - C. is proportional to  $1/L$
  - D. is proportional to  $1/L^2$
  - E. is non-zero and independent of  $L$
- ans: B
52. In order to tunnel through a potential barrier a particle must:
- A. have energy greater than the barrier height
  - B. have spin
  - C. be massive
  - D. have a wavelength longer than the barrier width
  - E. none of the above
- ans: E
53. An electron with energy  $E$  is incident on a potential energy barrier of height  $E_{\text{pot}}$  and thickness  $L$ . The probability of tunneling increases if:
- A.  $E$  decreases without any other changes
  - B.  $E_{\text{pot}}$  increases without any other changes
  - C.  $L$  decreases without any other changes
  - D.  $E$  and  $E_{\text{pot}}$  increase by the same amount
  - E.  $E$  and  $E_{\text{pot}}$  decrease by the same amount
- ans: C

54. Identical particles, each with energy  $E$ , are incident on the following four potential energy barriers:

1. barrier height =  $5E$ , barrier width =  $2L$
2. barrier height =  $10E$ , barrier width =  $L$
3. barrier height =  $17E$ , barrier width =  $L/2$
4. barrier height =  $26E$ , barrier width =  $L/3$

Rank the barriers in terms of the probability that the particles tunnel through them, from least probability to greatest probability.

- A. 1, 2, 3, 4
- B. 4, 3, 2, 1
- C. 1 and 2 tied, then 3, then 4
- D. 2, then 1 and 3 tied, then 4
- E. 3, 2, 1, 4

ans: A