

Local Anesthetic

- Local anesthetics are drugs used to reversibly depress CNS to prevent or relieve pain in specific regions of the body without loss of consciousness
- Unlike General Anesthetics, they generally don't block sense of touch, pressure or temperature or relax skeletal muscles
- Uses
 - Dentistry
 - Podiatry (treatment of disorders of the foot, ankle, and leg)
 - ENT operations
 - Surgery of skin
 - Labor pain
 - Postoperative pain
 - Skin Trauma

General Vs. Local Anesthetic

General Anesthetic	Local Anesthetic
Blocks pain in entire body	Blocks pain in specific region on body
Blocks sensation of temperature, touch and pressure	Selective to blocking pain only.
Muscle relaxation occurs	Muscle relaxation not caused
Drug intended to penetrate brain	Drug not intended to penetrate brain but act on local nerves branches
Receptor is ligand gated chlorine channel and binding site is outside of cell membrane	Receptor is voltage gated Sodium channel and binding site is inside of cell membrane

HISTORY

3000 B.C.: cocaine “Niemann”

1905: procaine “Einhorn”

1932: Tetracaine “Eisler”

1943: Lidocaine “Lofgren”

1957: Mepivacaine “Ekenstam”

1960: Prilocaine “Löfgren.”

1963: Bupivacaine “Ekenstam”

1972: Etidocaine


1996: Ropivacaine

1999: Levobupivacaine


- The first local anesthetic introduced into medical practice
Cocaine, was isolated from coca leaves by Albert Niemann in Germany in the 1860s.
- The very first clinical use of Cocaine was in 1884 by Sigmund Freud who used it to wean a patient from morphine addiction.
- Freud and his colleague Karl Kollar first noticed its anesthetic effect and introduced it to clinical ophthalmology as a topical ocular anesthetic.

How Nerve conduction occurs?

- Nerve conduction is a both electrical and chemical component
- The electrical component occurs within the axon. It involves membrane bound voltage gated ionic channels (responds to change in voltage across the membrane)
- The chemical component occurs at **synapse**. It involves *membrane bound ligand gated ionic channel* (responds to binding of a NT)





Electrical transmission of stimuli occurring through axon by movement of ion in and out



Chemical transmission of stimuli occurring at synapse through NT binding to their receptors


Concept of potential difference and its generation (-70mV) in resting cell



- When charges are separated by distance, potential **difference** is created
- The **inside** of cell has **more** potassium and **less** Sodium and **large** anionic Protein (A-) that cannot cross the lipid membrane and is thus **localized inside the cell**
- The **outside** of the cell has **more** Sodium and **less** potassium
- This difference is maintained by **Na/K** pumps that pump in **2K+** ion in cell and pump out **3Na+** ion out. Thus one extra positive charge out
- Overall, the **inside** of the cell has more **negative charge** and the **outside** has more **positive charge**
- This gives the cell a resting potential of -70mV




Steps in a action potential

1. A stimulus from a sensory cell or another neuron causes the target cell to depolarize toward the threshold potential.
2. If the threshold of excitation (-55mV) is reached, all Na^+ channels open and the membrane **depolarizes**.
3. At the peak action potential ($+30\text{mV}$) K^+ channels open and K^+ begins to leave the cell. The membrane starts to **repolarized**. At the same time, all Na^+ channels close.
4. The membrane becomes **hyperpolarized** (-90mV) as K^+ ions continue to leave the cell. The hyperpolarized membrane is in a refractory period and cannot initiate another action potential
5. The K^+ channels close and the Na^+/K^+ transporter restores the **resting potential** (-70mV).







- ① **Resting Potential**
Na⁺/K⁺ pump



- ② **Depolarisation**
Voltage-gated Na⁺ channel



- ③ **Repolarisation**
Voltage-gated K⁺ channel



- ④ **Resting Potential**
Na⁺/K⁺ pump

Changes in the resting membrane potential

Cell state	Active receptors	Potential
Resting potential	Na/K ATPase pump active	-70mV
Stimuli causes Depolarization beyond threshold potential (-55mV)	Voltage gated Na ⁺ channel open Na comes inside cell	+30mV
Repolarized state	Voltage gated Na ⁺ channel close Voltage gated K ⁺ channel open K goes outside cell	+30 to -70mV
Hyperpolarized state	Voltage gated K ⁺ channel close slowly	-90mV
Resting potential	Na/K ATPase pump active	-70mV

PROPERTIES OF AN IDEAL L.A

1. Its actions must be reversible.
2. It should be non irritating to the tissues.
3. It should have long shelf life.
4. It should be rapid in action
5. It should have a low degree of systemic toxicity.
6. It should have sufficient potency to provide complete local anesthesia.
7. It should be of sufficient duration to be advantageous.
8. It should have sufficient penetrating properties.
9. It should be either sterile or capable of being sterilized by heat without deterioration.
10. It should be stable in solution and undergo bio transformation readily within the body.
11. It should not produce any local reactions.
12. It should not produce any permanent damage.
13. It should not produce allergic reactions.
14. It should be stable in light.
15. It should be non addictive.
16. It should be combined with other agents.
17. It should have high therapeutic ratio.

CLASSIFICATION

- Method of administration.
- Classification on the basis of mode of application
- Based on duration of action
- Based on origin
- Based on chemical structure

➤ **Method of administration.**

Method of adm.	Definition	e.g. Clinical use
Surface anesthesia	application LA to the surface of the skin or mucosa	Eye surgery Dentistry, Surgery of skin
Infiltration anesthesia	injection of LA into the tissue	minor surgical and dental procedures
Nerve block	Injected of LA in the vicinity of major nerve or major branch nerve	surgical, dental, and diagnostic procedures and for pain management
epidural anesthesia	injected of LA into the epidural space where it acts primarily on the spinal nerve roots	Labor pain Postoperative pain
Spinal anesthesia	injected LA into the cerebrospinal fluid, usually at the lower back, where it acts on spinal nerve roots and part of the spinal cord.	operations below the umbilicus and Leg
Sympathetic block	injected LA around sympathetic nerves	Block some kind of pain (Cancer)

➤ Applications of LA

- ❑ **A topical anesthetic** is a local anesthetic that is used to numb the surface of a body part. They can be used to numb any area of the skin as well as the front of the eyeball, the inside of the nose, ear or throat, the anus and the genital area. Topical anesthetics are available in creams, ointments, aerosols, sprays, lotions, and jellies. Examples include benzocaine, lidocaine, oxybuprocaine,

- ❑ **Infiltration** It is the application of LA in intradermal or subcutaneous layers, where only the nerve fibers near the injected site is affected. The adequate dosage required depends on the extent of the area to be anesthetized and the expected duration of the surgical procedure.
 - Patients frequently experience pain immediately after infiltration injection of local anesthetic solutions. This response is due in part to the acidic nature of these solutions.
 - Used for postoperative pain control at incision site and suturing

❑ Epidural block

- Application of LA in the epidural space, ie just outside of the sac of cerebrospinal fluid, and thus blocking the transmission of pain signals from peripheral sensory neuron

Uses

- More effective and safe than N₂O during labor pain
- Management of back pain for hospitalized patient
- As a supplement to general anesthesia so that use of opioid analgesics can be avoided
- gynaecological surgery, orthopaedic (muscle n bone) surgery, vascular (artery n veins) surgery

❑ Spinal block

- Application of LA in the region containing cerebrospinal fluid ,which holds the spinal cord, thus blocking the transmission of pain signals from peripheral sensory neuron

Uses

- Total Hip Replacement, Total Knee Replacement, Caesarean sections, Lower limb Vascular surgery
- *limited to procedures involving regions below the upper abdomen.*
- Use in higher levels may affect the **ability to breathe** by paralyzing the intercostal respiratory muscles and the **disturb heart rate** by paralyzing cardiac nerve fibres

➤ **Based on duration of action**

Ultra short duration

- 2% lignocaine without vasoconstrictor

Short duration

- Procaine, 2% lignocaine with 0.0001% Epinephrine

Intermediate duration

- Articaine, mepivacaine, prilocaine, 2% lignocaine with 0.0002% Epinephrine

Long duration

- Bupivacaine (400-500 min), Etidocaine, 5% with lignocaine with 0.0002% Epinephrine

➤ **Based on origin**

NATURAL

- cocaine

**SYNTHETIC
NITROGENOUS
COMPOUND**


- Amino esters of PABA- procaine
- Alkyl esters of PABA- benzocaine
- Amino esters of MABA- unacaine
- Amino amides- xylocaine , bupivacaine

**SYNTHETIC NON
NITROGENOUS
COMPOUND**

- benzyl alcohol

MISCELLANEOUS DRUGS ----- Clove Oil, Phenol






➤ On the basis of chemical structure, local anaesthetics are classified as follows:



1st chemical structure classification




❖ Amino Ester

- Cocaine
- Procaine
- Tetracaine






❖ Amino Amides

- Lidocaine
- Bupivacaine
- Etidocaine





❖ Amino Ether

- Pramoxine
- Pramocaine



❖ Amino Ketones

- Dyclonine



The difference between an ester and amide local anesthetic

AMIDES

- longer lasting analgesia.
- Produce more intense analgesia.
- Rarely cause hypersensitivity reactions- no cross reactivity with ESTER LA s.
- Bind to alpha1 acid glycoprotein in plasma
- Not hydrolyzed by Plasma Cholinesterase, more slowly destroyed by liver microsomal P450 enzymes.

ESTERS


- Short duration of action
- Less intense analgesia
- Higher risk of hypersensitivity
ESTER linked LA s are rarely used.
- Hydrolyzed by Plasma Cholinesterase in blood.
- Rarely used for Infiltration anesthesia
- But useful for topical ana on mucous membranes.

Classes: The rule of “i”

- **Amides** will contain an “i” in the generic name prior to “-caine”. (i.e. lidocaine, mepivacaine, prilocaine, bupivacaine, ropivacaine, and levo-bupivacaine).
- **Ester’s** do not contain an “i” in the generic name prior to “-caine”. (i.e. procaine, chlorprocaine, cocaine, benzocaine, and tetracaine).

STRUCTURE ACTIVITY RELATIONSHIP

- All local anesthetics have an **amine** on one end to an **aromatic ring** on the other
- The **amine** end is **hydrophilic**, and the **aromatic** end is **lipophilic**
- The two groups are connected by mostly an **ester** or an **amide** group and less commonly by ether or ketones



Lipophilic group



- **Lipophilicity is important to penetrate the lipid layer and reach the binding site on the side of the cell.**

RULE 1:

The presence of electron withdrawing group in *ortho* or *para* position will decrease the lipophilicity of the drug but still **important in increases the activity for the ester group.**



Procaine is more potent if it has a e-donating amine group in the aromatic ring.

Other e-donating groups: -OH, OCH₃, CH₃





RULE 2:

Zwitterion: this increasing the activity (**for ester only**) which enhance activity due to formation of quaternary amine witch important for binding



RULE 3:

Halogens: the presence of electron withdrawing halogens in **ortho** position only can decrease duration of by making the ester more Likely for a nucleophilic attack



Chlorine in ortho group makes the carbonyl carbon more positive and more likely to be attacked by nucleophiles that causes breakdown of compound.

Nucleophile contain a lone pair of electron. they attack atoms with positive charges. The more positive atom, the better for attack

RULE 4:

For amide only, presence of di-ortho substituted group prevent breakdown of amide and thus increase it's stability in both liquid formulation and the body enzymes



Structure 1:

CH₃ group:

- Make it difficult to hydrolyze
- Stable in water and blood
- More duration of action

Structure 2:


- No protection against hydrolysis
- Unstable in water and blood
- Not enough duration of action




Note: instead of CH₃, other groups such as OCH₃ can also be used

Linker


- Linker group has short alkynene ($-\text{CH}_2-$) chain containing few carbon atom and various functional groups as Amids, ester, ether, or ketone.**




Ketone: Falcaine



Ether: Pramocaine




Amide: Lidocaine



Ester: Procaine

The increasing of the alkylene chain increase the pKa which reduces because more drug get ionized out-membrane and thus can't penetrate binding site.

Increasing CH₂ group = increased pka (increased basicity) = increased ionized form = decreased potency




All LA are basic drugs because they contain amine group

As pKa increases, the drugs become more basic and thus more ionized in the blood

In ionized form, they can't cross the lipid membrane and reach their binding site

Bridge X:



X: carbon, oxygen, nitrogen, or sulfur

The anesthetic potency decreased in the following order:


Sulphur, Oxygen, Carbon, Nitrogen.

These modification also affect duration of action and toxicity.


In general, amides (X=N) are more resistant to metabolic hydrolysis than esters (X=O). Thioesters (X=S) may cause dermatitis.

Hydrophilic group

- Usually be a secondary or tertiary amine group.
- It's important because it's believed that when they enter the cell they will accept a proton and form positively charged quaternary form (water soluble salt) that is needed for binding to voltage gated ion channel.




Procaine believed to bind to its receptor when the amine group is positively charge quaternary form




Charged Quaternary form is believed to bind to receptor


- However, *Benzocaine* has no amine portion but is still an effective topical LA
- Thus the use of Amine part could only be for proper water solubility and not directly related to proper binding




Amino Esters



Benzocaine: In *para* position of the lipophilic center there is an amino group. Lacks the basic aliphatic amine function yet has potent local anesthetic activity. Used topically.




Tetracaine: In *para* position of the lipophilic center there is an alkylamino group.




Proparacaine: There is an amino group in the *meta* position. This group will decrease lipophilicity of the molecule.


Amino Amides




Lidocaine: The o,o-dimethyl groups are required to provide suitable protection from amide hydrolysis to ensure a desirable duration of action



Mepivacaine: The o,o-dimethyl groups are required to provide suitable protection from amide hydrolysis to ensure a desirable duration of action. No carbon bridge. Cyclic amine (piperidine).















Bupivacaine: The o,o-dimethyl groups are required to provide suitable protection from amide hydrolysis to ensure a desirable duration of action. No carbon bridge. Cyclic amine (piperidine).





















Amino Ether-Pramoxine: Lipophilic group has an alkoxy Substituent. Nitrogen is in a morpholino ring

Amino Ketone-Dyclonine: Lipophilic group has an alkoxy substituent Nitrogen is in a piperidine ring

Amino Esters


	Lipophilic	Intermediate	Amine	Potency	Duration
Cocaine				2	Medium
Procaine				1	Short
2-Chloroprocaine					
Tetracaine				16	Long

Amino Amides

	Lipophilic	Intermediate	Amine	Potency	Duration
Lidocaine				4	Medium
Prilocaine				3	Medium
Etidocaine				16	Long
Mepivacaine				2	Medium
Bupivacaine				16	Long
Ropivacaine				16	Long

CLASSIFICATION AS:

1. Benzoic acid derivatives



	R	R'
Cocaine	H	
Hexylcaine	H	
Mepylcaine	H	
Isobucaine	H	
Cyclomethycaine		
Piperocaine	H	

2nd chemical structure classification

a. Benzoic acid derivatives

a.1 Derivatives of benzoic acid

a.2. Derivatives of para-amino benzoic acid

i. Freely soluble: Procaine, Amethocaine.

ii. Poorly soluble: Benzocaine, Orthocaine


b. Derivatives of acetanilide: Lignocaine, Mepivacaine, Bupivacaine, Prilocaine, Etidocaine.

c. Miscellaneous: Dimethisoquin, Dibucaine, and Dyclonine.

d. Newer drugs: Ropivacaine, Levobupivacaine

a. Benzoic acid derivatives

a.1. Benzoic acid derivatives



Hexylcaine

Meprylcaine


Isobucaine

Piperocaine

Cyclomethycaine

Name	R ₁	R ₂
Hexylcaine	-H	
Meprylcaine	-H	
Isobucaine	H	
Piperocaine	-H	
Cyclomethycaine		

a.2. p-Amino benzoic acid derivatives



Benzocaine

Tetracaine

Butacaine

Procaine


Chlorprocaine

Name	R ₁	R ₂	R ₃	R ₄	R ₅
Benzocaine	-H	-H	-H	-CH ₂ -CH ₃	-
Butamben	-H	-H	-H	-(CH ₂) ₃ CH ₃	-
Procaine	-H	-H	-H	-CH ₂ -CH ₂ -	-N(C ₂ H ₅) ₂
Chlorprocaine	-H	-H	-Cl	-CH ₂ -CH ₂ -	-N(C ₂ H ₅) ₂
Tetracaine	- ⁿ Butyl	-H	-H	-CH ₂ -CH ₂ -	-N(CH ₃) ₂
Butacaine	-H	-H	-H	-CH ₂ -CH ₂ -CH ₂ -	-N(C ₄ H ₉) ₂
Binoxinate	-H	ⁿ Butoxy	-H	-CH ₂ -CH ₂ -	-N(C ₂ H ₅) ₂
Propoxycaine	-H	-H	ⁿ Propoxy	-CH ₂ -CH ₂ -	-N(C ₂ H ₅) ₂

SAR of benzoic acid derivatives

Aryl group

- The clinically useful local anaesthetics of this series possess an aryl radical attached directly to the carbonyl group.
- Substitution of aryl group with substituents that increase the electron density of the carbonyl oxygen enhances activity.
- Favorable substituents in aryl ring include (electron-donating groups) alkoxy (propoxycaine), amino (procaine), and alkylamino (tetracaine) groups in the *para* or *ortho* positions. This homologous series increases partition coefficients with increasing number of methylene group (-CH₂-). Local anaesthetics activity peaked with the C4-, C5-, or C6-homologous: e.g., tetracaine, cyclomethycaine.
- Aryl aliphatic radicals that contain a methylene group between the aryl radical and the carbonyl group result in compounds that have not found clinical use.





SAR of benzoic acid derivatives

- Most of these local anaesthetics are tertiary amines available as HCl salts with pKa in the range of 7.5–9.0.
- Any structural modification of the local anaesthetic that causes change in pKa will have pronounced effect to reach hypothetical receptor or the binding sites.

Lipophilic “Aryl group”


1. The clinically useful local anaesthetics of this class possess an **aryl radical** that is attached directly to the carbonyl group and **are highly lipophilic**. They appear to play an **important role in the binding of local anaesthetics to the channel receptor protein**.





2. **Amino** (procaine, butacaine) **alkyl amino** (tetracaine) **alkoxyl** (cyclomethycaine) group can contribute to electron density in the aromatic ring by both resonance and inductive effects. Hence the **increase in local anaesthetic property**.

Tetracaine is more potent than procaine (40–50 times). Although the butyl group present in it increases lipid solubility


cyclomethycaine



Amino




alkyl amino




3. Any substitution that enhances zwitterion formation will be more potent. Hence *m*-position substitution decreases the activity.


Tetracaine is more potent than procaine (40–50 times). Although the potentiation is partly due to electron releasing property of the *n*-butyl group via inductive effect, which intend to increase the formation of the **Zwitterion**.



4. Structural modification leads to change in physical and chemical properties. Electron withdrawing substituents in *ortho* or *para* or at both the positions leads to an increase of its local anaesthetic property.





Chlorprocaine



Procaine



Intermediate “Bridge X”

- The bridge X may be carbon, oxygen, nitrogen, or sulphur.
- In an procaine series, anaesthetic potency decreased in the following order: sulphur, oxygen, carbon, nitrogen.
- These modifications also affect duration of action and toxicity. In general, amides (X=N) are more resistant to metabolic hydrolysis than esters (X=O). Thioesters (X=S) may cause dermatitis.
- In procaine-like analogues, branching (especially at the alpha carbon) will increase duration of action. This effect is not seen in the lidocaine series.
- Increasing the chain length will increase potency but will also increase toxicity.






Hydrophilic portion “Aminoalkyl”

- The aminoalkyl group is not necessary for local anaesthetic activity, but it is used to form water-soluble salts (HCl salts).
- Tertiary amines result in more useful agents. The secondary amines appear to be of longer activity, but they are more irritating; primary amines are not very active and cause irritation.
- The tertiary amino group may be diethylamino, piperidine, or pyrrolidino, leading to the products that exhibit essentially the same degree of activity.
- The more hydrophilic morpholino group usually leads to diminished potency.
- Some analogues have no amino group at all, such as benzocaine. They are active but have poor water solubility.



b. Anilide derivatives





Name	R ₁	R ₂
Lidocaine/Lignocane	-CH ₃	-CH ₂ -N $\begin{matrix} / \text{C}_2\text{H}_5 \\ \backslash \text{C}_2\text{H}_5 \end{matrix}$
Mepivacaine	-CH ₃	
Prilocaine	-H	$\begin{matrix} \text{CH}_3 \\ \\ \text{---C---NH---}(\text{CH}_2)_2\text{---CH}_3 \\ \\ \text{H} \end{matrix}$
Bupivacaine	-CH ₃	
Etidocaine	-CH ₃	$\begin{matrix} \text{C}_2\text{H}_5 \\ \\ \text{---C---N---CH}_2\text{CH}_2\text{CH}_3 \\ \\ \text{H} \\ \text{C}_2\text{H}_5 \end{matrix}$

SAR of Anilides

Aryl group

- The clinically useful local anaesthetics of this type possess a phenyl group attached to the sp^2 carbon atom through a nitrogen bridge.
- Placement of substituents on the phenyl ring with a methyl group in the 2 (or) 2 and 6-position enhances the activity. In addition, the methyl substituent “ as lidocaine “ provides steric hindrance to hydrolysis of the amide bond and enhances the coefficient of distribution.
- Any substitution on the aryl ring that enhances zwitterion formation will be more potent.





Substituent X


- X may be carbon, oxygen, or nitrogen. Among them Lidocaine series (X=O) has provided more useful products.

Aminoalkyl group

- The amino function has the capacity for salt formation and is considered the hydrophilic portion of the molecules.
- Tertiary amines (diethylamine, piperidines) are more useful because the primary and secondary amines are more irritating to tissue.


c. Miscellaneous

Phenacaine




Structurally, it is related to anilides in that the aromatic ring is attached to a sp² carbon through a nitrogen bridge. It is one of the oldest synthetic local anaesthetics. It is used mainly for producing local anaesthesia of the eye.

Diperodon




Soluble in water, and potent surface anaesthetic; used primarily for anus. Very toxic in nature.

Pramoxine HCl (Traonaolene)




It is a surface anaesthetic, which possesses very low degree of toxicity and sensitization. It is applied locally as 1% solution in rectal surgery, itching, and minor burns.

Dyclonine (Dyclone)




Containing lozenges are used to relieve minor sore throat and mouth discomfort. It is used to anesthetize mucous membranes of mouth, trachea, and urethra prior to various endoscopic procedures.

Dibucaine (Nupercaine)



It is several times more potent than procaine when injected subcutaneously and five times more toxic than cocaine, when injected intravenously. It is the most potent toxic and long-acting local anaesthetics used as infiltration, surface and spinal anaesthesia.


Dimethisoquin (Synonym: Quinisocaine, Quotane)




It is a surface anaesthetic used as an ointment or lotion for relief from irritation, itching, pain, or burning.

d. Newer drugs: Ropivacaine, Levobupivacaine

Newer local anaesthetics were introduced with the goal of reducing local tissue irritation, minimizing systemic cardiac and central nervous system (CNS) toxicity, and achieving a faster onset and longer duration of action.




Ropivacaine



Levobupivacaine


PROCAINE

- Procaine is a local anesthetic drug of the ester group
- effective parental but are relatively weak when applied topically
- slow onset (4-5 min), short duration, pKa=8.8
- It has the advantage of lacking of local irritation, minimal systemic toxicity, longer duration of action, and low cost.
- It can be effectively used for causing anaesthesia by infiltration, nerve block, epidural block, or spinal anaesthesia.
- is metabolized by the plasma enzymes to form para-amino benzoic acid (PABA) which is causes allergic effect
- MOA – blocks pain by depressing CNS by antagonizing voltage gated Na⁺ channel thus inhibiting generation of action potential




Synthesis


Route I. From: *p*-Amino benzoic acid: direct reaction of the 4-aminobenzoic acid with 2-diethylamino-ethanol in the presence of sodium ethoxide.



Route II. From: oxidizing 4-nitrotoluene to 4-nitrobenzoic acid, which is further reacted with thionyl chloride




Tetracaine




It is an all-purpose local anaesthetic drug used frequently in surface, infiltration block, caudal, and spinal anaesthesia. It is reported to be 10 times more toxic and potent than procaine. Its duration of action is twice than that of procaine.

Propoxycaine (Blockhain)



Its local anaesthetic potency is reported to be 7 or 8 times more than that of procaine. It is a structural isomer of proparacaine, and is less toxic with slightly lower potency than proparacaine. It is mainly used for infiltration and nerve block anaesthesia.


Proparacaine



An effective ester-type surface anaesthetic with potency about equal to that of tetracaine. It is a useful anaesthetic in ophthalmology and induces little or no initial irritation. It is useful for most ocular procedures that require topical anaesthesia.


Chloroprocaine

Is a very short-acting, amino ester-type local anesthetic used to provide regional anesthesia by infiltration as well as by peripheral and central nerve block, including lumbar and caudal epidural blocks. The presence of a chlorine atom *ortho* to the carbonyl of the ester function increases its lipophilicity and its rate of hydrolysis by plasma cholinesterase at least threefold compared to procaine and benzocaine.




Benzocaine

Like most amino ester-type local anesthetics, it is easily hydrolyzed by plasma cholinesterase. However, because of its low pK_a, it is un-ionized under most physiologic conditions and, therefore, can only bind to the lipid site in the sodium channel




LIDOCAINE

- Most commonly used potent amide type local anesthetic for both parenteral and topical use
- has a rapid onset of action (Intravenous - 45 to 90 seconds).
- It is a potent local anaesthetic. It is reported to be twice as active as procaine hydrochloride in the same concentrations
- Di-ortho methyl groups make the amide group resistant to hydrolysis thus it has moderate duration of action (1-2 hrs)
- Produces eutectic mixture with prilocaine
- MOA – blocks pain by depressing CNS by antagonizing voltage gated Na⁺ channel thus inhibiting generation of action potential




Lidocaine synthesis




Lidocaine is synthesised from 2,6-dimethylaniline upon reaction with chloroacetyl chloride, which gives α -chloro-2,6-dimethylacetanilide, and its subsequent reaction with diethylamine affords lidocaine.

Metabolism of Lidocaine

Undergoes *N*-de-ethylation to yield mono-ethylglycinexylide followed by amidase action to *N*-ethyl glycine and 2, 6-dimethylaniline.




Mepivacaine (Polocaine)




The duration of action is significantly longer than that of lidocaine, even without adrenaline. It is of particular importance in subjects showing contraindication to adrenaline. It is a local anaesthetic used for infiltration, peridural, nerve block, and caudal anaesthesia.

Prilocaine (Citanest hydrochloride)



Its duration of action is in between the shorter-acting lidocaine and longer-acting mepivacaine. The solution of prilocaine HCl is specifically used for such patients who cannot tolerate vasopressor agents, patients having cardiovascular disorders, diabetes, hypertension, and thyrotoxicosis.

Bupivacaine (Marcaine)



It is a long-acting local anaesthetic of the amide type, similar to mepivacaine and lidocaine, but about four times more potent. The effect of bupivacaine last longer than lidocaine hydrochloride. It is longacting local anaesthetic mainly employed for regional nerve block.

Ropivacaine

- *S*-Ropivacaine hydrochloride is the first optically active, **amino amide** type local anesthetic marketed in recent years. It combines the anesthetic **potency and long duration of action** of **bupivacaine** with a **side effect profile intermediate between those of bupivacaine and lidocaine**.
- Although ropivacaine has a *pK*_a nearly identical to that of bupivacaine, it is two to threefold **less lipid soluble and has a smaller volume of distribution, a greater clearance, and a shorter elimination half-life than bupivacaine in humans**.
- The metabolism of ropivacaine in humans is mediated by **hepatic** CYP1A2 and, to a minor extent, by CYP3A4. The major metabolite is **3-hydroxyropivacaine**, and the minor metabolite is *S*-2',6'-pipecoloxylidide (an *N*-dealkylated product).

