


# 7 | Carbohydrates and Glycobiology

© 2017 W. H. Freeman and Company



# CHAPTER 7

## Carbohydrates and Glycobiology

---

### *Learning goals:*

- Structures and names of monosaccharides
- Open-chain and ring forms of monosaccharides
- Structures and properties of disaccharides
- Biological function of polysaccharides
- Biological function of glycoconjugates

# Carbohydrates

---

- Named so because many have formula  $C_n(H_2O)_n$
- Produced from  $CO_2$  and  $H_2O$  via **photosynthesis** in plants
- Range from as small as glyceraldehyde ( $M_w = 90$  g/mol) to as large as amylopectin ( $M_w > 200,000,000$  g/mol)
- Fulfill a variety of functions, including:
  - energy source and energy storage
  - structural component of cell walls and exoskeletons
  - informational molecules in cell-cell signaling
- Can be covalently linked with proteins and lipids

# Carbohydrates

---

- **Monosaccharides** (simple sugars) – one polyhydroxy aldehyde or ketone unit (glucose)
- **Disaccharides** – two monosaccharide units joined together by a glycosidic linkage (sucrose)
- **Oligosaccharides** – few monosaccharide units joined together (in cells, most oligosaccharides are joined to nonsugar molecule)
- **Polysaccharides** – sugar polymers consisting of >20 monosaccharide units

# Carbohydrates

---

- Basic nomenclature:
  - number of carbon atoms in the carbohydrate + -ose
  - example: three carbons = triose
- Common functional groups:
  - All carbohydrates initially had a carbonyl functional group.
  - aldehydes = aldose
  - ketones = ketose


# Monosaccharide Carbon backbone

---


- 3 C → triose
- 4 C → tetrose
- 5 C → pentose
- 6 C → hexose
- 7 C → heptose

# Aldoses and Ketoses

- An **aldose** contains an **aldehyde** functional group
- A **ketose** contains a **ketone** functional group



**Glyceraldehyde,  
an aldotriose**



**Dihydroxyacetone,  
a ketotriose**

# Stereoisomers

---


- **All monosaccharides** (except dihydroxyacetone) are chiral compounds, with **at least** one chiral carbon
- A molecule with  $n$  chiral centers can have  $2^n$  stereoisomers
- E.g. glyceraldehyde has  $2^1 = 2$   
aldohexoses have  $2^4 = 16$  stereoisomers




# Carbohydrates Can Be Stereoisomers

---

- **Enantiomers**
  - Stereoisomers that are non-superimposable mirror images
- In sugars that contain many chiral centers, only the one that is **most distant from the carbonyl carbon** is designated as D (right) or L (left)
- **D and L isomers of a sugar are enantiomers**
  - For example, L and D glucose have the same water solubility
- Most hexoses in living organisms are D stereoisomers
- Some simple sugars occur in the L-form, such as L-arabinose




L-Glyceraldehyde



D-Glyceraldehyde

Non-  
superimposable  
mirror images




L-Glucose



Non-


superimposable  
mirror images



D-Glucose


# Carbohydrates Can Be Stereoisomers

- **Epimers** are two sugars that differ only in the configuration around **one carbon atom**
- Epimers are NOT mirror images, and therefore are NOT enantiomers.
- Epimers are diastereomers; diastereomers have different physical properties (i.e., water solubility, melting temp).
  - example: D-Threose is the C-2 epimer of D-erythrose.
  - Both are D sugars because they both have the same orientation around the last chiral carbon in the chain.



# Epimers

- D-Mannose and D-galactose are both epimers of D-glucose.
- D-Mannose and D-galactose vary at more than one chiral center and are diastereomers, but not epimers.




# Structures to Know

---


- **Glyceraldehyde & dihydroxyacetone** are the standard three-carbon sugars
- **Ribose** is the standard five-carbon sugar
- **Glucose** is the standard six-carbon sugar
- **Galactose** is an **epimer** of glucose
- **Mannose** is an **epimer** of glucose
- **Fructose** is the **ketose** form of glucose

*Aldotriose*




D-Glyceraldehyde

*Aldotetroses*




D-Erythrose




D-Threose


*Aldopentoses*




D-Ribose



D-Arabinose




D-Xylose




D-Lyxose


*Aldohexoses*




D-Allose




D-Altrose




D-Glucose




D-Mannose




D-Gulose



D-Idose




D-Galactose



D-Talose


*Ketotriose*




Dihydroxyacetone



*Ketotetrose*



D-Erythrulose





# Common Monosaccharides Have Cyclic Structures

---

- In solution, monosaccharides with 4 or more carbons form cyclic compounds (ring structures)
- Carbonyl C is attacked by the hydroxyl O forming a covalent bond

# Hemiacetals and Hemiketals

- Aldehyde and ketone carbons are **electrophilic**
- Alcohol oxygen atom is **nucleophilic**
- When **aldehydes** are attacked by alcohols, **hemiacetals** form
- When **ketones** are attacked by alcohols, **hemiketals** form
- These reactions form the basis of cyclization of sugars.









Figure 7-5  
Lehninger Principles of Biochemistry, Sixth Edition  
© 2013 W. H. Freeman and Company

*When the second alcohol is part of another sugar molecule, the bond produced is a **glycosidic bond**.*

# Cyclization of Monosaccharides


---

- Pentoses and hexoses readily undergo intramolecular ring formation
- The former carbonyl carbon becomes a new chiral center, called the **anomeric carbon**
- The former carbonyl oxygen becomes a hydroxyl group; the position of this group determines if the anomer is  $\alpha$  or  $\beta$
- If the *hydroxyl group* is on the **opposite side** (**trans**) of the ring as the  $CH_2OH$  moiety the configuration is  $\alpha$
- If the *hydroxyl group* is on the **same side** (**cis**) of the ring as the  $CH_2OH$  moiety, the configuration is  $\beta$



$\alpha$ -D-Glucopyranose  
(Haworth projection)

or




$\beta$ -D-Glucopyranose  
(Haworth projection)


**Mutarotation** – the interconversion of  $\alpha$  and  $\beta$  anomers.

# Pyranoses and Furanoses


- Six-membered oxygen-containing rings are called **pyranoses**
- Five-membered oxygen-containing rings are called **furanoses**
- The **anomeric** carbon is usually drawn on the **right side**
- Cyclic sugar structures are more accurately represented in **Haworth perspective formulas**




**D-Glucose**  
**Fischer projection**




**α-D-Glucopyranose**  
**Haworth perspective**




**$\alpha$ -D-Glucopyranose**




**$\alpha$ -D-Fructofuranose**




**$\beta$ -D-Glucopyranose**



**$\beta$ -D-Fructofuranose**



**Pyran**




**Furan**

# Converting linear to cyclic structures


---

- For any linear hexose:
  1. Draw 6-membered ring (O at the upper right)
  2. Number C's clockwise (start with anomeric)
  3. Insert the  $\text{-OH}$ 
    - a. if  $\text{-OH}$  is on the **right**, draw it pointing **down**
    - b. if  $\text{-OH}$  is on the **left**, draw it pointing **up**
  4. Insert the terminal  $\text{-CH}_2\text{OH}$ 
    - a. if the sugar is the **D-isomer**, draw it **upwards**
    - b. if it is the **L-isomer**, draw it **downwards**
  5. Draw the  $\text{-OH}$  on the anomeric C
    - a. if it faces the **same side** as the  $\text{-CH}_2\text{OH}$ , it's  $\beta$
    - b. if it faces the **opposite side**, it's  $\alpha$

# Example




$\beta$ -D-glucose  
(C<sup>1</sup>-OH points up,  
*cis* to C<sup>6</sup>H<sub>2</sub>OH)






$\alpha$ -L-glucose  
(C<sup>1</sup>-OH points up,  
*trans* to C<sup>6</sup>H<sub>2</sub>OH)







D-Ribose  
(Fischer projection)




$\alpha$ -D-Ribopyranose  
(Haworth projection)



$\beta$ -D-Ribopyranose  
(Haworth projection)





$\alpha$ -D-Ribofuranose  
(Haworth projection)



$\beta$ -D-Ribofuranose  
(Haworth projection)

# The ring is not planar

- In solution pyranose ring is not planar but it can assume two chair conformations
- Boat conformation is possible but very rare
- Multiple “chair” conformations are possible but require energy for interconversion (~46 kJ/mole)



$\alpha$ -D-Glucopyranose (b)






Figure 7-8  
Lehninger Principles of Biochemistry, Sixth Edition  
© 2013 W. H. Freeman and Company

(a)




Haworth projection

(b)




C<sub>2</sub>-endo envelope conformation

(c)



C<sub>3</sub>-endo envelope conformation

(d)



Twist conformation

# Chain-Ring Equilibrium and Reducing Sugars

---

- The ring forms exist in equilibrium with the open-chain forms
- Reducing sugars have a free anomeric carbon
- Aldehyde can reduce  $\text{Cu}^{2+}$  to  $\text{Cu}^+$  (Fehling's test)
- Aldehyde can reduce  $\text{Ag}^+$  to  $\text{Ag}^0$  (Tollens' test)
- Allows detection of reducing sugars, such as glucose (by measuring the amount of oxidizing agent reduced by a sugar solution, the sugar concentration can be estimated)

The cuprous ion ( $\text{Cu}^+$ ) produced forms a red cuprous oxide precipitate.




Figure 7-10







Lehninger Principles of Biochemistry, Fifth Edition

© 2008 W. H. Freeman and Company

# Blood glucose measurements in the diagnosis and treatment of diabetes

---

- Untreated diabetes has several long-term consequences: kidney failure, cardiovascular disease, blindness, impaired wound healing, etc.
- Average  $[\text{glc}]_{\text{blood}}$  over days can be measured because of a nonenzymatic reaction between  $\text{glc}$  and primary amino groups in Hb
- The amount of glycated Hb (GHb) reflects the average  $[\text{glc}]_{\text{blood}}$  over the circulating lifetime of RBC (~120 days)
- Normal levels ~5% of Hb is GHb
- Diabetic people may have it as high as 13%
- Advanced glycation endproducts (AGEs) contribute to the long term problems associated with diabetes




**Damage to kidneys, retinas, cardiovascular system**


**Box 7-1 figure 1**  
*Lehninger Principles of Biochemistry, Sixth Edition*  
 © 2013 W. H. Freeman and Company

# Organisms contain many hexose derivatives


## Glucose family




$\beta$ -D-Glucose




$\beta$ -D-Glucose 6-phosphate




$\beta$ -D-Glucosamine




*N*-Acetyl- $\beta$ -D-glucosamine



Muramic acid



*N*-Acetylmuramic acid



## Amino sugars




$\beta$ -D-Galactosamine




$\beta$ -D-Mannosamine


## Deoxy sugars




$\beta$ -L-Fucose




$\alpha$ -L-Rhamnose



$\beta$ -D-Glucuronate




D-Gluconate



D-Glucono- $\delta$ -lactone

## Acidic sugars



*N*-Acetylneuraminic acid  
(a sialic acid)




Figure 7-9

Lehninger Principles of Biochemistry, Sixth Edition


© 2013 W. H. Freeman and Company



# The Glycosidic Bond

---

- Two sugar molecules can be joined via a **glycosidic bond** between an anomeric carbon and a hydroxyl carbon
- The resulting compound is a **glycoside**
- The **glycosidic bond** (an **acetal**) between monomers is less reactive than the hemiacetal at the second monomer
  - Second monomer, with the hemiacetal, is reducing
  - Anomeric carbon involved in the glycosidic linkage is nonreducing
- The disaccharide formed upon condensation of two glucose molecules via 1 → 4 bond is called **maltose**



**Figure 7-10**  
*Lehninger Principles of Biochemistry, Sixth Edition*  
 © 2013 W. H. Freeman and Company

# Nonreducing Disaccharides

---


- Two sugar molecules can be also joined via a **glycosidic bond** between two anomeric carbons
- The product has two acetal groups and no hemiacetals
- There are **no reducing ends**, this is a nonreducing sugar
- Trehalose is a constituent of hemolymph of insects
  - Provides protection from drying

# Naming disaccharides

---


1. Find the nonreducing end
2. Give the configuration ( $\alpha$  or  $\beta$ ) at the anomeric C joining the first monosaccharide to the other
3. Name the nonreducing residue (use “furano” for 5-membered rings or “pyrano” for 6-membered rings)
4. Add the glycosidic bond in parenthesis (from which C to which C)
5. Name the second residue
6. If there are more residues, repeat step 2

\* Abbreviations can be used (Glc, Fru, Gal, Man, GlcN, etc.)




**Lactose ( $\beta$  form)**

$\beta$ -D-galactopyranosyl-(1 $\rightarrow$ 4)- $\beta$ -D-glucopyranose  
Gal( $\beta$ 1 $\rightarrow$ 4)Glc




$\beta$  anomer of cellobiose

( $\beta$ -D-Glucopyranosyl-(1 $\rightarrow$ 4)- $\beta$ -D-glucopyranose)



**Sucrose**

$\beta$ -D-fructofuranosyl  $\alpha$ -D-glucopyranoside  
Fru(2 $\beta$  $\leftrightarrow$  $\alpha$ 1)Glc  $\equiv$  Glc( $\alpha$ 1 $\leftrightarrow$ 2 $\beta$ )Fru



**Trehalose**


$\alpha$ -D-glucopyranosyl  $\alpha$ -D-glucopyranoside  
Glc( $\alpha$ 1 $\leftrightarrow$ 1 $\alpha$ )Glc

# Sweet tooth? Anyone?



Box 7-2 Figure 1  
Lehninger Principles of Biochemistry, Sixth Edition  
© 2013 W. H. Freeman and Company


- Most people like sweets
- Sweetness is one of 5 tastes humans can taste
- Due to receptors on gustatory cells on the surface of tongues... T1R2 and T1R3
- Stevioside (few hundred times sweeter than sugar). Brazzein protein ~17000x sweeter
- Binding of molecules to these receptors, signals are transduced to give the sweet taste



Box 7-2 figure 2  
Lehninger Principles of Biochemistry, Sixth Edition  
© 2013 W. H. Freeman and Company

to give the sweet taste

# L,L-Aspartame is sweet; D,L-Aspartame is bitter



**Box 7-2 figure 3**

*Lehninger Principles of Biochemistry, Sixth Edition*

© 2013 W. H. Freeman and Company

# Polysaccharides


---

- Natural carbohydrates are usually found as polymers
- These polysaccharides can be
  - homopolysaccharides
  - heteropolysaccharides
  - linear
  - branched
- Polysaccharides do not have a defined molecular weight.
  - This is in contrast to proteins because unlike proteins, no template is used to make polysaccharides



## Homopolysaccharides


Unbranched      Branched



## Heteropolysaccharides

Two  
monomer  
types,  
unbranched

Multiple  
monomer  
types,  
branched



Can be:

- 1) Storage forms of monosaccharides, used for fuel (starch and glycogen)
- 2) Structural elements in plant cell walls and animal exoskeletons (cellulose and chitin)

Provide extracellular support for organisms of all kingdoms,

E.g.

- 1) Bacterial cell envelope is composed in part of a heteropolysaccharide
- 2) Extracellular matrix in animal cells

**Table 8.2 Structures of some common polysaccharides**

<b>Polysaccharide<sup>a</sup></b>	<b>Component(s)<sup>b</sup></b>	<b>Linkage(s)</b>
<b>Storage homoglycans</b>		
Starch		
Amylose	Glc	$\alpha$ -(1 → 4)
Amylopectin	Glc	$\alpha$ -(1 → 4), $\alpha$ -(1 → 6) (branches)
Glycogen	Glc	$\alpha$ -(1 → 4), $\alpha$ -(1 → 6) (branches)
<b>Structural homoglycans</b>		
Cellulose	Glc	$\beta$ (1 → 4)
Chitin	GlcNAc	$\beta$ (1 → 4)
<b>Heteroglycans</b>		
Glycosaminoglycans	Disaccharides (amino sugars, sugar acids)	Various
Hyaluronic acid	GlcUA and GlcNAc	$\beta$ (1 → 3), $\beta$ (1 → 4)

<sup>a</sup>Polysaccharides are unbranched unless otherwise indicated.

<sup>b</sup>Glc, Glucose; GlcNAc, *N*-acetylglucosamine; GlcUA, *D*-glucuronate.

TABLE 7-2

## Structures and Roles of Some Polysaccharides

Primer	Type <sup>a</sup>	Repeating unit <sup>b</sup>	Size (number of monosaccharide units)	Roles/significance
Starch				Energy storage: in plants
Amylose	Homo-	( $\alpha 1 \rightarrow 4$ ) Glc, linear	50–5,000	
Amylopectin	Homo-	( $\alpha 1 \rightarrow 4$ ) Glc, with ( $\alpha 1 \rightarrow 6$ ) Glc branches every 24–30 residues	Up to $10^6$	
Glycogen	Homo-	( $\alpha 1 \rightarrow 4$ ) Glc, with ( $\alpha 1 \rightarrow 6$ ) Glc branches every 8–12 residues	Up to 50,000	Energy storage: in bacteria and animal cells
Cellulose	Homo-	( $\beta 1 \rightarrow 4$ ) Glc	Up to 15,000	Structural: in plants, gives rigidity and strength to cell walls
Chitin	Homo-	( $\beta 1 \rightarrow 4$ ) GlcNAc	Very large	Structural: in insects, spiders, crustaceans, gives rigidity and strength to exoskeletons
Dextran	Homo-	( $\alpha 1 \rightarrow 6$ ) Glc, with ( $\alpha 1 \rightarrow 3$ ) branches	Wide range	Structural: in bacteria, extracellular adhesive
Peptidoglycan	Hetero-; peptides attached	4)Mur2Ac( $\beta 1 \rightarrow 4$ ) GlcNAc( $\beta 1$ )	Very large	Structural: in bacteria, gives rigidity and strength to cell envelope
Agarose	Hetero-	3)D-Gal ( $\beta 1 \rightarrow 4$ )3,6- anhydro-L-Gal( $\alpha 1$ )	1,000	Structural: in algae, cell wall material
Hyaluronan (a glycosaminoglycan)	Hetero-; acidic	4)GlcA ( $\beta 1 \rightarrow 3$ ) GlcNAc( $\beta 1$ )	Up to 100,000	Structural: in vertebrates, extracellular matrix of skin and connective tissue; viscosity and lubrication in joints



<sup>a</sup>Each polymer is classified as a homopolysaccharide (homo-) or heteropolysaccharide (hetero-).

<sup>b</sup>The abbreviated names for the peptidoglycan, agarose, and hyaluronan repeating units indicate that the polymer contains repeats of this disaccharide unit. For example, in peptidoglycan, the GlcNAc of one disaccharide unit is ( $\beta 1 \rightarrow 4$ )-linked to the first residue of the next disaccharide unit.

# Glycogen

---

- Glycogen is a branched homopolysaccharide of **glucose**
  - Glucose monomers form ( $\alpha 1 \rightarrow 4$ ) linked chains
  - **Branch-points** with ( $\alpha 1 \rightarrow 6$ ) linkers every 8–12 residues
  - Molecular weight reaches several millions
  - Functions as the main **storage polysaccharide in animals**




# Starch


---

- Starch is a **mixture** of two homopolysaccharides of **glucose**
  - **Amylose** is an unbranched polymer of ( $\alpha 1 \rightarrow 4$ ) linked residues
  - **Amylopectin** is branched like glycogen but the branch-points with ( $\alpha 1 \rightarrow 6$ ) linkers occur every 24–30 residues
    - Molecular weight of amylopectin is up to 200 million
- Starch is the main **storage polysaccharide in plants**

# Glycosidic Linkages in Glycogen and Starch




**Figure 7-13a**  
*Lehninger Principles of Biochemistry, Sixth Edition*  
© 2013 W. H. Freeman and Company



**Figure 7-13b**  
*Lehninger Principles of Biochemistry, Sixth Edition*  
© 2013 W. H. Freeman and Company

# Mixture of Amylose and Amylopectin in Starch



**Figure 7-13c**  
*Lehninger Principles of Biochemistry, Sixth Edition*  
© 2013 W. H. Freeman and Company

*In starch granules, strands of amylopectin (blue) form double-helical structures with each other or with amylose strands (black). Glucose residues at the nonreducing ends of the outer branches are removed enzymatically during the mobilization of starch for energy production.*

*Glycogen has a similar structure but is more highly branched and more compact.*



# Metabolism of Glycogen and Starch

---

- Glycogen and starch often form **granules** in cells
- Granules contain enzymes that synthesize and degrade these polymers
- Glycogen and amylopectin have one reducing end but **many nonreducing ends** (a polymer with  $n$  branches has  $n + 1$  nonreducing ends and only 1 reducing end)
- Enzymatic processing occurs simultaneously in many nonreducing ends

# Dextrans

---


- Bacterial and yeast polysaccharides
- ( $\alpha 1 \rightarrow 6$ )-linked poly-D-glucose
- All units have ( $\alpha 1 \rightarrow 3$ ) branches
- Some have also ( $\alpha 1 \rightarrow 2$ ) or ( $\alpha 1 \rightarrow 4$ ) branches
- Dental plaque (formed by bacteria on the surface of teeth) is rich in dextrans

# Cellulose


---

- Cellulose is an **unbranched homopolysaccharide** of glucose
  - Glucose monomers form ( $\beta 1 \rightarrow 4$ ) linked chains
  - **Hydrogen bonds** form between adjacent monomers
  - Additional H-bonds between chains
  - Structure is now tough and water-insoluble
  - Most abundant polysaccharide in nature
  - Cotton is nearly pure fibrous cellulose


(a)



(b)



# Hydrogen Bonding in Cellulose



**$(\beta 1 \rightarrow 4)$ -linked D-glucose units**

**Figure 7-14**

*Lehninger Principles of Biochemistry, Sixth Edition*

© 2013 W. H. Freeman and Company

# Cellulose Metabolism

---

- The fibrous structure and water-insolubility make cellulose a difficult substrate to act on
- Fungi, bacteria, and protozoa secrete **cellulase**, which allows them to use wood as source of glucose
- Most animals cannot use cellulose as a fuel source because they lack the enzyme to hydrolyze ( $\beta 1 \rightarrow 4$ ) linkages
- **Ruminants and termites** live symbiotically with microorganisms that produces cellulase
- Cellulases hold promise in the fermentation of biomass into biofuels

A wood fungus growing on an oak log



**Figure 7-16**

*Lehninger Principles of Biochemistry, Fifth Edition*

© 2008 W. H. Freeman and Company

# Chitin Is a Homopolysaccharide

- Chitin is a linear homopolysaccharide of *N*-acetylglucosamine.
  - *N*-acetylglucosamine monomers form ( $\beta 1 \rightarrow 4$ )-linked chains.
  - forms extended fibers that are similar to those of cellulose
  - hard, insoluble, cannot be digested by vertebrates
  - structure is tough but flexible, and water insoluble
  - found in cell walls in mushrooms and in exoskeletons of insects, spiders, crabs, and other arthropods




Figure 7-16a

Lehninger Principles of Biochemistry, Seventh Edition

© 2017 W. H. Freeman and Company





**Figure 7-16b**

*Lehninger Principles of Biochemistry, Sixth Edition*


© 2013 W. H. Freeman and Company

# Agar and Agarose

---

- Agar is a complex mixture of complex heteropolysaccharides composed of **agarose** and **agarpectin** (containing **modified galactose** units)
- Agar serves as a component of cell wall in some seaweeds
- Agar solutions form gels that are commonly used in the laboratory as a surface for **growing bacteria**
- Agar is also used for capsules in which some drugs and vitamins are packaged
- Agarose solutions form gels that are commonly used in the laboratory for **separation DNA** by electrophoresis

# Agar and Agarose



**Agarose**

**3)D-Gal( $\beta$ 1  $\rightarrow$ 4)3,6-anhydro-L-Gal<sup>2S</sup>( $\alpha$ 1 repeating units)**

# Glycosaminoglycans

---

- Linear polymers of **repeating disaccharide units**
- One monomer is either
  - N-acetyl-glucosamine or
  - N-acetyl-galactosamine
- Negatively charged
  - **Uronic acids** (C6 oxidation)
  - **Sulfate esters**
- Extended hydrated molecule
  - Minimizes charge repulsion
- Forms meshwork with fibrous proteins to form **extracellular matrix**
  - Connective tissue
  - Lubrication of joints


## Glycosaminoglycan

## Repeating disaccharide

Number of  
disaccharides  
per chain

Hyaluronate

~50,000



Chondroitin  
4-sulfate

20 – 60




Figure 7-22 part 1a

*Lehninger Principles of Biochemistry, Sixth Edition*


© 2013 W. H. Freeman and Company

# Glycosaminoglycan

# Repeating disaccharide

Number of  
disaccharides  
per chain


Keratan  
sulfate  
~25



Gal

GlcNAc6S

Heparin  
15-90



IdoA2S

GlcNS3S6S

Figure 7-22 part 1b


Lehninger Principles of Biochemistry, Sixth Edition

© 2013 W. H. Freeman and Company

# Heparin and Heparan Sulfate

---

- Heparin is linear polymer, 3–40 kDa.
- Heparan sulfate is heparin-like polysaccharide but attached to proteins.
- Highest negative-charge density biomolecules
- **Prevent blood clotting** by activating protease inhibitor antithrombin
- Binding to various cells **regulates development and formation of blood vessels**.
- Can also bind to viruses and bacteria and **decrease their virulence**



*The affinity of antithrombin for thrombin is three orders of magnitude greater in the presence of heparin than in its absence.*

**Figure 7-27**

Lehninger Principles of Biochemistry, Seventh Edition  
© 2017 W. H. Freeman and Company




# Glycoconjugates: Glycoprotein


---

- A protein with small oligosaccharides attached
  - Carbohydrate attached via its **anomeric carbon**
  - About half of mammalian proteins are glycoproteins
  - Carbohydrates play role in **protein-protein recognition**
  - Only some bacteria glycosylate few of their proteins
  - Viral proteins heavily glycosylated; helps **evade the immune system**


(a) O-linked



Examples:



(b) N-linked



Examples:





Figure 7-30  
Lehninger Principles of Biochemistry, Sixth Edition  
© 2013 W. H. Freeman and Company

# Glycoconjugates: Glycolipids

---

- A lipid with covalently bound oligosaccharide
  - Parts of plant and animal cell membranes
  - In vertebrates, ganglioside carbohydrate composition determines **blood groups**
  - In gram-negative bacteria, **lipopolysaccharides** cover the peptidoglycan layer

H antigen

Fuc  $\alpha$ -(1  $\rightarrow$  2)

Gal  $\beta$ -(1  $\rightarrow$  3)- GlcNAc  $\beta$ ...

A enzyme

B enzyme

Fuc  $\alpha$ -(1  $\rightarrow$  2)

Gal  $\beta$ -(1  $\rightarrow$  3)- GlcNAc  $\beta$ ...

GalNAc  $\alpha$ -(1  $\rightarrow$  3)




A antigen

Fuc  $\alpha$ -(1  $\rightarrow$  2)

Gal  $\beta$ -(1  $\rightarrow$  3)- GlcNAc  $\beta$ ...

Gal  $\alpha$ -(1  $\rightarrow$  3)

B antigen



# Glycoconjugates: Proteoglycans

---

- Sulfated **glucoseaminoglycans** attached to a large rod-shaped protein in cell membrane
  - Syndecans: protein has a single transmembrane domain
  - Glypicans: protein is anchored to a lipid membrane
  - Interact with a variety of receptors from neighboring cells and regulate cell growth

# Syndecan

# Glypican

Heparan sulfate

Chondroitin sulfate

Outside

Membrane

Inside

$+NH_3$

Core protein

Cleavage site

$COO^-$

Globular domain

$H_3N^+$


S-S

S-S

$COO^-$

GPI anchor

Figure 7-26a  
Lehninger Principles of Biochemistry, Sixth Edition  
© 2013 W. H. Freeman and Company




# Proteoglycan Aggregates

---

- Hyaluronan and aggrecan form huge ( $M_r > 2 \cdot 10^8$ ) noncovalent aggregates
- Hold lots of water (1000× its weight); provides lubrication
- Very low friction material
- Covers joint surfaces: **articular cartilage**
  - Reduced friction
  - Load balancing





**Figure 7-28**

*Lehninger Principles of Biochemistry*, Sixth Edition  
© 2013 W. H. Freeman and Company

# Oligosaccharides in Recognition




Figure 7-37  
Lehninger Principles of Biochemistry, Seventh Edition  
© 2017 W. H. Freeman and Company