


## Chapter 5

# The Structure and Function of Large Biological Molecules



Lectures by  
Erin Barley  
Kathleen Fitzpatrick

© 2011 Pearson Education, Inc.

### Overview: The Molecules of Life

---

- All living things are made up of **four classes of large biological molecules**: carbohydrates, lipids, proteins, and nucleic acids
- Within cells, small organic molecules are joined together to form larger molecules
- **Macromolecules** are large molecules composed of **thousands of covalently connected atoms**
- Molecular structure and function are inseparable

---

Copyright © 2008 Pearson Education, Inc., publishing as Pearson Benjamin Cummings

Fig. 5-1



Copyright © 2008 Pearson Education, Inc., publishing as Pearson Benjamin Cummings.

### Concept 5.1: **Macromolecules** are **polymers**, built from **monomers**

---

- A **polymer** is a long molecule consisting of many similar building blocks
- These small building-block molecules are called **monomers**
- Three of the **four classes** of life's organic molecules are **polymers**:
  - **Carbohydrates**
  - **Proteins**
  - **Nucleic acids**

---

Copyright © 2008 Pearson Education, Inc., publishing as Pearson Benjamin Cummings


## The Synthesis and Breakdown of Polymers

- A **condensation reaction** or more specifically a **dehydration reaction** occurs when two monomers bond together through the **loss of a water molecule**
- **Enzymes** are macromolecules that speed up the dehydration process
- Polymers are disassembled to monomers by **hydrolysis**, a reaction that is essentially the reverse of the dehydration reaction

**PLAY** Animation: Polymers

Copyright © 2008 Pearson Education, Inc., publishing as Pearson Benjamin Cummings


Fig. 5-2a



(a) Dehydration reaction in the synthesis of a polymer

Copyright © 2008 Pearson Education, Inc., publishing as Pearson Benjamin Cummings.

Fig. 5-2b



## The Diversity of Polymers

- Each cell has **thousands of different kinds of macromolecules**
- Macromolecules **vary among cells of an organism**, **vary more within a species**, and **vary even more between species**
- What is the basis of this diversity??
- An immense variety of polymers can be built from a small set of monomers

## Concept 5.2: Carbohydrates serve as fuel and building material

---

- **Carbohydrates** include sugars and the polymers of sugars
- The simplest carbohydrates are **monosaccharides**, or single sugars
- Carbohydrate macromolecules are **polysaccharides**, polymers composed of many sugar building blocks

Copyright © 2008 Pearson Education, Inc., publishing as Pearson Benjamin Cummings

## Sugars


---

- **Monosaccharides** have molecular formulas that are usually multiples of  $(\text{CH}_2\text{O})_n$
- **Glucose** ( $\text{C}_6\text{H}_{12}\text{O}_6$ ) is the most common monosaccharide
- Most names of sugars end in **-ose**
- Monosaccharides are classified by
  - The **location of the carbonyl group** (as aldose or ketose)
  - The **number of carbons** in the carbon skeleton

Copyright © 2008 Pearson Education, Inc., publishing as Pearson Benjamin Cummings

Figure 5.3

The structure and classification of some monosaccharides.




© 2011 Pearson Education, Inc.

- Though often drawn as **linear skeletons**, in aqueous solutions many sugars **form rings**
- **Monosaccharides** serve as a **major fuel** for cells and as **raw material for building molecules**

Fig. 5-4

## Linear and ring forms of glucose




- A **disaccharide** is formed when a dehydration reaction joins **two monosaccharides**
- This covalent bond is called a **glycosidic linkage**
- **Examples of disaccharides:**
  - **Sucrose: Glucose + fructose**
  - **Maltose: glucose + glucose**
  - **Lactose: glucose + galactose**


**PLAY** Animation: Disaccharides

Fig. 5-5

## Examples of disaccharide synthesis



(a) Dehydration reaction in the synthesis of maltose



(b) Dehydration reaction in the synthesis of sucrose

Copyright © 2008 Pearson Education, Inc., publishing as Pearson Benjamin Cummings.

## Polysaccharides

- **Polysaccharides**, the polymers of sugars, have **storage and structural roles**
- The **structure and function** of a polysaccharide are determined by its **sugar monomers** and the **positions of glycosidic linkages**

Copyright © 2008 Pearson Education, Inc., publishing as Pearson Benjamin Cummings




## *Storage Polysaccharides*

- **Starch:**
- **A storage polysaccharide of plants**, consists entirely of  **$\alpha$ -glucose monomers**
- Plants store surplus starch as granules within **chloroplasts and other plastids**

Copyright © 2008 Pearson Education, Inc., publishing as Pearson Benjamin Cummings

Fig. 5-6



Copyright © 2008 Pearson Education, Inc., publishing as Pearson Benjamin Cummings.

- 
- **Glycogen:**
  - Is a **storage polysaccharide in animals**
  - Humans and other vertebrates **store glycogen mainly in liver and muscle cells**

Copyright © 2008 Pearson Education, Inc., publishing as Pearson Benjamin Cummings

### *Structural Polysaccharides*

---

- **Cellulose:**
- The polysaccharide **cellulose** is a major component of the tough wall of plant cells
- Like starch, cellulose is a polymer of glucose, but the **glycosidic linkages** differ
- The difference is based on **two ring forms for glucose: alpha ( $\alpha$ ) and beta ( $\beta$ )**


**PLAY**

Animation: Polysaccharides

Copyright © 2008 Pearson Education, Inc., publishing as Pearson Benjamin Cummings

Fig. 5-7a


## Starch and cellulose structures




(a)  $\alpha$  and  $\beta$  glucose ring structures

Copyright © 2008 Pearson Education, Inc., publishing as Pearson Benjamin Cummings.

Fig. 5-7bc



(b) Starch: 1-4 linkage of  $\alpha$  glucose monomers



(c) Cellulose: 1-4 linkage of  $\beta$  glucose monomers


Copyright © 2008 Pearson Education, Inc., publishing as Pearson Benjamin Cummings.

- Polymers with  **$\alpha$  glucose** are helical
- Polymers with  **$\beta$  glucose** are straight
- In straight structures, **H atoms** on one strand can bond with **OH groups** on other strands
- Parallel cellulose molecules held together this way are grouped into **microfibrils**, which form **strong building materials for plants**

Copyright © 2008 Pearson Education, Inc., publishing as Pearson Benjamin Cummings

Fig. 5-8

## The arrangement of cellulose in plant cell walls




Copyright © 2008 Pearson Education, Inc., publishing as Pearson Benjamin Cummings.

- 
- Enzymes that digest starch by hydrolyzing  $\alpha$  linkages can't hydrolyze  $\beta$  linkages in cellulose
  - Cellulose in human food passes through the digestive tract as insoluble fiber
  - Some **microbes secrete enzymes** to digest cellulose
  - Many **herbivores**, from **cows to termites**, have symbiotic relationships with these microbes
- 

Copyright © 2008 Pearson Education, Inc., publishing as Pearson Benjamin Cummings

Fig. 5-9




Copyright © 2008 Pearson Education, Inc., publishing as Pearson Benjamin Cummings

- **Chitin**
- Another structural polysaccharide, found in the **exoskeleton of arthropods**
- Chitin also provides structural support for the cell walls of many fungi

Copyright © 2008 Pearson Education, Inc., publishing as Pearson Benjamin Cummings

Fig. 5-10


## Chitin, a structural polysaccharide



(a) The structure of the chitin monomer.



(b) Chitin forms the exoskeleton of arthropods.



(c) Chitin is used to make a strong and flexible surgical thread.

Copyright © 2008 Pearson Education, Inc., publishing as Pearson Benjamin Cummings.

N-acetyl-  
Glucosamine

### Concept 5.3: **Lipids** are a diverse group of hydrophobic molecules

---

- **Lipids** are the one class of large biological molecules that **do not form polymers**
- The unifying feature of lipids is **having little or no affinity for water**
- Lipids are hydrophobic because they **consist mostly of hydrocarbons**, which form **nonpolar covalent bonds**
- The most biologically important lipids are **fats**, **phospholipids**, and **steroids**

Copyright © 2008 Pearson Education, Inc., publishing as Pearson Benjamin Cummings


### **Fats**

---

- **Fats** are constructed from two types of smaller molecules: **glycerol and fatty acids**
- **Glycerol** is a three-carbon alcohol with a **hydroxyl group** attached to each carbon
- A **fatty acid** consists of a **carboxyl group** attached to a long carbon skeleton

Copyright © 2008 Pearson Education, Inc., publishing as Pearson Benjamin Cummings

Fig. 5-11a




**Glycerol**

**(a) Dehydration reaction in the synthesis of a fat**

Copyright © 2008 Pearson Education, Inc., publishing as Pearson Benjamin Cummings.

Fig. 5-11b



**(b) Fat molecule (triacylglycerol)**

Copyright © 2008 Pearson Education, Inc., publishing as Pearson Benjamin Cummings.



- 
- Fats **separate from water** because water molecules form hydrogen bonds with each other and **exclude the fats**
  - **In a fat**, three fatty acids are joined to a glycerol by an **ester linkage**, creating a **triacylglycerol**, or **triglyceride**

Copyright © 2008 Pearson Education, Inc., publishing as Pearson Benjamin Cummings


- 
- **Fatty acids vary in length** (number of carbons) and in the number and locations of double bonds
  - **Saturated fatty acids** have the maximum number of hydrogen atoms possible and no double bonds
  - **Unsaturated fatty acids** have one or more double bonds

**PLAY** Animation: Fats


Copyright © 2008 Pearson Education, Inc., publishing as Pearson Benjamin Cummings

Figure 5.11

### (a) Saturated fat



### (b) Unsaturated fat



- Fats made from saturated fatty acids are called **saturated fats**, and are **solid at room temperature**
- Most animal fats are saturated
- Fats made from unsaturated fatty acids are called **unsaturated fats or oils**, and are **liquid at room temperature**
- Plant fats and fish fats are usually unsaturated

- 
- A diet rich in **saturated fats** may contribute to **cardiovascular disease** through plaque deposits
  - **Hydrogenation** is the process of converting unsaturated fats to saturated fats by adding hydrogen
  - Hydrogenating vegetable oils also creates unsaturated fats with **trans double bonds**
  - These **trans fats** may contribute more than saturated fats to **cardiovascular disease**

Copyright © 2008 Pearson Education, Inc., publishing as Pearson Benjamin Cummings

- 
- The **major function of fats** is **energy storage**
  - Humans and other mammals **store** their fat in **adipose cells**
  - **Adipose tissue** also **cushions vital organs and insulates the body**

---


Copyright © 2008 Pearson Education, Inc., publishing as Pearson Benjamin Cummings

## Phospholipids

- In a **phospholipid**, **two fatty acids** and a **phosphate group** are attached to **glycerol**
- The two fatty acid tails are hydrophobic, but the **phosphate group** and its attachments form a **hydrophilic head**

Copyright © 2008 Pearson Education, Inc., publishing as Pearson Benjamin Cummings

Fig. 5-13ab




Copyright © 2008 Pearson Education, Inc., publishing as Pearson Benjamin Cummings.

- 
- When phospholipids are added to water, they **self-assemble into a bilayer**, with the hydrophobic tails pointing toward the interior
  - The structure of phospholipids results in a **bilayer** arrangement found in **cell membranes**
  - Phospholipids are the **major component of all cell membranes**
- 

Copyright © 2008 Pearson Education, Inc., publishing as Pearson Benjamin Cummings

Fig. 5-14




Copyright © 2008 Pearson Education, Inc., publishing as Pearson Benjamin Cummings.

## Steroids

- **Steroids** are lipids characterized by a **carbon skeleton consisting of four fused rings**
- **Cholesterol**, an important steroid, is a component in **animal cell membranes**
- Although cholesterol is essential in animals, **high levels in the blood** may contribute to **cardiovascular disease**

Copyright © 2008 Pearson Education, Inc., publishing as Pearson Benjamin Cummings

Fig. 5-15



Copyright © 2008 Pearson Education, Inc., publishing as Pearson Benjamin Cummings.

## Concept 5.4: **Proteins** have many structures, resulting in a **wide range of functions**

- Proteins account for more than **50% of the dry mass of most cells**
- Protein functions include: **structural support**, **storage**, **transport**, **cellular communications**, **movement**, and **defense against foreign substances**


Copyright © 2008 Pearson Education, Inc., publishing as Pearson Benjamin Cummings

Table 5-1

| Table 5.1 An Overview of Protein Functions |  |   |
|--|--|---|
| Type of Protein                            | Function                                     | Examples  |
| Enzymatic proteins                         | Selective acceleration of chemical reactions | Digestive enzymes   |
| Structural proteins                        | Support                                      | Silk fibers; collagen and elastin in animal connective tissues; keratin in hair, horns, feathers, and other skin appendages |
| Storage proteins                           | Storage of amino acids                       | Ovalbumin in egg white; casein, the protein of milk; storage proteins in plant seeds  |
| Transport proteins                         | Transport of other substances                | Hemoglobin, transport proteins  |
| Hormonal proteins                          | Coordination of an organism's activities     | Insulin, a hormone secreted by the pancreas   |
| Receptor proteins                          | Response of cell to chemical stimuli         | Receptors in nerve cell membranes   |
| Contractile and motor proteins             | Movement                                     | Actin and myosin in muscles, proteins in cilia and flagella   |
| Defensive proteins                         | Protection against disease                   | Antibodies combat bacteria and viruses.   |


Copyright © 2008 Pearson Education, Inc., publishing as Pearson Benjamin Cummings.

Figure 5.15-a



© 2011 Pearson Education, Inc.

Figure 5.15-b



© 2011 Pearson Education, Inc.




Figure 5.15a

### Enzymatic proteins

**Function:** Selective acceleration of chemical reactions

**Example:** Digestive enzymes catalyze the hydrolysis of bonds in food molecules.




© 2011 Pearson Education, Inc.

Figure 5.15b

### Storage proteins

**Function:** Storage of amino acids

**Examples:** Casein, the protein of milk, is the major source of amino acids for baby mammals. Plants have storage proteins in their seeds. Ovalbumin is the protein of egg white, used as an amino acid source for the developing embryo.




© 2011 Pearson Education, Inc.

Figure 5.15c

### Hormonal proteins

**Function:** Coordination of an organism's activities

**Example:** Insulin, a hormone secreted by the pancreas, causes other tissues to take up glucose, thus regulating blood sugar concentration




© 2011 Pearson Education, Inc.

Figure 5.15d

### Contractile and motor proteins

**Function:** Movement

**Examples:** Motor proteins are responsible for the undulations of cilia and flagella. Actin and myosin proteins are responsible for the contraction of muscles.




© 2011 Pearson Education, Inc.

Figure 5.15e

### Defensive proteins

**Function:** Protection against disease

**Example:** Antibodies inactivate and help destroy viruses and bacteria.




© 2011 Pearson Education, Inc.

Figure 5.15f

### Transport proteins

**Function:** Transport of substances

**Examples:** Hemoglobin, the iron-containing protein of vertebrate blood, transports oxygen from the lungs to other parts of the body. Other proteins transport molecules across cell membranes.




© 2011 Pearson Education, Inc.

Figure 5.15g

### Receptor proteins

**Function:** Response of cell to chemical stimuli

**Example:** Receptors built into the membrane of a nerve cell detect signaling molecules released by other nerve cells.




© 2011 Pearson Education, Inc.

Figure 5.15h

### Structural proteins

**Function:** Support

**Examples:** Keratin is the protein of hair, horns, feathers, and other skin appendages. Insects and spiders use silk fibers to make their cocoons and webs, respectively. Collagen and elastin proteins provide a fibrous framework in animal connective tissues.



© 2011 Pearson Education, Inc.


- **Enzymes** are a type of protein that acts as a **catalyst to speed up chemical reactions**
- Enzymes can perform their functions repeatedly, functioning as workhorses that carry out the processes of life

**PLAY**

Animation: Enzymes

Copyright © 2008 Pearson Education, Inc., publishing as Pearson Benjamin Cummings

Fig. 5-16



## Polypeptides

---

- **Polypeptides** are polymers built from the same set of **20 amino acids**
- A **protein** consists of one or more polypeptides

---

Copyright © 2008 Pearson Education, Inc., publishing as Pearson Benjamin Cummings

## *Amino Acid Monomers*

---


- **Amino acids** are organic molecules with **carboxyl and amino groups**
- Amino acids **differ in their properties** due to differing side chains, called **R groups**

---

Copyright © 2008 Pearson Education, Inc., publishing as Pearson Benjamin Cummings


Figure 5.UN01

Side chain (R group)



© 2011 Pearson Education, Inc.

Figure 5.16




## Amino Acid Polymers

- Amino acids are **linked by peptide bonds**
- A polypeptide is a polymer of amino acids
- Polypeptides range in length from **a few to more than a thousand monomers**
- Each polypeptide has a **unique linear sequence of amino acids**

Copyright © 2008 Pearson Education, Inc., publishing as Pearson Benjamin Cummings

Figure 5.17



© 2011 Pearson Education, Inc.



## Protein Structure and Function


---

- A functional protein consists of one or more polypeptides twisted, folded, and coiled into a unique shape

---


Copyright © 2008 Pearson Education, Inc., publishing as Pearson Benjamin Cummings

Figure 5.18



(a) A ribbon model

© 2011 Pearson Education, Inc.




(b) A space-filling model

- 
- The sequence of amino acids determines a protein's three-dimensional structure
  - A protein's structure determines its function

---

Copyright © 2008 Pearson Education, Inc., publishing as Pearson Benjamin Cummings

Fig. 5-20



Copyright © 2008 Pearson Education, Inc., publishing as Pearson Benjamin Cummings.

## *Four Levels of Protein Structure*

- The **primary structure** of a protein is its *unique sequence of amino acids*
- **Secondary structure**, found in most proteins, consists of *coils and folds in the polypeptide chain*
- **Tertiary structure** is determined by *interactions among various side chains (R groups)*
- **Quaternary structure** results when a protein consists of *multiple polypeptide chains*

**PLAY** Animation: Protein Structure Introduction

Copyright © 2008 Pearson Education, Inc., publishing as Pearson Benjamin Cummings

Figure 5.20a






Figure 5.20b



- **Primary structure**, the sequence of amino acids in a protein, is like the order of letters in a long word
- Primary structure is determined by **inherited genetic information**

**PLAY** Animation: Primary Protein Structure

Fig. 5-21a



- The coils and folds of **secondary structure** result from **hydrogen bonds** between repeating constituents of the **polypeptide backbone**
- Typical secondary structures are a coil called an  **$\alpha$  helix** and a folded structure called a  **$\beta$  pleated sheet**

**PLAY**

Animation: Secondary Protein Structure

Copyright © 2008 Pearson Education, Inc., publishing as Pearson Benjamin Cummings

Figure 5.20c

### Secondary structure





Fig. 5-21d




- 
- **Tertiary structure** is determined by **interactions between R groups**, rather than interactions between backbone constituents
  - These interactions between R groups include **hydrogen bonds**, **ionic bonds**, **hydrophobic interactions**, and **van der Waals interactions**
  - **Strong covalent bonds** called **disulfide bridges** may reinforce the protein's structure

**PLAY**

Animation: Tertiary Protein Structure


Copyright © 2008 Pearson Education, Inc., publishing as Pearson Benjamin Cummings

Figure 5.20e



© 2011 Pearson Education, Inc.

Figure 5.20f



- **Quaternary structure** results when **two or more polypeptide chains form one macromolecule**
- **Collagen** is a **fibrous protein** consisting of **three polypeptides** coiled like a rope
- **Hemoglobin** is a **globular protein** consisting of **four polypeptides: two alpha and two beta chains**


PLAY

Animation: Quaternary Protein Structure



Figure 5.20g

**Quaternary structure**




**Transthyretin  
protein  
(four identical  
polypeptides)**

© 2011 Pearson Education, Inc.


Figure 5.20h

**Collagen**



© 2011 Pearson Education, Inc.

Figure 5.20i











### *Sickle-Cell Disease: A Change in Primary Structure*

---


- A slight change in primary structure can affect a protein's structure and ability to function
- Sickle-cell disease, an inherited blood disorder, results from a single amino acid substitution in the protein hemoglobin

Figure 5.21

|                        | Primary Structure   | Secondary and Tertiary Structures  | Quaternary Structure  | Function  | Red Blood Cell Shape   |
|------------------------|---|--|---|---|--|
| Normal hemoglobin      | <ol style="list-style-type: none"> <li>1 Val</li> <li>2 His</li> <li>3 Leu</li> <li>4 Thr</li> <li>5 Pro</li> <li>6 Glu</li> <li>7 Glu</li> </ol> |  <p>β subunit</p>                                   | <p>Normal hemoglobin</p>       | <p>Molecules do not associate with one another; each carries oxygen.</p>         |  <p>10 μm</p> |
| Sickle-cell hemoglobin | <ol style="list-style-type: none"> <li>1 Val</li> <li>2 His</li> <li>3 Leu</li> <li>4 Thr</li> <li>5 Pro</li> <li>6 Val</li> <li>7 Glu</li> </ol> | <p>Exposed hydrophobic region</p>  <p>β subunit</p> | <p>Sickle-cell hemoglobin</p>  | <p>Molecules crystallize into a fiber; capacity to carry oxygen is reduced.</p>  |  <p>10 μm</p> |

© 2011 Pearson Education, Inc.

Fig. 5-22c



**Normal red blood cells are full of individual hemoglobin molecules, each carrying oxygen.**

**Fibers of abnormal hemoglobin deform red blood cell into sickle shape.**


Copyright © 2008 Pearson Education, Inc., publishing as Pearson Benjamin Cummings.

## *What Determines Protein Structure?*

- In addition to **primary structure**, **physical and chemical conditions** can affect structure
- Alterations in **pH**, **salt concentration**, **temperature**, or other **environmental factors** can cause a protein to unravel
- *This loss of a protein's native structure is called **denaturation***
- A **denatured protein** is **biologically inactive**

Copyright © 2008 Pearson Education, Inc., publishing as Pearson Benjamin Cummings

Fig. 5-23




Copyright © 2008 Pearson Education, Inc., publishing as Pearson Benjamin Cummings.

## *Protein Folding in the Cell*

- It is hard to predict a protein's structure from its primary structure
- Most proteins probably go through several states on their way to a stable structure
- **Chaperonins** are protein molecules that assist the proper folding of other proteins

Copyright © 2008 Pearson Education, Inc., publishing as Pearson Benjamin Cummings

Fig. 5-24




Copyright © 2008 Pearson Education, Inc., publishing as Pearson Benjamin Cummings

- Scientists use **X-ray crystallography** to determine a protein's structure
- Another method is **nuclear magnetic resonance (NMR) spectroscopy**, which does not require protein crystallization
- **Bioinformatics** uses computer programs to predict protein structure from amino acid sequences


Copyright © 2008 Pearson Education, Inc., publishing as Pearson Benjamin Cummings

Fig. 5-25

#### EXPERIMENT



#### RESULTS



Copyright © 2008 Pearson Education, Inc., publishing as Pearson Benjamin Cummings

## Concept 5.5: **Nucleic acids** store and transmit hereditary information

---

- The **amino acid sequence** of a polypeptide is programmed by a unit of inheritance called a **gene**
- **Genes are made of DNA**, a **nucleic acid**

---

Copyright © 2008 Pearson Education, Inc., publishing as Pearson Benjamin Cummings

## The Roles of Nucleic Acids

---

- There are two types of nucleic acids:
  - **Deoxyribonucleic acid (DNA)**
  - **Ribonucleic acid (RNA)**
- DNA provides directions for **its own replication**
- DNA directs synthesis of messenger RNA (mRNA) and, through mRNA, controls protein synthesis
- Protein synthesis occurs in ribosomes

---

Copyright © 2008 Pearson Education, Inc., publishing as Pearson Benjamin Cummings

Fig. 5-26-1




Fig. 5-26-2







Fig. 5-26-3



## The Structure of Nucleic Acids


- Nucleic acids are polymers called **polynucleotides**
- Each polynucleotide is made of monomers called **nucleotides**
- Each nucleotide consists of: a **nitrogenous base**, a **pentose sugar**, and a **phosphate group**
- The portion of a nucleotide without the phosphate group is called a **nucleoside**

Copyright © 2008 Pearson Education, Inc., publishing as Pearson Benjamin Cummings

## Nucleotide Monomers

- **Nucleoside = nitrogenous base + sugar**
- There are **two families of nitrogenous bases**:
  - **Pyrimidines** (cytosine, thymine, and uracil) have a single six-membered ring
  - **Purines** (adenine and guanine) have a six-membered ring fused to a five-membered ring
- In **DNA**, the sugar is **deoxyribose**; in **RNA**, the sugar is **ribose**
- **Nucleotide = nucleoside + phosphate group**

Copyright © 2008 Pearson Education, Inc., publishing as Pearson Benjamin Cummings



## *Nucleotide Polymers*

---

- Nucleotide polymers are linked together to build a polynucleotide
  - Adjacent nucleotides are joined by covalent bonds that form **between the –OH group** on the **3' carbon of one nucleotide** and the **phosphate on the 5' carbon on the next**
  - These links create a **backbone of sugar-phosphate** units with **nitrogenous bases as appendages**
  - The sequence of bases along a DNA or mRNA polymer is unique for each gene
- 



Copyright © 2008 Pearson Education, Inc., publishing as Pearson Benjamin Cummings

## *The DNA Double Helix*

---

- A **DNA molecule has two polynucleotides** spiraling around an imaginary axis, forming a **double helix**
  - In the DNA double helix, the two backbones run in **opposite 5' → 3' directions** from each other, an arrangement referred to as **antiparallel**
  - The **nitrogenous bases in DNA pair up and form hydrogen bonds**: **adenine (A) always with thymine (T)**, and **guanine (G) always with cytosine (C)**
- 

Copyright © 2008 Pearson Education, Inc., publishing as Pearson Benjamin Cummings











## DNA and Proteins as Tape Measures of Evolution

- The linear sequences of nucleotides in DNA molecules are passed from parents to offspring
- Two closely related species are more similar in DNA than are more distantly related species
- Molecular biology can be used to assess evolutionary kinship







Copyright © 2008 Pearson Education, Inc., publishing as Pearson Benjamin Cummings

Figure 5.UN02

| Large Biological Molecules   | Components   | Examples   | Functions   |
|--|--|--|---|
| <p><b>CONCEPT 5.2</b><br/>Carbohydrates serve as fuel and building material</p>                                  |  <p>Monosaccharide monomer</p>  | <p><b>Monosaccharides:</b> glucose, fructose</p> <p><b>Disaccharides:</b> lactose, sucrose</p> <p><b>Polysaccharides:</b></p> <ul style="list-style-type: none"> <li>• Cellulose (plants)</li> <li>• Starch (plants)</li> <li>• Glycogen (animals)</li> <li>• Chitin (animals and fungi)</li> </ul>  | <p>Fuel; carbon sources that can be converted to other molecules or combined into polymers</p> <ul style="list-style-type: none"> <li>• Strengthens plant cell walls</li> <li>• Stores glucose for energy</li> <li>• Stores glucose for energy</li> <li>• Strengthens exoskeletons and fungal cell walls</li> </ul>   |
| <p><b>CONCEPT 5.3</b><br/>Lipids are a diverse group of hydrophobic molecules</p>                                | <p><b>Glycerol</b></p>  <p>3 fatty acids</p> <p><b>Phospholipids:</b> phosphate group + 2 fatty acids</p>  <p>Head with P<br/>2 fatty acids</p> <p><b>Steroid backbone</b></p>  | <p><b>Triacylglycerols (fats or oils):</b> glycerol + 3 fatty acids</p> <p><b>Phospholipids:</b> phosphate group + 2 fatty acids</p> <p><b>Steroids:</b> four fused rings with attached chemical groups</p>  | <p>Important energy source</p>  <p>Lipid bilayers of membranes</p>  <p>Hydrophilic heads<br/>Hydrophobic tails</p> <ul style="list-style-type: none"> <li>• Component of cell membranes (cholesterol)</li> <li>• Signaling molecules that travel through the body (hormones)</li> </ul> |
| <p><b>CONCEPT 5.4</b><br/>Proteins include a diversity of structures, resulting in a wide range of functions</p> | <p><b>Amino acid monomer (20 types)</b></p>   | <ul style="list-style-type: none"> <li>• Enzymes</li> <li>• Structural proteins</li> <li>• Storage proteins</li> <li>• Transport proteins</li> <li>• Hormones</li> <li>• Receptor proteins</li> <li>• Motor proteins</li> <li>• Defensive proteins</li> </ul>  | <ul style="list-style-type: none"> <li>• Catalyze chemical reactions</li> <li>• Provide structural support</li> <li>• Store amino acids</li> <li>• Transport substances</li> <li>• Coordinate organismal responses</li> <li>• Receive signals from outside cell</li> <li>• Function in cell movement</li> <li>• Protect against disease</li> </ul>  |
| <p><b>CONCEPT 5.5</b><br/>Nucleic acids store, transmit, and help express hereditary information</p>             | <p><b>Nucleotide monomer</b></p>  <p>Phosphate group<br/>Sugar<br/>Nitrogenous base</p>   | <p><b>DNA:</b></p> <ul style="list-style-type: none"> <li>• Sugar = deoxyribose</li> <li>• Nitrogenous bases = C, G, A, T</li> <li>• Usually double-stranded</li> </ul> <p><b>RNA:</b></p> <ul style="list-style-type: none"> <li>• Sugar = ribose</li> <li>• Nitrogenous bases = C, G, A, U</li> <li>• Usually single-stranded</li> </ul> | <p>Stores hereditary information</p> <p>Various functions during gene expression, including carrying instructions from DNA to ribosomes</p>   |





© 2011 Pearson Education, Inc.

Figure 5.UN02a

| Large Biological Molecules   | Components   | Examples   | Functions   |
|--|--|--|---|
| <p><b>CONCEPT 5.2</b></p> <p>Carbohydrates serve as fuel and building material</p>   |  <p>Monosaccharide monomer</p>          | <p><b>Monosaccharides:</b> glucose, fructose</p>   | <p>Fuel: carbon sources that can be converted to other molecules or combined into polymers</p> <ul style="list-style-type: none"> <li>• Strengthens plant cell walls</li> <li>• Stores glucose for energy</li> <li>• Stores glucose for energy</li> <li>• Strengthens exoskeletons and fungal cell walls</li> </ul> |
|  |  | <p><b>Disaccharides:</b> lactose, sucrose</p> <p><b>Polysaccharides:</b></p> <ul style="list-style-type: none"> <li>• Cellulose (plants)</li> <li>• Starch (plants)</li> <li>• Glycogen (animals)</li> <li>• Chitin (animals and fungi)</li> </ul> |   |
| <p><b>CONCEPT 5.3</b></p> <p>Lipids are a diverse group of hydrophobic molecules</p> | <p>Glycerol</p>  <p>3 fatty acids</p>   | <p><b>Triacylglycerols</b> (fats or oils): glycerol + 3 fatty acids</p>  | <p>Important energy source</p>   |
|  |  <p>Head with (P)<br/>2 fatty acids</p> | <p><b>Phospholipids:</b> phosphate group + 2 fatty acids</p>   | <p>Lipid bilayers of membranes</p>  <p>Hydrophilic heads<br/>Hydrophobic tails</p>  |
|  |  <p>Steroid backbone</p>                | <p><b>Steroids:</b> four fused rings with attached chemical groups</p>   | <ul style="list-style-type: none"> <li>• Component of cell membranes (cholesterol)</li> <li>• Signaling molecules that travel through the body (hormones)</li> </ul>  |

© 2011 Pearson Education, Inc.

Figure 5.UN02b

| Large Biological Molecules  | Components   | Examples  | Functions  |
|---|--|---|--|
| <p><b>CONCEPT 5.4</b></p> <p>Proteins include a diversity of structures, resulting in a wide range of functions</p> |  <p>Amino acid monomer (20 types)</p> | <ul style="list-style-type: none"> <li>• Enzymes</li> <li>• Structural proteins</li> <li>• Storage proteins</li> <li>• Transport proteins</li> <li>• Hormones</li> <li>• Receptor proteins</li> <li>• Motor proteins</li> <li>• Defensive proteins</li> </ul> | <ul style="list-style-type: none"> <li>• Catalyze chemical reactions</li> <li>• Provide structural support</li> <li>• Store amino acids</li> <li>• Transport substances</li> <li>• Coordinate organismal responses</li> <li>• Receive signals from outside cell</li> <li>• Function in cell movement</li> <li>• Protect against disease</li> </ul>   |
|   |  | <p><b>Nitrogenous base</b></p>  <p>Phosphate group<br/>Sugar<br/>Nucleotide monomer</p>  | <p><b>DNA:</b> </p> <ul style="list-style-type: none"> <li>• Sugar = deoxyribose</li> <li>• Nitrogenous bases = C, G, A, T</li> <li>• Usually double-stranded</li> </ul> <p><b>RNA:</b> </p> <ul style="list-style-type: none"> <li>• Sugar = ribose</li> <li>• Nitrogenous bases = C, G, A, U</li> <li>• Usually single-stranded</li> </ul> |

© 2011 Pearson Education, Inc.