





# PARABOLAS

## Parabola Vertex (0, 0)




Concept	Equation	Example
Parabola with vertex (0, 0) and vertical axis	$x^2 = 4py$ $p > 0$ : opens upward $p < 0$ : opens downward Focus: $(0, p)$ Directrix: $y = -p$	$x^2 = -2y$ has $4p = -2$ or $p = -\frac{1}{2}$ The parabola opens downward with vertex $(0, 0)$ , focus $(0, -\frac{1}{2})$ , and directrix $y = \frac{1}{2}$ 
Parabola with vertex (0, 0) and horizontal axis	$y^2 = 4px$ $p > 0$ : opens to the right $p < 0$ : opens to the left Focus: $(p, 0)$ Directrix: $x = -p$	$y^2 = 4x$ has $4p = 4$ or $p = 1$ The parabola opens to the right with vertex $(0, 0)$ , focus $(1, 0)$ , and directrix $x = -1$ 

# Parabola Vertex (h, k)






Concept	Equation	Example
<p>Parabola with vertex (h, k) and horizontal axis</p>	<p><math>(y - k)^2 = 4p(x - h)</math></p> <p><math>p &gt; 0</math>: opens to the right</p> <p><math>p &lt; 0</math>: opens to the left</p> <p>Focus: <math>(h + p, k)</math></p> <p>Directrix: <math>x = h - p</math></p>	<p><math>(y + 1)^2 = -2(x + 2)</math> has <math>p = -\frac{1}{2}</math></p> <p>The parabola opens to the left with vertex <math>(-2, -1)</math>, focus <math>(-\frac{5}{2}, -1)</math>, and directrix <math>x = -\frac{3}{2}</math></p> 
<p>Parabola with vertex (h, k) and vertical axis</p>	<p><math>(x - h)^2 = 4p(y - k)</math></p> <p><math>p &gt; 0</math>: opens upwards</p> <p><math>p &lt; 0</math>: opens downwards</p> <p>Focus: <math>(h, k + p)</math></p> <p>Directrix: <math>y = k - p</math></p>	<p><math>(x - 1)^2 = 8(y - 3)</math> has <math>p = 2</math>.</p> <p>The parabola opens upward with vertex <math>(1, 3)</math>, focus <math>(1, 5)</math>, and directrix <math>y = 1</math>.</p> 

# HYPERBOLA

Concept	Equation	Example
<p>Hyperbola with center (0, 0)</p> 	<p>Standard equation</p> <p>Transverse axis: horizontal</p> $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$ <p>Transverse axis: vertical</p> $\frac{y^2}{a^2} - \frac{x^2}{b^2} = 1$	<p><math>\frac{x^2}{4} - \frac{y^2}{9} = 1; a = 2, b = 3</math></p> <p>Transverse axis: vertical</p> <p>Vertices (0, ±2); foci: (0, ±√13)</p> <p>(c<sup>2</sup> = a<sup>2</sup> + b<sup>2</sup> = 4 + 9 = 13, so c = √13.)</p> <p>Asymptotes: <math>y = \pm \frac{2}{3}x</math></p> 
<p>Hyperbola with center (h, k)</p>	<p>Standard Equation</p> <p>Transverse axis: horizontal</p> $\frac{(x-h)^2}{a^2} - \frac{(y-k)^2}{b^2} = 1$ <p>Transverse axis: vertical</p> $\frac{(y-k)^2}{a^2} - \frac{(x-h)^2}{b^2} = 1$	<p><math>\frac{(x-1)^2}{4} - \frac{(y+1)^2}{9} = 1; a = 2, b = 3</math></p> <p>Transverse axis: horizontal; center (1, -1)</p> <p>Vertices (1 ± 2, -1); foci: (1 ± √13, -1)</p> <p>(c<sup>2</sup> = a<sup>2</sup> + b<sup>2</sup> = 4 + 9 = 13, so c = √13.)</p> <p>Asymptotes: <math>y = \pm \frac{3}{2}(x-1) - 1</math></p> 

# ELLIPSE, HYPERBOLA AND PARABOLA

## ELLIPSE

Concept	Equation	Example
<b>Ellipse with Center (0, 0)</b>	Standard equation with $a > b > 0$  Horizontal major axis: $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$  Vertical major axis: $\frac{x^2}{b^2} + \frac{y^2}{a^2} = 1$	$\frac{x^2}{4} + \frac{y^2}{9} = 1$ ; $a = 3$ , $b = 2$ Center (0, 0); major axis: vertical Vertices: (0, $\pm 3$ ); foci: (0, $\pm \sqrt{5}$ ) ( $c^2 = a^2 - b^2 = 9 - 4 = 5$ , so $c = \sqrt{5}$ .) 
<b>Ellipse with center (h, k)</b>	Standard equation with $a > b > 0$  Horizontal major axis: $\frac{(x-h)^2}{a^2} + \frac{(y-k)^2}{b^2} = 1$  Vertical major axis: $\frac{(x-h)^2}{b^2} + \frac{(y-k)^2}{a^2} = 1$	$\frac{(x-1)^2}{4} + \frac{(y+1)^2}{9} = 1$ ; $a = 3$ , $b = 2$ 
<b>Circle with center (h, k) and radius r</b>	Standard equation $(x-h)^2 + (y-k)^2 = r^2$  A circle is an ellipse with $a = b = r$ .	$(x-2)^2 + (y+2)^2 = 9$ Center: (2, -2); radius: $r = 3$ 
Area inside an ellipse	$A = \pi ab$	The area inside the ellipse give by $\frac{x^2}{49} + \frac{y^2}{9} = 1$ is  $A = \pi(7)(3) = 21\pi$ square units.