

11.2 Calculus with parametric curves

Note Title

۳۳/۰۶/۱۰

Tangents and Areas:

If $x = f(t)$ and $y = g(t)$ are differentiable funcs of t , then by Chain rule we have that

$$\frac{dy}{dt} = \frac{dy}{dx} \cdot \frac{dx}{dt},$$

so if $\frac{dx}{dt} \neq 0$, we get that

$$\boxed{\frac{dy}{dx} = \frac{dy/dt}{dx/dt}}$$

Similarly, we can prove that

$$\frac{d^2y}{dx^2} = \frac{(dy'/dt)}{(dx/dt)}$$

و لپ کپ لپ لپ $y' = \frac{dy}{dx}$ نگو، دانه سی t

Examples: 1) a) Find the tangent to the curve

$x = \sec t$, $y = \tan t$ $-\frac{\pi}{2} < t < \frac{\pi}{2}$
at the point $(\sqrt{2}, 1)$ when $t = \frac{\pi}{4}$.


Sol: $\frac{dy}{dx} = \frac{(dy/dt)}{(dx/dt)} = \frac{\sec^2 t}{\sec t \tan t} = \frac{\sec t}{\tan t}$

$$= \csc t$$

$$\therefore \text{at } t = \frac{\pi}{4}, \quad \frac{dy}{dx} = \sqrt{2}$$

Eqn of the Tangent: $m_T = \sqrt{2}$, $P(\sqrt{2}, 1)$, so

$$y = y_0 + m_T(x - x_0) = 1 + \sqrt{2}(x - \sqrt{2}) = \boxed{\sqrt{2}x - 1}$$



b) Find $\frac{d^2y}{dx^2}$ at $t = \frac{\pi}{4}$.

sol: $y'(t) = \frac{dy}{dx} = \csc t \Rightarrow$

$$\frac{d^2y}{dx^2} = \frac{(dy'/dt)}{(dx/dt)} = \frac{-\csc t \cot t}{\sec t \tan t} = -\cot^3 t$$

at $t = \frac{\pi}{4}$, $\frac{d^2y}{dx^2} = -\cot^3 \frac{\pi}{4} = \boxed{-1}$

2) Find d^2y/dx^2 as a function of t if

$$x = t - t^2, \quad y = t - t^3, \quad -\infty < t < \infty$$

sol: $y'(t) = \frac{dy}{dx} = \frac{dy/dt}{dx/dt} = \frac{1 - 3t^2}{1 - 2t}$

Now $\frac{dy'}{dt} = \frac{(1-2t)(-6t) - (1-3t^2)(-2)}{(1-2t)^2} = \frac{2-6t+6t^2}{(1-2t)^2}$

$$\therefore \frac{d^2y}{dx^2} = \frac{dy'/dt}{dx/dt} = \frac{(2-6t+6t^2)/(1-2t)^2}{(1-2t)} = \frac{2-6t+6t^2}{(1-2t)^3}$$

3) Find the normal to the curve

$$x = 2t^2 + 3, \quad y = t^4 \quad \text{at } t = -1$$

sol: The slope of the tangent to the curve at $t = -1$ is


$$m_T = \left. \frac{dy}{dx} \right|_{t=-1} = \left. \frac{dy/dt}{dx/dt} \right|_{t=-1} = \left. \frac{4t^3}{4t} \right|_{t=-1} = 1$$

So the slope of the normal line is $m_{\perp} = \frac{-1}{m_T} = -1$

Moreover, at $t = -1$, $(x, y) = (5, 1)$

\therefore Normal line:

$$y = 1 + (-1)(x - 5) = \boxed{6 - x}$$



Area: تذكر أنه إذا كانت $y = f(x) \geq 0$ لمساحة تحتها (تحتها) $a \leq x \leq b$ فإنها \sim

$$\text{Area} = \int_a^b y \, dx$$


فإنها كلمة لدينا معادلات بسيطة أفاها برأعي (التكويبي بدلًا من التكاليفي).

Example: Find the area enclosed by the ellipse

$$x = a \cos t, \quad y = b \sin t, \quad 0 \leq t \leq 2\pi$$

sol: Clearly,

$$\text{Area} = 4 \int_0^a y \, dx$$



but $x=0$ when $y=b$ at $t=\pi/2$ and $x=a$ at $t=0$. Moreover, $y = b \sin t$ and

$$dx = a \cdot -\sin t \, dt = -a \sin t \, dt \implies$$

$$\begin{aligned} \text{Area} &= 4 \int_{\pi/2}^0 b \sin t \cdot (-a \sin t) \, dt = 4ab \int_0^{\pi/2} \sin^2 t \, dt \\ &= \frac{4ab}{2} \int_0^{\pi/2} (1 - \cos 2t) \, dt = 2ab \left[t - \frac{\sin 2t}{2} \right]_0^{\pi/2} \\ &= 2ab \left[\left(\frac{\pi}{2} - 0 \right) - (0 - 0) \right] = \boxed{ab\pi} \end{aligned}$$

Length of a Parametrically Defined Curve:

DEFINITION If a curve C is defined parametrically by $x = f(t)$ and $y = g(t)$, $a \leq t \leq b$, where f' and g' are continuous and not simultaneously zero on $[a, b]$, and C is traversed exactly once as t increases from $t = a$ to $t = b$, then **the length of C** is the definite integral

$$L = \int_a^b \sqrt{[f'(t)]^2 + [g'(t)]^2} \, dt.$$

Examples: 1) prove that the length of the circle of radius r is $2\pi r$.

sol: Using the parametrization

$$x = r \sin t, \quad y = r \cos t, \quad 0 \leq t \leq 2\pi$$

or any other parametrization

we get that (هناك عدد لا نهائي من التمثيلات المتكافئة لنفس الدائرة) $\left(\frac{dx}{dt} = r \cos t, \quad \frac{dy}{dt} = -r \sin t \right)$ which are continuous

$$\frac{dx}{dt} = r \cos t, \quad \frac{dy}{dt} = -r \sin t$$

and not simultaneously zero, so

$$L = \int_0^{2\pi} \sqrt{r^2 \cos^2 t + r^2 \sin^2 t} dt = \int_0^{2\pi} r dt = \boxed{2\pi r}$$

2) Find the length of the curve

$$x = 8 \cos t + 8t \sin t, \quad y = 8 \sin t - 8t \cos t, \quad 0 \leq t \leq \frac{\pi}{2}$$

sol: $\frac{dx}{dt} = -8 \sin t + 8 \sin t + 8t \cos t = 8t \cos t$

$$\frac{dy}{dt} = 8 \cos t - 8 \cos t + 8t \sin t = 8t \sin t$$

$$\begin{aligned} \therefore L &= \int_0^{\pi/2} \sqrt{64t^2 \cos^2 t + 64t^2 \sin^2 t} dt = \int_0^{\pi/2} 8t dt = 4t^2 \Big|_0^{\pi/2} \\ &= 4 \left[\frac{\pi^2}{4} - 0 \right] = \boxed{\pi^2} \end{aligned}$$


EXAMPLE 5 Find the length of the astroid (Figure 11.13)

$$x = \cos^3 t, \quad y = \sin^3 t, \quad 0 \leq t \leq 2\pi.$$

sol: $\frac{dx}{dt} = -3 \cos^2 t \sin t$

$$\frac{dy}{dt} = 3 \sin^2 t \cos t$$

لاحظ أنه تردد (كثافة) من اتجاه t وليس من الاتجاه x (لأنه من المماسات)



$$\begin{aligned}
\therefore L &= 4 \int_0^{\pi/2} \sqrt{9 \cos^4 t \sin^2 t + 9 \sin^4 t \cos^2 t} dt \\
&= 4 \int_0^{\pi/2} \sqrt{9 \sin^2 t \cos^2 t} dt = 4 \int_0^{\pi/2} 3 |\sin t \cos t| dt \\
&= \frac{12}{2} \int_0^{\pi/2} \sin 2t dt \quad \left(\sin 2t = 2 \sin t \cos t, \text{ and } \sin 2t > 0 \right. \\
&\quad \left. \text{in the interval } [0, \pi/2] \right) \\
&= 6 \cdot \left[-\frac{\cos 2t}{2} \right]_0^{\pi/2} = -3 (-1 - 1) = \boxed{6}
\end{aligned}$$

ملوظة: سابقاً تم دراسة طول الكحن و كان قانونه يستخدم المعادلات العادية

$$L = \int_a^b \sqrt{1 + \left(\frac{df}{dx}\right)^2} dx$$

لاحظ انه هذا القانون هو حالة خاصة من القانون الجديد تكون فيه المعادلات (بارامترية)

$$x = t, \quad y = f(t), \quad a < t < b$$

$$\Rightarrow \frac{dx}{dt} = 1, \quad dx = dt, \quad \frac{dy}{dt} = f'(t) \quad \text{and} \quad \frac{dy}{dx} = \frac{dy/dt}{dx/dt} = \frac{dy}{dt}$$

$$\Rightarrow L = \int_a^b \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} dt = \int_a^b \sqrt{1 + \left(\frac{dy}{dx}\right)^2} dx$$

Area of Surfaces of Revolution

Area of Surface of Revolution for Parametrized Curves

If a smooth curve $x = f(t), y = g(t), a \leq t \leq b$, is traversed exactly once as t increases from a to b , then the areas of the surfaces generated by revolving the curve about the coordinate axes are as follows.

1. Revolution about the x-axis ($y \geq 0$):

$$S = \int_a^b 2\pi y \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} dt \quad (5)$$

2. Revolution about the y-axis ($x \geq 0$):

$$S = \int_a^b 2\pi x \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} dt \quad (6)$$

Example: Find the area of the surface generated by revolving the circle

$$x = \cos t, \quad y = 1 + \sin t, \quad 0 \leq t \leq 2\pi$$

about x -axis.

(نقطه‌ای معادله پارامتری به معادله دایره مرکزها $(0, 1)$ و شعاع 1)

sol: $\left(\frac{dx}{dt}\right)^2 = (-\sin t)^2 = \sin^2 t$



$$\left(\frac{dy}{dt}\right)^2 = \cos^2 t$$

$$\therefore S = \int_0^{2\pi} 2\pi y \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2} dt$$

$$2\pi \int_0^{2\pi} (1 + \sin t) \sqrt{\sin^2 t + \cos^2 t} dt$$

$$= 2\pi \int_0^{2\pi} 1 + \sin t dt = 2\pi [t - \cos t]_0^{2\pi}$$

$$= 2\pi [(2\pi - 1) - (0 - 1)] = \boxed{4\pi^2}$$



rest

1) Find the tangent to the curve

$$x = \frac{1}{t+1}, \quad y = \frac{t}{t-1}, \quad \text{at } t_0 = 2.$$

sol: At $t_0 = 2$, $(x_0, y_0) = (\frac{1}{3}, 2)$.

$$\text{Now, } m_T = \left. \frac{dy}{dx} \right|_{t=2} = \frac{dy/dt}{dx/dt} = \frac{(t-1-t)/(t-1)^2}{-1/(t+1)^2} = \frac{(t+1)^2}{(t-1)^2} \Big|_{t=2} = 9$$

Tangent line: $y = y_0 + m_T(x - x_0)$

$$= 2 + 9(x - \frac{1}{3}) \Rightarrow \boxed{y = 9x - 1}$$

Note that

$$\begin{aligned} \frac{d^2y}{dx^2} &= \frac{\left(\frac{dy}{dx}\right)'}{\left(\frac{dx}{dt}\right)'} = \frac{\left[\frac{d}{dt} \left(\frac{(t+1)^2}{(t-1)^2} \right)\right]}{-1/(t+1)^2} \\ &= \frac{4(t+1)^3}{(t-1)^3} \end{aligned}$$

2) Find the area of the surface generated by revolving the curve:

$x = \ln(\sec t + \tan t) - \sin t$, $y = \cos t$, $0 \leq t \leq \frac{\pi}{3}$
about x -axis.

sol: $\frac{dx}{dt} = \frac{\sin^2 t}{\cos t}$ (Do it) and $\frac{dy}{dt} = -\sin t$

$$\therefore S = 2\pi \int_0^{\frac{\pi}{3}} \cos t \sqrt{\frac{\sin^4 t}{\cos^2 t} + \sin^2 t} dt$$

$$= 2\pi \int_0^{\frac{\pi}{3}} \cos t \sqrt{\frac{\sin^4 t + \sin^2 t \cos^2 t}{\cos^2 t}} dt = 2\pi \int_0^{\frac{\pi}{3}} \cos t \frac{\sin t}{\cos t} dt$$

$$= -2\pi \cos t \Big|_0^{\pi/3} = -2\pi \left(\frac{1}{2} - 1\right) = \pi$$

in book

EXAMPLE 3 Find the area enclosed by the astroid (Figure 11.13)

$$x = \cos^3 t, \quad y = \sin^3 t, \quad 0 \leq t \leq 2\pi.$$

Solution By symmetry, the enclosed area is 4 times the area beneath the curve in the first quadrant where $0 \leq t \leq \pi/2$. We can apply the definite integral formula for area studied in Chapter 5, using substitution to express the curve and differential dx in terms of the parameter t . So,

$$A = 4 \int_0^1 y \, dx = 4 \int_{\pi/2}^0 \sin^3 t \cdot (-3\cos^2 t \sin t) \, dt$$

$$= 4 \int_0^{\pi/2} \sin^3 t \cdot 3 \cos^2 t \sin t \, dt$$

$$= 12 \int_0^{\pi/2} \left(\frac{1 - \cos 2t}{2}\right)^2 \left(\frac{1 + \cos 2t}{2}\right) \, dt$$

$$= \frac{3}{2} \int_0^{\pi/2} (1 - 2 \cos 2t + \cos^2 2t)(1 + \cos 2t) \, dt$$


$$= \frac{3}{2} \int_0^{\pi/2} (1 - \cos 2t - \cos^2 2t + \cos^3 2t) \, dt$$

$$= \frac{3}{2} \left[\int_0^{\pi/2} (1 - \cos 2t) \, dt - \int_0^{\pi/2} \cos^2 2t \, dt + \int_0^{\pi/2} \cos^3 2t \, dt \right]$$

$$= \frac{3}{2} \left[\left(t - \frac{1}{2} \sin 2t\right) - \frac{1}{2} \left(t + \frac{1}{4} \sin 2t\right) + \frac{1}{2} \left(\sin 2t - \frac{1}{3} \sin^3 2t\right) \right]_{\pi/2}^0$$

$$= \frac{3}{2} \left[\left(\frac{\pi}{2} - 0 - 0 - 0\right) - \frac{1}{2} \left(\frac{\pi}{2} + 0 - 0 - 0\right) + \frac{1}{2} (0 - 0 - 0 + 0) \right]$$

$$= \frac{3\pi}{8}.$$



$$\sin^4 t = \left(\frac{1 - \cos 2t}{2}\right)^2$$

Expand square term.

Multiply terms.

Section 8.2,
Example 3

Evaluate.

