


Chapter 4:



Q1) (a) $y = 1 - (x+1)^3$

* $y = 1 - (x+1)^3 \Rightarrow y' = -3(x+1)^2$
 $y' = \text{zero} \Rightarrow \text{zero} = -3(x+1)^2 \Rightarrow x = -1$



* $y' = -6(x+1) \Rightarrow y'' = \text{zero} \Rightarrow \text{zero} = -6(x+1)$
 $x = -1$

- $f(x)$ concave up on $(-\infty, -1]$
- $f(x)$ concave down on $[-1, \infty)$
- In inflection points $f(-1) = 1$.




(b) $y = \frac{x^2 + 1}{x} \quad x \neq 0$

$y = x + \frac{1}{x}$


$y' = 1 - \frac{1}{x^2} \Rightarrow \text{zero} = 1 - \frac{1}{x^2}$
 $1 = \frac{1}{x^2}$
 $x = -1, 1$

to help:-
 → No. H. asy
 → x = CI v. asy
 → y = x obliqu. asy




- $f(x)$ increasing on $(-\infty, 0)$
- $f(x)$ decreasing on $(0, \infty)$
- ~~f(x)~~ No extreme values.

$f''(x) = \frac{1}{x^3} \quad f'' \neq 0$



$\lim_{x \rightarrow 0^-} f(x) = -\infty$

$\lim_{x \rightarrow 0^+} f(x) = \infty$



c) $y = x^4 - 2x^2$

$y' = 4x^3 - 4x \Rightarrow y' = \text{zero}$

$\text{zero} = 4x^3 - 4x$

$\Rightarrow x^3 - x = 0$

$x(x^2 - 1) = 0 \Rightarrow x = 0, 1, -1$

$f(x)$ decreasing on $(-\infty, -1) \cup (0, 1)$

$f(x)$ increasing on $[-1, 0] \cup [1, \infty)$

$f(-1) = -1$ local minimum values

$f(0) = 0$ local maximum values & $f(1) = -1$ local minimum values

$y'' = 12x^2 - 4 \Rightarrow y'' = 0$

$0 = 12x^2 - 4 \Rightarrow 0 = 3x^2 - 1$

$\Rightarrow 3x^2 = 1 \Rightarrow x^2 = \frac{1}{3}$


$\Rightarrow x = \pm \sqrt{\frac{1}{3}}$

$f(x)$ concave up on $(-\infty, -\frac{1}{\sqrt{3}}) \cup (\frac{1}{\sqrt{3}}, \infty)$

$f(x)$ concave down on $(-\frac{1}{\sqrt{3}}, \frac{1}{\sqrt{3}})$

$f(-\frac{1}{\sqrt{3}}) = \frac{1}{9} - \frac{2}{3} = -\frac{5}{9}$ inflection points

$f(\frac{1}{\sqrt{3}}) = \frac{1}{9} - \frac{2}{3} = -\frac{5}{9}$ inflection points




d) $y = \frac{x^2 - 3}{x - 2}, x \neq 2$

$x=2$ v. asy
 $y=x+2$ oblique asy

$y' = \frac{x^2 - 4x + 3}{(x-2)^2}$ $y'' = \frac{2}{(x-2)^3}$




$y' = 0 \Rightarrow x = 1, 3 \rightarrow$ critical values 1, 3

y' DNE $\Rightarrow x = 2$ ($2 \notin D$)



$y'' = 0 \Rightarrow$ No values

y'' DNE $\Rightarrow x = 2$ ($2 \notin D$)



$x \neq 1, -1$
 $y = 0$ H.A.Sy
 $x = 1, x = -1$ V.Asy

f) $y = \frac{x}{x^2 - 1}$

$$y' = \frac{(x^2 - 1) - 2x^2}{(x^2 - 1)^2}$$

$$y' = \text{zero} \Rightarrow x^2 - 1 = 2x^2 \Rightarrow x^2 = -1$$


$f'(x)$ decreasing on $(-\infty, \infty)$

$$y' = \frac{1}{x^2 - 1} - \frac{2x^2}{(x^2 - 1)^2}$$

$$y'' = \frac{-2x}{(x^2 - 1)^2} - \left(\frac{(x^2 - 1)^2 4x - 2x^2 * 2x * 2(x^2 - 1)^2}{(x^2 - 1)^4} \right)$$

$$0 = -2x - 4x + 8x^3$$

$$-6x + 8x^3 = 0 \Rightarrow 2x(-3 + 4x^2) = 0$$



$$x = 0, -\frac{\sqrt{3}}{2}, \frac{\sqrt{3}}{2}$$

$f(x)$ concave up on $[-\frac{\sqrt{3}}{2}, 0] \cup [\frac{\sqrt{3}}{2}, \infty)$


$f(x)$ concave down on $(-\infty, -\frac{\sqrt{3}}{2}] \cup [0, \frac{\sqrt{3}}{2}]$

$(f(\frac{\sqrt{3}}{2}), \frac{\sqrt{3}}{2}) = (-\frac{\sqrt{3}}{2}, \frac{\sqrt{3}}{2})$

$(f(0), 0) = (0, 0)$

$(f(-\frac{\sqrt{3}}{2}), -\frac{\sqrt{3}}{2}) = (\frac{\sqrt{3}}{2}, -\frac{\sqrt{3}}{2})$

inflection points



g) $y = x\sqrt{8-x^2}$


$D \Rightarrow 8-x^2 \geq 0$
 $\sqrt{8} \geq x \geq -\sqrt{8}$

$y' = \sqrt{8-x^2} + x \frac{-2x}{2\sqrt{8-x^2}}$

$y' = \frac{8-x^2-2x^2}{\sqrt{8-x^2}} = \frac{8-2x^2}{\sqrt{8-x^2}}$


$y'=0 \Rightarrow x=2, -2$

- $f(x)$ increasing on $[-2, 2]$
- $f(x)$ decreasing on $[-\sqrt{8}, -2] \cup [2, \sqrt{8}]$



- $f(-2) = -4$ abs. min
- $f(2) = 4$ abs. max

$y'' = \frac{2x(x^2-12)}{(8-x^2)^{3/2}} \Rightarrow y''=0 \Rightarrow x=0 \rightarrow y=0$



- y concave up $[-\sqrt{2}, 0]$
- y concave down $[0, \sqrt{2}]$
- $(0, 0)$ inflection point

