

## 8.1 Integration by Parts:

$$\int f(x) g(x) dx$$

we let  $u = f(x)$   $dv = g(x) dx$  (The easiest to integrate)

$$\Rightarrow du = f'(x) dx \quad \longrightarrow \quad v = \int g(x) dx = h(x)$$

then  $\int f(x) g(x) dx = f(x) h(x) - \int h(x) f'(x) dx.$

Example:  $\int x \cos x dx$

Let  $u = x$   $dv = \cos x dx$

$$\Rightarrow du = dx \quad v = \sin x$$

then  $\int x \cos x dx = x \sin x - \int \sin x dx = x \sin x + \cos x + C$

Example:  $\int \ln x dx$

$$\left. \begin{array}{l} u = \ln x \\ du = \frac{1}{x} dx \end{array} \right\} \begin{array}{l} dv = dx \\ v = x \end{array} \Rightarrow \int \ln x = x \ln x - \int \frac{1}{x} \cdot x dx$$

$$\Rightarrow \int \ln x dx = x \ln x - x + C$$

Example:  $\int x^2 e^x dx$

$$u = x^2 \quad dv = e^x dx$$

$$du = 2x dx \quad v = e^x$$

$$\Rightarrow \int x^2 e^x dx = x^2 e^x - \int 2x e^x dx$$

Again let  $u = x$   $dv = e^x dx$   
 $du = dx$   $v = e^x$

$$\begin{aligned} \Rightarrow \int x^2 e^x dx &= x^2 e^x - x e^x + \int e^x dx \\ &= x^2 e^x - 2x e^x + 2e^x + C \end{aligned}$$

Example:  $\int e^x \cos x dx$

$$\text{Let } u = e^x, \quad dv = \cos x dx$$

$$du = e^x dx \quad v = \sin x$$

$$\Rightarrow \int e^x \cos x dx = e^x \sin x - \int e^x \sin x dx$$

Again let  $u = e^x$   $dv = \sin x dx$   
 $du = e^x dx$   $v = -\cos x$


$$\Rightarrow \int e^x \cos x dx = e^x \sin x + e^x \cos x - \int e^x \cos x dx + C$$

$$\Rightarrow 2 \int e^x \cos x dx = e^x \sin x + e^x \cos x + C$$

$$\Rightarrow \int e^x \cos x dx = \frac{1}{2} e^x \sin x + \frac{1}{2} e^x \cos x + C$$

## Tabular Integration:


We use this method to find  $\int x^n g(x) dx$  when  $n$  is large &  $g(x)$  is easy to Integrate. derivatives Integrations.




$$\Rightarrow \int x^n g(x) dx = x^n h(x) - n x^{n-1} (r(x)) + n(n-1) x^{n-2} s(x) \dots$$

Example:

$$\int x^2 e^x dx = x^2 e^x - 2x e^x + 2e^x + C$$




Example:  $\int x^3 \sin x dx = -x^3 \cos x + 3x^2 \sin x + 6x \cos x - 6 \sin x + C$



Example: Find the area of the region bounded

by the curve  $y = x e^{-x}$  and the  $x$ -axis,  $x=0$  or  $x=4$

$$A = \int_0^4 x e^{-x} dx.$$



$$\text{Let } u = x$$


$$dv = e^{-x} dx$$

$$du = dx$$

$$v = -e^{-x}$$

$$\Rightarrow A = -x e^{-x} \Big|_0^4 + \int_0^4 e^{-x} dx$$

$$= -4e^{-4} + -e^{-x} \Big|_0^4 = -4e^{-4} - e^{-4} + 1 = 1 - 5e^{-4}$$



We combine the products of the functions connected by the arrows according to the operation signs above the arrows to obtain

$$\int x^2 e^x dx = x^2 e^x - 2x e^x + 2e^x + C.$$

Compare this with the result in Example 3.

**EXAMPLE 8** Evaluate

$$\int x^3 \sin x dx.$$

**Solution** With  $f(x) = x^3$  and  $g(x) = \sin x$ , we list:

$f(x)$ and its derivatives		$g(x)$ and its integrals
$x^3$	(+)	$\sin x$
$3x^2$	(-)	$-\cos x$
$6x$	(+)	$-\sin x$
$6$	(-)	$\cos x$
$0$		$\sin x$

Again we combine the products of the functions connected by the arrows according to the operation signs above the arrows to obtain

$$\int x^3 \sin x dx = -x^3 \cos x + 3x^2 \sin x + 6x \cos x - 6 \sin x + C.$$

The Additional Exercises at the end of this chapter show how tabular integration can be used when neither function  $f$  nor  $g$  can be differentiated repeatedly to become zero.

### Exercises 8.1

#### Integration by Parts

Evaluate the integrals in Exercises 1–24 using integration by parts.

1.  $\int x \sin \frac{x}{2} dx$

2.  $\int_0^{\pi} \theta \cos \pi \theta d\theta$   $u = \theta, dv = \cos \pi \theta d\theta$

15.  $\int x^3 e^x dx$

16.  $\int p^4 e^{-p} dp$

3.  $\int t^2 \cos t dt$

4.  $\int_0^{\pi} x^2 \sin x dx$   $u = x, dv = \sin x dx$

17.  $\int (x^2 - 5x) e^x dx$

18.  $\int (r^2 + r + 1) e^r dr$

5.  $\int_1^2 x \ln x dx$

6.  $\int_1^e x^3 \ln x dx \rightarrow u = \ln x, dv = x^3 dx$

19.  $\int x^5 e^x dx$

20.  $\int t^2 e^{4t} dt$

7.  $\int x e^x dx$

8.  $\int x e^{3x} dx$

21.  $\int e^{\theta} \sin \theta d\theta$

22.  $\int e^{-y} \cos y dy$

9.  $\int x^2 e^{-x} dx$

10.  $\int (x^2 - 2x + 1) e^{2x} dx$

23.  $\int e^{2x} \cos 3x dx$

24.  $\int e^{-2x} \sin 2x dx$

11.  $\int \tan^{-1} y dy$

12.  $\int \sin^{-1} y dy$

#### Using Substitution

Evaluate the integrals in Exercises 25–30 by using a substitution prior to integration by parts.

13.  $\int x \sec^2 x dx$

14.  $\int 4x \sec^2 2x dx$

25.  $\int e^{\sqrt{3x+9}} dx$

26.  $\int_0^1 x \sqrt{1-x} dx$

11.  $u = \tan^{-1} y \Rightarrow du = \frac{1}{1+y^2} dy, \text{ \& } dv = dy, v = y$

$$= y \tan^{-1} y - \int \left( \frac{y}{1+y^2} \right) dy = y \tan^{-1} y - \frac{1}{2} \ln(1+y^2) + C$$

(28)  $u = \ln(x+n^2)$ ,  $dv = dx$   
 $du = \frac{1+2n}{x+n^2}$ ,  $v = x$

$\Rightarrow x \ln(x+n^2) - \int \frac{x+2nx}{x+n^2} dx = x \ln(x+n^2) - \int \frac{(2n+1)}{x+n^2} dx$   
 $= x \ln(x+n^2) - \int \frac{2(n+1)-1}{x+n^2} dx =$

$x \ln(x+n^2)$   
 $- 2x + \ln(x+n^2)$   
 $+ C$

442 Chapter 8: Techniques of Integration

27.  $\int_0^{\pi/3} x \tan^2 x dx$       28.  $\int \ln(x+x^2) dx$   
 29.  $\int \sin(\ln x) dx$       30.  $\int z(\ln z)^2 dz$

d. What pattern do you see? What is the area between the curve and the x-axis for


$\left(\frac{2n-1}{2}\right)\pi \leq x \leq \left(\frac{2n+1}{2}\right)\pi$ .

n an arbitrary positive integer? Give reasons for your answer.

**Evaluating Integrals**


Evaluate the integrals in Exercises 31–50. Some integrals do not require integration by parts.

31.  $\int x \sec x^2 dx$       32.  $\int \frac{\cos \sqrt{x}}{\sqrt{x}} dx$   
 33.  $\int x (\ln x)^2 dx$       34.  $\int \frac{1}{x (\ln x)^2} dx$   
 35.  $\int \frac{\ln x}{x^2} dx$       36.  $\int \frac{(\ln x)^3}{x} dx$   
 37.  $\int x^3 e^x dx$       38.  $\int x^3 e^{x^2} dx$   
 39.  $\int x^3 \sqrt{x^2+1} dx$       40.  $\int x^2 \sin x^3 dx$   
 41.  $\int \sin 3x \cos 2x dx$       42.  $\int \sin 2x \cos 4x dx$   
 43.  $\int e^x \sin e^x dx$       44.  $\int \frac{e^{\sqrt{x}}}{\sqrt{x}} dx$   
 45.  $\int \cos \sqrt{x} dx$       46.  $\int \sqrt{x} e^{\sqrt{x}} dx$   
 47.  $\int_0^{\pi/2} \theta^2 \sin 2\theta d\theta$       48.  $\int_0^{\pi/2} x^3 \cos 2x dx$   
 49.  $\int_{2\sqrt{3}}^2 t \sec^{-1} t dt$       50.  $\int_0^{1/\sqrt{2}} 2x \sin^{-1}(x^2) dx$



**Theory and Examples**

51. **Finding area** Find the area of the region enclosed by the curve  $y = x \sin x$  and the x-axis (see the accompanying figure) for  
 a.  $0 \leq x \leq \pi$ .  
 b.  $\pi \leq x \leq 2\pi$ .  
 c.  $2\pi \leq x \leq 3\pi$ .  
 d. What pattern do you see here? What is the area between the curve and the x-axis for  $n\pi \leq x \leq (n+1)\pi$ , n an arbitrary nonnegative integer? Give reasons for your answer.



52. **Finding area** Find the area of the region enclosed by the curve  $y = x \cos x$  and the x-axis (see the accompanying figure) for  
 a.  $\pi/2 \leq x \leq 3\pi/2$ .  
 b.  $3\pi/2 \leq x \leq 5\pi/2$ .  
 c.  $5\pi/2 \leq x \leq 7\pi/2$ .

53. **Finding volume** Find the volume of the solid generated by revolving the region in the first quadrant bounded by the coordinate axes, the curve  $y = e^x$ , and the line  $x = \ln 2$  about the line  $x = \ln 2$ .  
 54. **Finding volume** Find the volume of the solid generated by revolving the region in the first quadrant bounded by the coordinate axes, the curve  $y = e^{-x}$ , and the line  $x = 1$   
 a. about the y-axis.  
 b. about the line  $x = 1$ .  
 55. **Finding volume** Find the volume of the solid generated by revolving the region in the first quadrant bounded by the coordinate axes and the curve  $y = \cos x$ ,  $0 \leq x \leq \pi/2$ , about  
 a. the y-axis.  
 b. the line  $x = \pi/2$ .  
 56. **Finding volume** Find the volume of the solid generated by revolving the region bounded by the x-axis and the curve  $y = x \sin x$ ,  $0 \leq x \leq \pi$ , about  
 a. the y-axis.  
 b. the line  $x = \pi$ .  
 (See Exercise 51 for a graph.)  
 57. Consider the region bounded by the graphs of  $y = \ln x$ ,  $y = 0$ , and  $x = e$ .  
 a. Find the area of the region.  
 b. Find the volume of the solid formed by revolving this region about the x-axis.  
 c. Find the volume of the solid formed by revolving this region about the line  $x = -2$ .  
 d. Find the centroid of the region.  
 58. Consider the region bounded by the graphs of  $y = \tan^{-1} x$ ,  $y = 0$ , and  $x = 1$ .  
 a. Find the area of the region.  
 b. Find the volume of the solid formed by revolving this region about the y-axis.  
 59. **Average value** A retarding force, symbolized by the dashpot in the accompanying figure, slows the motion of the weighted spring so that the mass's position at time  $t$  is

$y = 2e^{-t} \cos t, \quad t \geq 0.$

(33)  $u = \ln x \Rightarrow \begin{cases} x = e^u \\ du = \frac{1}{x} dx \\ dx = e^u du \end{cases}$   
 $\Rightarrow \int e^{2u} u^2 du \Rightarrow$   
 $= \frac{1}{2} u^2 e^u - \frac{2u}{1} e^u + \frac{2e^u}{1} + C$

$u^2 \cdot e^{2u}$   
 $2u \cdot e^{2u} \cdot 2e^u$   
 $2 \cdot e^{2u} \cdot 2e^u$   
 $0 \cdot e^{2u} \cdot 2e^u$

(39)  $u = x^2 \Rightarrow du = 2x dx$   
 $dv = \sqrt{x^2+1} \cdot x dx$   
 $v = \frac{1}{3} (x^2+1)^{3/2}$   
 by parts

$$(39) \quad u = x^2$$

$$du = 2x dx$$

$$dv = \sqrt{x^2+1} \cdot x dx$$

$$v = \frac{1}{3} (x^2+1)^{3/2}$$

$$\Rightarrow \frac{1}{3} x^2 (x^2+1)^{3/2} - \int \frac{1}{3} (x^2+1)^{3/2} \cdot 2x dx$$

$$\frac{1}{3} x^2 (x^2+1)^{3/2} - \frac{2}{15} (x^2+1)^{5/2} + C$$

where  $\frac{d}{dx} \left( \frac{2}{15} (x^2+1)^{5/2} \right) = \frac{2}{15} \cdot \frac{5}{2} (2x) \cdot (x^2+1)^{3/2}$

$$(46) \quad \int \sqrt{x} e^{\sqrt{x}} dx$$

Let  $u = \sqrt{x}$

$$du = \frac{1}{2\sqrt{x}} dx$$

$$\Rightarrow dx = 2u du$$

$$\Rightarrow \int u e^u \cdot (2u du) = \int 2u^2 e^u du = 2[u^2 e^u - 2u e^u + 2e^u] + C$$
$$= 2[\sqrt{x} e^{\sqrt{x}} - 2\sqrt{x} e^{\sqrt{x}} + 2e^{\sqrt{x}}] + C$$

$u^2$		$e^u$
	$\circledast$	
$2u$		$e^u$
	$\circledast$	
$2$		$e^u$
	$\circledast$	
$0$		$e^u$

## 8.2 Trigonometric Integrals:

The purpose is to find  $\int \sin^m x \cos^n x dx$ :

Case 1 If  $m$  is odd, we write  $m = 2k+1$

then use  $\sin^2 x = 1 - \cos^2 x$

$$\rightarrow \sin^m x = (\sin^2 x)^k \sin x = (1 - \cos^2 x)^k \sin x.$$

Then we write  $\sin x dx$  as  $-d(\cos x)$

Example: Evaluate  $\int \sin^3 x \cos^2 x dx$ .

$$\sin^3 x = \sin^2 x \cdot \sin x = (1 - \cos^2 x) \sin x$$

$$\rightarrow \int \sin^3 x \cos^2 x dx = \int (1 - \cos^2 x) \cos^2 x \sin x dx$$

$$= \int (1 - \cos^2 x) \cos^2 x (-d(\cos x))$$

$$= \int \cos^2 x - \cos^4 x (-d \cos x) =$$

Let  $u = \cos x$

$$= - \int u^2 - u^4 du$$

$$= - \frac{u^3}{3} + \frac{u^5}{5} + C = - \frac{\cos^3 x}{3} + \frac{\cos^5 x}{5} + C$$

(1)



Case 2: If  $m$  is even &  $n$  is odd

we write  $n = 2k+1$  & use:

$$\cos^2 x = 1 - \sin^2 x$$

$$\Rightarrow \cos^n x = (\cos^2 x)^k \cos x = (1 - \sin^2 x)^k \cos x$$

then use:  $\cos x dx \rightarrow d(\sin x)$ .

Example: Evaluate  $\int \cos^5 x dx$ .

here  $m=0$  (even) &  $n=5$  (odd)

$$\Rightarrow \cos^5 x = (\cos^2 x)^2 \cos x$$

$$\Rightarrow \int \cos^5 x dx = \int (1 - \sin^2 x)^2 \cos x dx$$

$$= \int (1 - \sin^2 x)^2 d \sin x$$

Let  $u = \sin x$

$$= \int (1 - u^2)^2 du = \int 1 - 2u^2 + u^4 du$$

$$= u - \frac{2u^3}{3} + \frac{u^5}{5} + C$$

$$= \sin x - \frac{2 \sin^3 x}{3} + \frac{\sin^5 x}{5} + C$$

Case 3 If both  $m$  &  $n$  are even

then:  $\sin^2 x = \frac{1 - \cos 2x}{2}$ ,  $\cos^2 x = \frac{1 + \cos 2x}{2}$

Example:  $\int \sin^2 x \cos^4 x dx$ .

$$\begin{aligned} \int \sin^2 x \cos^4 x dx &= \int \left( \frac{1 - \cos 2x}{2} \right) \left( \frac{1 + \cos 2x}{2} \right)^2 dx \\ &= \frac{1}{8} \int (1 - \cos 2x)(1 + 2\cos 2x + \cos^2 2x) dx \\ &= \frac{1}{8} \int (1 + \cos 2x - \cos^2 2x - \cos^3 2x) dx \\ &= \frac{1}{8} \left[ x + \frac{\sin 2x}{2} - \int (\cos^2 2x + \cos^3 2x) dx \right] \end{aligned}$$

• Need to Evaluate  $\int \cos^2 2x dx$  +  $\int \cos^3 2x dx$

□  $\int \cos^2 2x dx = \int \frac{1 + \cos 4x}{2} dx = \frac{1}{2} \left( x + \frac{\sin 4x}{4} \right)$

\*  $\int \cos^3 2x dx$ ,  $n$  odd.

□  $\int \cos^3 2x dx = \int (1 - \sin^2 2x) \cos 2x dx$   $u = \sin 2x$   
 $du = 2\cos 2x dx$

$$= \int (1 - u^2) \cdot \frac{1}{2} du = \frac{1}{2} \left( u - \frac{u^3}{3} \right) = \frac{1}{2} \left( \sin 2x - \frac{\sin^3 2x}{3} \right)$$

then The whole Integration is:

$$\begin{aligned} &= \frac{1}{8} \left[ x + \frac{\sin 2x}{2} - \frac{1}{2} x - \frac{1}{8} \sin 4x - \frac{1}{2} \sin 2x + \frac{\sin^3 2x}{6} \right] + C \\ &= \frac{1}{16} \left[ x - \frac{1}{4} \sin 4x + \frac{1}{3} \sin^3 2x \right] + C \end{aligned}$$

## Eliminating Square Roots:

Example: Evaluate  $\int_0^{\frac{\pi}{4}} \sqrt{1 + \cos 4x} \, dx$

Recall:  $\cos 2x = 2\cos^2 x - 1 \Rightarrow \cos^2 x = \frac{\cos 2x + 1}{2}$

$\Rightarrow 2\cos^2 2x = \cos 4x + 1$  , So we can substitute.

$$\int_0^{\frac{\pi}{4}} \sqrt{1 + \cos 4x} \, dx = \int_0^{\frac{\pi}{4}} \sqrt{2} \sqrt{\cos^2 2x} \, dx = \int_0^{\frac{\pi}{4}} \sqrt{2} \underbrace{|\cos 2x|}_{\geq 0 \text{ on } [0, \frac{\pi}{4}]}} \, dx$$

$$= \int_0^{\frac{\pi}{4}} \sqrt{2} \cos 2x \, dx = \sqrt{2} \left[ \frac{\sin 2x}{2} \right]_0^{\frac{\pi}{4}} = \frac{\sqrt{2}}{2}$$

## Integrals of Powers of $\tan x$ & $\sec x$ :

Example: Evaluate  $\int \tan^4 x \, dx$ .

$$\int \tan^4 x \, dx = \int \tan^2 x \underbrace{\tan^2 x}_{\sec^2 x - 1} \, dx = \int \tan^2 x (\sec^2 x - 1) \, dx$$

$$= \int \tan^2 x \sec^2 x \, dx - \int \tan^2 x \, dx$$

$$= \int \tan^2 x \sec^2 x \, dx - \int (\sec^2 x - 1) \, dx$$

$$= \int \tan^2 x \sec^2 x \, dx - \int \sec^2 x \, dx + \int 1 \, dx$$

↓  
by parts

(4)

$$\int \tan^2 x \sec^2 x \, dx:$$

$$\text{Let } u = \tan x \quad du = \sec^2 x \, dx$$

$$\text{then: } \int u^2 \, du = \frac{1}{3} u^3 + C \quad \text{u's rule}$$

then: The whole Integral becomes:

$$\int \tan^4 x \, dx = \frac{1}{3} \tan^3 x - \tan x + x + C$$

Example: Evaluate  $\int \sec^3 x \, dx$

Integrate by parts:

$$\begin{aligned} u &= \sec x & dv &= \sec^2 x \, dx \\ du &= \sec x \tan x \, dx & v &= \tan x \end{aligned}$$

$$\begin{aligned} \Rightarrow \int \sec^3 x \, dx &= \sec x \tan x - \int \tan^2 x \sec x \, dx \\ &= \sec x \tan x - \int (\sec^2 x - 1) \sec x \, dx \\ &= \sec x \tan x - \int \sec^3 x \, dx + \int \sec x \, dx \end{aligned}$$

$$\Rightarrow 2 \int \sec^3 x \, dx = \sec x \tan x + \int \sec x \, dx$$

$$\Rightarrow \int \sec^3 x \, dx = \frac{\sec x \tan x}{2} + \frac{1}{2} \ln |\sec x + \tan x| + C$$

## Products of Sines and Cosines:

$$\sin m x \sin n x = \frac{1}{2} [\cos(m-n)x - \cos(m+n)x],$$

$$\sin m x \cos n x = \frac{1}{2} [\sin(m-n)x + \sin(m+n)x],$$

$$\cos m x \cos n x = \frac{1}{2} [\cos(m-n)x + \cos(m+n)x].$$

Example: Evaluate  $\int \sin 3x \cos 5x \, dx$ .

$$\int \sin 3x \cos 5x \, dx = \frac{1}{2} \int [\sin(-2x) + \sin(8x)] \, dx$$

$$= \frac{1}{2} \left[ + \frac{\cos 2x}{2} - \frac{\cos 8x}{8} \right] + C$$

$$= \frac{\cos 2x}{4} - \frac{\cos 8x}{16} + C$$

$$(8.2) (19) \int 16 \sin^2 x \cos^2 x dx$$

$$= \int 16 \left( \frac{1 - \cos 2x}{2} \right) \left( \frac{1 + \cos 2x}{2} \right) dx$$

$$= \int 4 (1 - \cos^2 2x) dx = 4 \int \left( 1 - \left[ \frac{1 + \cos 4x}{2} \right] \right) dx$$

$$= 4x - 2x - \frac{1}{2} \sin 4x + C = 2x - \frac{1}{2} \sin 4x + C$$

$$(34) \int \sec x \tan^2 x dx = \int \sec x \tan x \tan x dx$$

Let

$$\begin{array}{ll} u = \tan x & dv = \sec x \tan x dx \\ du = \sec^2 x dx & v = \sec x \end{array}$$

$$\begin{aligned} \Rightarrow &= \tan x \sec x - \int \sec^3 x dx \\ &= \tan x \sec x - \int \sec^2 x \sec x dx \\ &= \tan x \sec x - \int (\tan^2 x + 1) \sec x dx \\ &= \tan x \sec x - \int \tan^2 x \sec x dx + \int \sec x dx \\ &= \sec x \tan x - \ln |\sec x + \tan x| - \int \tan^2 x \sec x dx \end{aligned}$$

$$\Rightarrow 2 \int \sec x \tan^2 x dx = \sec x \tan x - \ln |\sec x + \tan x| + C$$

$$\Rightarrow \int \sec x \tan^2 x dx = \frac{1}{2} \sec x \tan x - \frac{1}{2} \ln |\sec x + \tan x| + C$$

$$\begin{aligned}
 (52) \quad & \int \sin 2x \cos 3x \, dx \\
 &= \frac{1}{2} \int (\underbrace{\sin(-x)}_{\text{odd}} + \sin 5x) \, dx = \frac{1}{2} \cos x - \frac{1}{10} \cos 5x + C
 \end{aligned}$$

$$\begin{aligned}
 (64) \quad & \int \frac{\sin^3 x}{\cos^4 x} \, dx = \int \frac{\sin^2 x \sin x}{\cos^4 x} \, dx \\
 &= \int \frac{(1 - \cos^2 x) \sin x}{\cos^4 x} \, dx = \int \frac{\sin x}{\cos^4 x} \, dx - \int \frac{\cos^2 x \sin x}{\cos^4 x} \, dx
 \end{aligned}$$

$$\begin{aligned}
 \textcircled{35} \quad &= \int \sec^3 x \tan x \, dx - \int \sec x \tan x \, dx \\
 &= \int \sec^2 x \sec x \tan x \, dx - \int \sec x \tan x \, dx
 \end{aligned}$$

$$= \frac{1}{3} \sec^3 x - \sec x + C$$

$\textcircled{35}$  Let  $u = \sec x$   
 $du = \sec x \tan x \, dx$

$$(67) \quad \int x \sin^2 x \, dx = \int x \left( \frac{1 - \cos 2x}{2} \right) dx$$

$$= \frac{1}{2} \int x \, dx - \frac{1}{2} \int x \cos 2x \, dx \quad \left[ \begin{array}{l} u = x \\ du = dx \end{array} \right. \quad \begin{array}{l} dv = \cos 2x \, dx \\ v = \frac{1}{2} \sin 2x \end{array}$$

$$= \frac{1}{2} \frac{x^2}{2} - \frac{1}{2} \left[ \frac{1}{2} x \sin 2x - \frac{1}{2} \int \sin 2x \, dx \right]$$

$$= \frac{1}{4} x^2 - \frac{1}{4} x \sin 2x - \frac{1}{8} \cos 2x + C$$