


2.5 Continuity

36

- * Points are 3 kinds:
- ① interior points (c)
 - ② left endpoints (a)
 - ③ right endpoints (b)








Definition: (Continuity at point)

A function f is continuous at an interior point $x=c$ of its domain if

$$\lim_{x \rightarrow c} f(x) = f(c)$$

Example: Discuss the continuity at $x=0$ to the following functions



f is continuous at $x=0$ because $\lim_{x \rightarrow 0} f(x) = f(0) = 1$

g is not continuous at $x=0$ because $\lim_{x \rightarrow 0} g(x) = 1 \neq g(0) \rightarrow \text{DNE}$

h is not continuous at $x=0$ because $\lim_{x \rightarrow 0} h(x) = 1 \neq h(0) = 2$

l is not continuous at $x=0$ because $1 = f(0) \neq \lim_{x \rightarrow 0} l(x) \rightarrow \text{DNE}$

$y = \frac{1}{x^2}$ is not continuous at $x=0$ because $\lim_{x \rightarrow 0} \frac{1}{x^2} = \infty$ and $f(0)$ is not defined.

(b) and (c) are called removable discontinuity because in

(b) g would be continuous if $g(0) = 1$


(c) h would be continuous if $h(0) = 1$ instead of 2

(d) is called jump discontinuity.

(e) is called infinite discontinuity.

Definition: A function f is

- continuous at left endpoint $x=a$ of its domain if $\lim_{x \rightarrow a^+} f(x) = f(a)$
- continuous at right endpoint $x=b$ of its domain if $\lim_{x \rightarrow b^-} f(x) = f(b)$




Example: $f(x) = \sqrt{4-x^2}$ is continuous at every point of its domain $[-2, 2]$


→ f is continuous at -2 because $\lim_{x \rightarrow -2^+} \sqrt{4-x^2} = f(-2) = 0$

→ f is continuous at 2 because $\lim_{x \rightarrow 2^-} \sqrt{4-x^2} = f(2) = 0$

→ f is continuous on an interval $[-2, 2]$.



Example: $f(x) = \begin{cases} 1, & x \geq 0 \\ 0, & x < 0 \end{cases}$



• f is right continuous at $x=0$ because

$$\lim_{x \rightarrow 0^+} f(x) = f(0) = 1$$

• f is not left continuous at $x=0$ because $\lim_{x \rightarrow 0^-} f(x) = 0 \neq f(0) = 1$

• f is not continuous at $x=0$ because $\lim_{x \rightarrow 0} f(x) \neq f(0) = 1$

DNE


Test of Continuity:

$f(x)$ is continuous at $x=c$ ^{an interior point} iff the following three conditions hold:

- 1) $f(c)$ exists where $c \in D(f)$ domain of f
- 2) $\lim_{x \rightarrow c} f(x)$ exists
- 3) $\lim_{x \rightarrow c} f(x) = f(c)$

Example: Discuss the continuity of f at $x = 0, 1, 2, 3, 4$, where f is as given in the graph defined over the domain $[0, 4]$.

(38)



(a) f is continuous at $x = 0$ because

$$\lim_{x \rightarrow 0^+} f(x) = f(0) = 1 \quad \left(\begin{array}{l} \text{right-hand limit exists} \\ \text{at the left endpoint} \end{array} \right)$$

(b) f is discontinuous at $x = 1$ because

$$\lim_{x \rightarrow 1} f(x) \text{ DNE}$$

(c) f is discontinuous at $x = 2$ because $\lim_{x \rightarrow 2} f(x) = 1 \neq f(2) = 2$

(d) f is continuous at $x = 3$ because $\lim_{x \rightarrow 3} f(x) = f(3) = 2$

(e) f is discontinuous at $x = 4$ because $\lim_{x \rightarrow 4^-} f(x) = 1 \neq f(4) = \frac{1}{2}$ (left-hand limit exists at the right endpoint).

Theorem: If the functions f and g are continuous at $x = c$, then the following functions are continuous at $x = c$:

1) $f \pm g$ 2) fg 3) Kf , where $K \in \mathbb{R}$

4) $\frac{f}{g}$ where $g(c) \neq 0$

5) $[f(x)]^{\frac{m}{n}}$ where $(f(x))^{\frac{m}{n}}$ is defined on an interval containing c , and $m, n \in \mathbb{Z}$

Example:
 • Every polynomial is continuous at every point of the real line.
 • Every rational function is continuous at every point where it's defined (denominator is different from zero).


Example: $f(x) = x^3 - 2x^2 + 1$ is continuous at every point x .

$$g(x) = \frac{f(x)}{x^2 - 4} = \frac{x^3 - 2x^2 + 1}{(x-2)(x+2)}$$

x except $x = 2$ and $x = -2$ where the denominator is zero.

Example $f(x) = |x|$ is continuous

$$f(x) = \begin{cases} x & \text{if } x \geq 0 \\ -x & \text{if } x < 0 \end{cases}$$



(39)

if $x > 0$ then $f(x) = x$ polynomial which is continuous

if $x < 0$ then $f(x) = -x$ polynomial which is continuous

if $x = 0$ then $\lim_{x \rightarrow 0} f(x) = \lim_{x \rightarrow 0} |x| = f(0) = 0$

Continuity of trigonometric functions:

* The functions $\sin x$ and $\cos x$ are continuous at every value x .


* The functions $\tan x = \frac{\sin x}{\cos x}$, $\cot x = \frac{\cos x}{\sin x} = \frac{1}{\tan x}$, $\sec x = \frac{1}{\cos x}$ and $\csc x = \frac{1}{\sin x}$

are continuous at every point except where they are not defined.




Theorem (Continuity of Composition)

If f is continuous at c and g is continuous at $f(c)$, then

$g \circ f$ is continuous at c .



Example: Let $f(x) = \sqrt{x}$ and $g(x) = x^2 - 1$ show that $g \circ f$ is continuous at $x = 4$



1. $f(x)$ is continuous at $x = 4$ because

$$\lim_{x \rightarrow 4} \sqrt{x} = f(4) = 2 \quad \text{and}$$

$g(x)$ is continuous at $x = f(4) = 2$ because

$$\lim_{x \rightarrow 2} (x^2 - 1) = g(2) = 3.$$

Thus by Theorem above $g \circ f$ is continuous at $x = 4$.

2. Note that $(g \circ f)(x) = g(f(x)) = g(\sqrt{x}) = (\sqrt{x})^2 - 1 = x - 1$.

This is polynomial and continuous everywhere. Thus continuous at $x = 4$.

Continuous Extension to point

(40)

* A rational function f may have a limit L at point $x=c$ even if $f(c)$ is not defined (the denominator is zero).

Example: $f(x) = \frac{x^2 - 4}{x - 2}$

• If $x=2 \Rightarrow f(2)$ is not defined but

• If $x \neq 2 \Rightarrow f(x) = \frac{x^2 - 4}{x - 2} = \frac{(x-2)(x+2)}{(x-2)} = x+2$

The function $F(x) = x+2$ is the same as $f(x) = \frac{x^2 - 4}{x - 2}$ for all $x \neq 2$

The only difference is that $F(x)$ is continuous at $x=2$ because

$$\lim_{x \rightarrow 2} F(x) = \lim_{x \rightarrow 2} (x+2) = 4 = F(2)$$

but $f(x)$ is not continuous at $x=2$ because

$$\lim_{x \rightarrow 2} f(x) = 4 \neq f(2)$$

Thus, $F(x)$ is called the continuous extension of $f(x)$ at $x=c$, and we write

$$F(x) = \begin{cases} f(x) & \text{if } x \neq c \text{ and } x \in D(f) \\ L & \text{if } x = c \end{cases}$$

where $\lim_{x \rightarrow c} f(x) = L$.

Example: show that $f(x) = \frac{x^2 + x - 6}{x^2 - 4}$ has a continuous extension at $x=2$, and find that extension.

• If $x=2 \Rightarrow f(2)$ is not defined.

• If $x \neq 2 \Rightarrow f(x) = \frac{x^2 + x - 6}{x^2 - 4} = \frac{(x-2)(x+3)}{(x-2)(x+2)} = \frac{x+3}{x+2}$

The function $F(x) = \frac{x+3}{x+2}$ is the same as $f(x)$ for all $x \neq 2$

⇒ But $F(x)$ is continuous at $x=2$ because



(41)

$$\lim_{x \rightarrow 2} F(x) = \lim_{x \rightarrow 2} \frac{x+3}{x+2} = \frac{5}{4} = F(2)$$

and $f(x)$ is not continuous at $x=2$ because

$$\lim_{x \rightarrow 2} f(x) = \lim_{x \rightarrow 2} \frac{x^2+x-6}{x^2-4} = \lim_{x \rightarrow 2} \frac{x+3}{x+2} = \frac{5}{4} \neq f(2)$$

Thus, F is the continuous extension of f to $x=2$. $F(x) = \begin{cases} \frac{x^2+x-6}{x^2-4}, & x \neq 2 \\ \frac{5}{4}, & x = 2 \end{cases}$



Continuity on Intervals

• Let $D(f)$ be the domain of the function f :

→ A function f is continuous if it is continuous ^{at} everywhere in $D(f)$.





→ A function f is continuous on an interval $I \subset D(f)$ if f is continuous at every point in I .

• → If the function f is continuous on an interval I , then f is continuous on any interval $J \subset I$.

Example! → Polynomials _{functions} are continuous on every interval.

→ Rational functions are continuous on every interval on which they are defined.



Example*



42

Theorem: Suppose $f(x)$ is continuous on an interval $I = [a, b]$.
The Intermediate Value Theorem (IVT): If y_0 is any number between $f(a)$ and $f(b)$, then there exists a number c between a and b such that $f(c) = y_0$.

The continuity of f on I is essential to IVT.
 If f is discontinuous at even one point of I , then IVT may fail



Consequences of the IVT

• IVT is the reason that the graph of a function continuous on an interval I can not have any breaks. It will be **connected** and **single unbroken curve**. For instance (a) and (d) in Example*. It will not have jumps like (c) in Example* or separate branches like (b) in Example*.

Theorem: (limits of continuous functions)

(43)

If g is continuous at the point b , and

$$\lim_{x \rightarrow c} f(x) = b, \text{ then } \lim_{x \rightarrow c} g(f(x)) = g(b)$$

$$= g\left(\lim_{x \rightarrow c} f(x)\right)$$

Example: $\lim_{x \rightarrow \frac{\pi}{2}} \cos\left(2x + \sin\left(\frac{3\pi}{2} + x\right)\right) = \cos\left(\lim_{x \rightarrow \frac{\pi}{2}} 2x + \lim_{x \rightarrow \frac{\pi}{2}} \sin\left(\frac{3\pi}{2} + x\right)\right)$

$$= \cos\left(\pi + \sin 2\pi\right)$$
$$= \cos \pi = -1$$

Example: Show that \exists a root of the equation $x^3 - x - 1 = 0$ between 1 and 2.

$$\text{let } f(x) = x^3 - x - 1$$

$$f(1) = 1 - 1 - 1 = -1 < 0$$

$$f(2) = 8 - 2 - 1 = 5 > 0$$

Since $0 = y_0$ is between $f(1)$ and $f(2)$

Since f is continuous (polynomial). Thus, by IVT, \exists

a (zero) root of f between 1 and 2.

$$x = 1.32$$