


# 4.4 Concavity and Curve Sketching (86)

Def Let  $f(x)=y$  be a differentiable function on interval  $I$

(a) If  $f'$  is increasing on  $I$ , then  $f$  is concave up on the open interval  $I$ .

(b) If  $f'$  is decreasing on  $I$ , then  $f$  is concave down on the open interval  $I$ .

Example  $y = x^3$




## The 2<sup>nd</sup> Derivative Test for Concavity:


\* Let  $y = f(x)$  be twice-differentiable on an interval  $I$ .

(a) If  $f'' > 0$  on  $I$ , then  $f$  is concave up on  $I$ .

(b) If  $f'' < 0$  on  $I$ , then  $f$  is concave down on  $I$ .

Example: Determine the concavity of

①  $y = x^2 \Rightarrow y' = 2x \Rightarrow y'' = 2 > 0$    $\Rightarrow y$  is concave up on every  $I$ .

②  $y = x^3 \Rightarrow y' = 3x^2 \Rightarrow y'' = 6x$  

$y$  is concave up on  $(0, \infty)$

$y$  is concave down on  $(-\infty, 0)$


③  $y = 3 + \sin x$  on  $[0, 2\pi]$

$y' = \cos x$

$y'' = -\sin x$

$y$  concave up on  $(\pi, 2\pi)$

$y$  concave down on  $(0, \pi)$



Def A point  $x=c$  is called inflection point of the function  $f$  if the function  $f$  has a tangent line at  $x=c$  and changes concavity.   
*May be horizontal or vertical or oblique*

If the function  $f$  has an inflection point at  $(c, f(c))$  then either  $f'(c) = 0$  or  $f''(c)$  fails to exist (undefined)


Example 1 ( $f'(c)$  exists but  $f''(c)$  fails to exist) Find the inflection point?

$$f(x) = x^{\frac{5}{3}}, \quad f'(x) = \frac{5}{3}x^{\frac{2}{3}}, \quad f''(x) = \frac{10}{9}x^{-\frac{1}{3}} = \frac{10}{9\sqrt[3]{x}}$$

at  $x=0$ , we have a tangent

since  $f'(0) = 0$  (exists)

②  $f$  changes concavity since



$f''(0)$  fails to exist

Thus,  $(0, f(0)) = (0, 0)$  is an inflection point.

Example 2 ( $f'(c)$  and  $f''(c)$  exist)

Find an inflection point of  $y = x^4$


$$y' = 4x^3, \quad y'' = 12x^2 = 0 \Leftrightarrow x = 0 \text{ so } f''(0) = 0$$

at  $x=0$ , we have a tangent

since  $f'(0) = 0$  (exists)

②  $f$  does not change concavity

since  $\begin{matrix} \text{up} & & \text{up} \\ \uparrow\uparrow\uparrow & & \uparrow\uparrow\uparrow \\ & 0 & \end{matrix} y''$



Thus,  $(0, f(0)) = (0, 0)$  is not inflection point

Example 3 ( $f'(c)$  and  $f''(c)$  do not exist) Find the inflection point of  $y = x^{\frac{1}{3}}$

$$y' = \frac{1}{3}x^{-\frac{2}{3}} = \frac{1}{3\sqrt[3]{x^2}}$$


$$y'' = -\frac{2}{9}x^{-\frac{5}{3}} = \frac{-2}{9\sqrt[3]{x^5}}$$

at  $x=0$ , we have a vertical tangent since  $f'(0)$  fails to exist.

②  $f$  changes concavity at  $x=0$  since

$\begin{matrix} \text{up} & & \text{down} \\ \uparrow\uparrow\uparrow & & \downarrow\downarrow\downarrow \\ & 0 & \end{matrix} y''$

Thus  $(0, 0)$  is inflection point



## The (2<sup>nd</sup> derivative Test for local extrem)

(88)

Suppose  $f'$  is continuous on an open interval that contain  $c$ .

- (1) If  $f'(c) = 0$  and  $f''(c) < 0$ , then  $f$  has local max at  $c$ .
- (2) If  $f'(c) = 0$  and  $f''(c) > 0$ , then  $f$  has local min at  $c$ .
- (3) If  $f'(c) = 0$  and  $f''(c) = 0$ , then the test fails, and the function  $f$  may have local max = local min, or neither. (see example (1), (2))

Example (a) Find the local extrem of  $f(x) = x^3 - 3x + 3$

$$f' = 3x^2 - 3 = 0 \Leftrightarrow x = \pm 1 \quad \text{(critical points)}$$


(1, f(1)) and (-1, f(-1))

$$f''(x) = 6x$$


- $f'(1) = 0$  and  $f''(1) = 6 > 0 \Rightarrow f$  has local min at  $x = 1$
- $f'(-1) = 0$  and  $f''(-1) = -6 < 0 \Rightarrow f$  has local max at  $x = -1$ .

(b) Graph the function  $f$

- $f(0) = 3$
- local Max at  $(-1, 5)$
- local Min at  $(1, 1)$



$$f' = 3x^2 - 3 = 0 \Leftrightarrow x^2 - 1 = 0$$



•  $f$  is increasing on  $(-\infty, -1)$  and  $(1, \infty)$

•  $f$  is decreasing on  $(-1, 1)$

$$f'' = 6x = 0$$

down
up  
---
+++  
o
+

at  $x = 0$  we have inflection point  $(0, 3)$  since  $f(0) = 3$  and so, we have a tangent and  $f$  changes concavity.

\* To graph a function  $y=f(x)$


89

- 1) Find the Domain of  $f(x)$ .
- 2) Find  $y'$  and the critical points.
- 3) Find where  $f$  is increasing and decreasing, local Max & local Min.
- 4) Find  $y''$  and the inflection points.
- 5) Find where  $f$  is concave up and concave down.
- 6) Find the asymptotes of  $f$  (horizontal and vertical)
- 7) Plot Key points:  $x$ -intercepts and  $y$ -intercepts.


Example: Sketch the graph  $f(x) = \frac{(x+1)^2}{1+x^2}$

1)  $D(f) = (-\infty, \infty)$


2)  $f'(x) = \frac{2(1-x^2)}{(1+x^2)^2} = 0 \iff x = \pm 1$ , so the critical points are  $(1, 2)$  and  $(-1, 0)$



- $f$  is increasing on  $(-1, 1)$
- $f$  is decreasing on  $(-\infty, -1)$  and  $(1, \infty)$
- $f$  has local min  $(-1, 0)$
- $f$  has local Max at  $(1, 2)$



4)  $f''(x) = \frac{4x(x^2-3)}{(1+x^2)^3} = 0 \iff x = 0, \sqrt{3}, -\sqrt{3}$  are inflection points because  $f$  changes concavity



6) No vertical asymptotes  
Horizontal asymptote at  $x=1$   
because  $\lim_{x \rightarrow \infty} f(x) = 1$

7)  $(-1, 0), (0, 1)$