


## CHAPTER 6

### Continuous Random Variable


- A continuous random variable  $X$  is a random variable that assume any value over an interval or union of intervals.
- The probability that  $X$  assumes a value in any interval lies in the range 0 to 1.
- The total probability of all the (mutually exclusive) intervals within which  $x$  can assume a value is 1.
- The probability that a continuous random variable  $X$  assumes a value within a certain interval is given by the area under the curve between the two limits of the interval.
- For a continuous probability distribution, the probability is always calculated for an interval.
- The probability that a continuous random variable  $x$  assumes a single value is always zero.
- $P(a \leq X \leq b) = P(a < X < b) =$  **Area under the curve** from  $x = a$  to  $x = b$ .



## 6.1 The Uniform Distribution

A **uniform random variable**  $X$  over an interval  $[a, b]$ , denoted by  $X = U [a, b]$  is a continuous random variable with the following probability (density) function:

$$f(x) = \begin{cases} \frac{1}{b-a} & a \leq x \leq b \\ 0 & \text{elsewhere} \end{cases}$$



Let  $X = U [a, b]$ , then

- $P(x_1 < X < x_2) = \frac{1}{b-a} \cdot (x_2 - x_1) = \frac{x_2 - x_1}{b-a}$
- $E(X) = \frac{a+b}{2}$
- $\text{Var}(X) = \frac{(b-a)^2}{12}$
- $\sigma(X) = \frac{b-a}{\sqrt{12}}$

### Example

The time needed to finish STAT 2361 final exam is uniformly distributed between 60 and 100 minutes

1. Write the probability function for this distribution.

$$X = U [60, 100]$$

$$f(x) = \begin{cases} \frac{1}{40} & 60 \leq x \leq 100 \\ 0 & \text{elsewhere} \end{cases}$$

2. What is the mean time for this exam? What is the standard deviation?

$$E(X) = \frac{a+b}{2} = \frac{60+100}{2} = 80$$

$$\sigma(X) = \frac{b-a}{\sqrt{12}} = \frac{100-60}{\sqrt{12}} = 11.55$$

3. Find  $P(70 < X < 85)$

$$P(70 < X < 85) = \frac{85-70}{40} = 0.375$$

4. Find  $P(X > 90)$

$$P(X > 90) = \frac{100-90}{40} = 0.25$$

5. Find  $P(50 < X < 65)$

6.  $P(50 < X < 65) = \frac{65-50}{40} = 0.125$

## **Normal Distribution**


Data obtained from many experiments and studies often follow a common pattern called a **normal distribution**. Because it occurs so often in practical situations it is generally regarded as the most important and most widely used of all distributions.

### **Properties of a Normal Distribution**

1. The normal distribution is a **bell-shaped curve**.
2. The normal curve is symmetrical about the mean (The mean is at the middle and divides the area into two halves)
3. The total area under the curve is equal to 1.
4. The distribution is completely described by its mean  $\mu$  and standard deviation  $\sigma$ .
5. If  $X$  is a normal random variable with mean  $\mu$  and standard deviation  $\sigma$ , then  $X$  is denoted by  $X = N(\mu, \sigma)$
6. The probability density function for  $X = N(\mu, \sigma)$  is given by


$$f(x) = \frac{1}{\sigma\sqrt{2\pi}} e^{-\frac{(x-\mu)^2}{2\sigma^2}}$$

7.  $P(a \leq X \leq b)$  ( $P(a < X < b)$ ) is the area from  $x = a$  to  $x = b$  under the normal curve.



**P (a < x < b)**

8. The standard deviation of a normal curve gives the spread of the curve. As the standard deviation increases the spread of the curve increases. The following graph shows three normal curves with the same mean and different standard deviations.



### 9. **Recall : The Empirical Rule:**

- ❖ **68%** of data values are within **1 standard deviation** of the mean (-1 to +1).
- ❖ **95%** of data values are within **2 standard deviations** of the mean (-2 to +2).
- ❖ **99.7%** of data values are within **3 standard deviations** of the mean (-3 to +3).

### 10. **Standard Normal Curve**

A normal distribution curve is determined by two factors, the mean  $\mu$  and the standard deviation  $\sigma$ . That is, normal distributions with different means or different standard deviations give different normal curves. In such cases it is difficult to calculate the areas under the curves, or even to construct separate tables for areas under a normal curve for every value of  $\mu$  and of  $\sigma$ . Fortunately, we are able to transform all normal distributions into one distribution called **standard normal distribution**.

- ❖ The **standard normal** curve is a normal curve with mean  $\mu = 0$  and standard deviation  $\sigma = 1$ ,  $N(0, 1)$ .
- ❖ To transform from a normal curve into a standard normal curve we use the standardized value (z-score). The **z-score** is given by:


$$z = \frac{x - \mu}{\sigma}$$

- ❖ To find the area under a standard normal curve we use the standard normal distribution table, also called the **Z table**.

- ❖ The Z table lists the areas under the standard normal curve to the left of  $z$  that is,  $P(Z < z)$  for  $z \geq 0$ .




- ❖ We use the Z- table to find values on the right of the mean ( $\mu = 0$ ).
- ❖ Table entries for  $z$  represent the area under the bell curve to the left of  $z$ . (it is at least 0.5)
- ❖ To find areas to the right of  $z$  and to the left (right) negative value(s) we will use the **symmetric** of the curve and the fact that the **total area** under the curve is 1.



|     | 0.00   | 0.01   | 0.02   | 0.03   | 0.04   | 0.05   | 0.06   | 0.07   | 0.08   | 0.09   |
|-----|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|
| 0.0 | 0.5000 | 0.5040 | 0.5080 | 0.5120 | 0.5160 | 0.5199 | 0.5239 | 0.5279 | 0.5319 | 0.5359 |
| 0.1 | 0.5398 | 0.5438 | 0.5478 | 0.5517 | 0.5557 | 0.5596 | 0.5636 | 0.5675 | 0.5714 | 0.5753 |
| 0.2 | 0.5793 | 0.5832 | 0.5871 | 0.5910 | 0.5948 | 0.5987 | 0.6026 | 0.6064 | 0.6103 | 0.6141 |
| 0.3 | 0.6179 | 0.6217 | 0.6255 | 0.6293 | 0.6331 | 0.6368 | 0.6406 | 0.6443 | 0.6480 | 0.6517 |
| 0.4 | 0.6554 | 0.6591 | 0.6628 | 0.6664 | 0.6700 | 0.6736 | 0.6772 | 0.6808 | 0.6844 | 0.6879 |
| 0.5 | 0.6915 | 0.6950 | 0.6985 | 0.7019 | 0.7054 | 0.7088 | 0.7123 | 0.7157 | 0.7190 | 0.7224 |
| 0.6 | 0.7257 | 0.7291 | 0.7324 | 0.7357 | 0.7389 | 0.7422 | 0.7454 | 0.7486 | 0.7517 | 0.7549 |
| 0.7 | 0.7580 | 0.7611 | 0.7642 | 0.7673 | 0.7704 | 0.7734 | 0.7764 | 0.7794 | 0.7823 | 0.7852 |
| 0.8 | 0.7881 | 0.7910 | 0.7939 | 0.7967 | 0.7995 | 0.8023 | 0.8051 | 0.8078 | 0.8106 | 0.8133 |
| 0.9 | 0.8159 | 0.8186 | 0.8212 | 0.8238 | 0.8264 | 0.8289 | 0.8315 | 0.8340 | 0.8365 | 0.8389 |
| 1.0 | 0.8413 | 0.8438 | 0.8461 | 0.8485 | 0.8508 | 0.8531 | 0.8554 | 0.8577 | 0.8599 | 0.8621 |
| 1.1 | 0.8643 | 0.8665 | 0.8686 | 0.8708 | 0.8729 | 0.8749 | 0.8770 | 0.8790 | 0.8810 | 0.8830 |
| 1.2 | 0.8849 | 0.8869 | 0.8888 | 0.8907 | 0.8925 | 0.8944 | 0.8962 | 0.8980 | 0.8997 | 0.9015 |
| 1.3 | 0.9032 | 0.9049 | 0.9066 | 0.9082 | 0.9099 | 0.9115 | 0.9131 | 0.9147 | 0.9162 | 0.9177 |
| 1.4 | 0.9192 | 0.9207 | 0.9222 | 0.9236 | 0.9251 | 0.9265 | 0.9279 | 0.9292 | 0.9306 | 0.9319 |
| 1.5 | 0.9332 | 0.9345 | 0.9357 | 0.9370 | 0.9382 | 0.9394 | 0.9406 | 0.9418 | 0.9429 | 0.9441 |
| 1.6 | 0.9452 | 0.9463 | 0.9474 | 0.9484 | 0.9495 | 0.9505 | 0.9515 | 0.9525 | 0.9535 | 0.9545 |
| 1.7 | 0.9554 | 0.9564 | 0.9573 | 0.9582 | 0.9591 | 0.9599 | 0.9608 | 0.9616 | 0.9625 | 0.9633 |
| 1.8 | 0.9641 | 0.9649 | 0.9656 | 0.9664 | 0.9671 | 0.9678 | 0.9686 | 0.9693 | 0.9699 | 0.9706 |
| 1.9 | 0.9713 | 0.9719 | 0.9726 | 0.9732 | 0.9738 | 0.9744 | 0.9750 | 0.9756 | 0.9761 | 0.9767 |
| 2.0 | 0.9772 | 0.9778 | 0.9783 | 0.9788 | 0.9793 | 0.9798 | 0.9803 | 0.9808 | 0.9812 | 0.9817 |
| 2.1 | 0.9821 | 0.9826 | 0.9830 | 0.9834 | 0.9838 | 0.9842 | 0.9846 | 0.9850 | 0.9854 | 0.9857 |
| 2.2 | 0.9861 | 0.9864 | 0.9868 | 0.9871 | 0.9875 | 0.9878 | 0.9881 | 0.9884 | 0.9887 | 0.9890 |
| 2.3 | 0.9893 | 0.9896 | 0.9898 | 0.9901 | 0.9904 | 0.9906 | 0.9909 | 0.9911 | 0.9913 | 0.9916 |
| 2.4 | 0.9918 | 0.9920 | 0.9922 | 0.9925 | 0.9927 | 0.9929 | 0.9931 | 0.9932 | 0.9934 | 0.9936 |
| 2.5 | 0.9938 | 0.9940 | 0.9941 | 0.9943 | 0.9945 | 0.9946 | 0.9948 | 0.9949 | 0.9951 | 0.9952 |
| 2.6 | 0.9953 | 0.9955 | 0.9956 | 0.9957 | 0.9959 | 0.9960 | 0.9961 | 0.9962 | 0.9963 | 0.9964 |
| 2.7 | 0.9965 | 0.9966 | 0.9967 | 0.9968 | 0.9969 | 0.9970 | 0.9971 | 0.9972 | 0.9973 | 0.9974 |
| 2.8 | 0.9974 | 0.9975 | 0.9976 | 0.9977 | 0.9977 | 0.9978 | 0.9979 | 0.9979 | 0.9980 | 0.9981 |
| 2.9 | 0.9981 | 0.9982 | 0.9982 | 0.9983 | 0.9984 | 0.9984 | 0.9985 | 0.9985 | 0.9986 | 0.9986 |
| 3.0 | 0.9987 | 0.9987 | 0.9987 | 0.9988 | 0.9988 | 0.9989 | 0.9989 | 0.9989 | 0.9990 | 0.9990 |

# Table of Standard Normal Probabilities for Positive Z-Scores



| z   | .00   | .01   | .02   | .03   | .04   | .05   | .06   | .07   | .08   | .09   |
|-----|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|
| 0.0 | .5000 | .5040 | .5080 | .5120 | .5160 | .5199 | .5239 | .5279 | .5319 | .5359 |
| 0.1 | .5398 | .5438 | .5478 | .5517 | .5557 | .5596 | .5636 | .5675 | .5714 | .5753 |
| 0.2 | .5793 | .5832 | .5871 | .5910 | .5948 | .5987 | .6026 | .6064 | .6103 | .6141 |
| 0.3 | .6179 | .6217 | .6255 | .6293 | .6331 | .6368 | .6406 | .6443 | .6480 | .6517 |
| 0.4 | .6554 | .6591 | .6628 | .6664 | .6700 | .6736 | .6772 | .6808 | .6844 | .6879 |
| 0.5 | .6915 | .6950 | .6985 | .7019 | .7054 | .7088 | .7123 | .7157 | .7190 | .7224 |
| 0.6 | .7257 | .7291 | .7324 | .7357 | .7389 | .7422 | .7454 | .7486 | .7517 | .7549 |
| 0.7 | .7580 | .7611 | .7642 | .7673 | .7704 | .7734 | .7764 | .7794 | .7823 | .7852 |
| 0.8 | .7881 | .7910 | .7939 | .7967 | .7995 | .8023 | .8051 | .8078 | .8106 | .8133 |
| 0.9 | .8159 | .8186 | .8212 | .8238 | .8264 | .8289 | .8315 | .8340 | .8365 | .8389 |
| 1.0 | .8413 | .8438 | .8461 | .8485 | .8508 | .8531 | .8554 | .8577 | .8599 | .8621 |
| 1.1 | .8643 | .8665 | .8686 | .8708 | .8729 | .8749 | .8770 | .8790 | .8810 | .8830 |
| 1.2 | .8849 | .8869 | .8888 | .8907 | .8925 | .8944 | .8962 | .8980 | .8997 | .9015 |
| 1.3 | .9032 | .9049 | .9066 | .9082 | .9099 | .9115 | .9131 | .9147 | .9162 | .9177 |
| 1.4 | .9192 | .9207 | .9222 | .9236 | .9251 | .9265 | .9279 | .9292 | .9306 | .9319 |
| 1.5 | .9332 | .9345 | .9357 | .9370 | .9382 | .9394 | .9406 | .9418 | .9429 | .9441 |




### Example

Find the area under the standard normal curve for each of the following cases.

(a)  $P(Z < 1.25)$ ,  $P(Z < 1)$

From the table directly:  $P(Z < 1.25) = 0.8994$ ,  $P(Z < 1) = 0.8413$ .

(b)  $P(1 < Z < 1.25)$



$$\begin{aligned} P(1 < Z < 1.25) &= P(Z < 1.25) - P(Z < 1) \\ &= 0.8994 - 0.8413 \\ &= 0.0581 \end{aligned}$$


(c)  $P(Z > 1.25) = 1 - P(Z < 1.25) = 1 - 0.8994 = 0.1006$



(d)  $P(Z < -1.25) = P(Z > 1.25) = 0.1006$

(e)  $P(-1.25 < Z < 0) = P(0 < Z < 1.25)$


$$= 0.8994 - 0.5$$



(f)  $P(-1 < Z < 1.25) =$

$$P(-1 < Z < 1.25) = P(Z < 1.25) - P(Z < -1) = P(Z < 1.25) - [1 - P(Z < 1)]$$

$$= P(Z < 1.25) + P(Z < 1) - 1$$



### Example

Find  $z$  from  $z$  – table such that the area to the right of  $z$  is 0.05. That is, find  $z$  such that:  $P(Z > z) = 0.05$

- $P(Z > z) = 0.05 \Rightarrow P(Z < z) = 1 - 0.05 = 0.95$
- Find the value of  $z$  from the  $Z$  table that corresponds (approximately) to the area 0.95.
- We have two values near to 0.95; **0.9495** and **0.9505**. The values of  $z$  corresponds to those areas are **1.64** and **1.65**. So, the  $z$ -score corresponds to the area 0.95 is

$$z = (1.64 + 1.65) / 2 = 1.645$$

- In the cases for which the table does not contain an exact value that corresponds to the given area, find the value closest to the given area.

### Exercise

1. Find  $z$  from  $z$  – table such that the area to the right of  $z$  is 0.025
2. Find  $z$  from  $z$  – table such that the area to the right of  $z$  is 0.02

### Example

The IQ scores of 5000 students have a normal distribution with  $\mu = 100$  and  $\sigma = 15$ .

- (a) How many students have an IQ between 115 and 130?
- (b) How many students have an IQ between 100 and 130?
- (c) How many students have an IQ more than 124?
- (d) How many students have an IQ less than 85?
- (e) Find the IQ score for which 3 % of the scores exceed.

### Solution:

The IQ scores are normally distributed  $\Rightarrow X = N(100, 15)$

**(a)**  $P(115 < X < 130)$

$$x = 115: z_1 = \frac{x - \mu}{\sigma} = \frac{115 - 100}{15} = 1$$

$$x = 130: z_2 = \frac{x - \mu}{\sigma} = \frac{130 - 100}{15} = 2$$

$$P(115 < X < 130) = P(1 < Z < 2) = P(Z < 2) - P(Z < 1) \\ = 0.9772 - 0.8413 = 0.1341$$

The number of students with IQ scores between 115 and 130 is

$$(5000)(0.1341) = 671 \text{ students}$$

**(b)**  $x = 100: z_1 = 0$

$$x = 130: z_2 = 2$$

The percentage of students with IQ scores between 100 and 130 is,

$$P(0 < Z < 2) = 0.4772$$

The number of students with IQ scores between 100 and 130 is

$$(5000)(0.4772) = 2386 \text{ students.}$$

**(c)**  $P(X > 124) = P(Z > 1.6)$

$$[x = 124: z = \frac{124 - 100}{15} = 1.6]$$

$$P(Z > 1.6) = 1 - 0.9452$$

$$= 0.0448$$

The number of students with IQ scores more than 124 is

$$(5000)(0.0448) = 224 \text{ students}$$

**(d)**  $P(X > 85) = P(Z > -1) = P(Z < 1) = 0.8413$

$$[x = 85: z = \frac{85 - 100}{15} = -1]$$

The number of students with IQ scores less than 85 is,

$$(5000)(0.8413) = 4207 \text{ students.}$$

**(e)** Find  $x$ -value so that the area to the right of  $x$  under the normal curve is 0.03.

First, we find the  $z$  value that corresponds to the required  $x$  value.

$$P(Z > z) = 0.03 \Rightarrow P(Z < z) = 1 - 0.03 = 0.97$$

The value of  $z$  corresponds to the area 0.97 is 1.88

Now, use the formula for the  $z$ -score to find  $x$ . that is

$$\begin{aligned}x &= \mu + z \sigma \\ &= 100 + (1.88) (15) = 128.2\end{aligned}$$

The IQ score for which 3 % of the scores exceed is 128.2.

14/240

14. Given that  $z$  is a standard normal random variable, find  $z$  for each situation.
- The area to the left of  $z$  is .9750.
  - The area between 0 and  $z$  is .4750.
  - The area to the left of  $z$  is .7291.
  - The area to the right of  $z$  is .1314.
  - The area to the left of  $z$  is .6700.
  - The area to the right of  $z$  is .3300.

**b.** The area between 0 and  $z = 0.4750$  means the area to the left of  $z$  is  $0.5 + 0.4750 = 0.9750$ . So,  $z = 1.96$

**d.** If the area to the right of  $z = 0.1314$ , then the area to the left of  $z$  is  $1 - 0.1314 = 0.8686$ , from the table  $z = 1.12$

23/242

23. The time needed to complete a final examination in a particular college course is normally distributed with a mean of 80 minutes and a standard deviation of 10 minutes. Answer the following questions.
- What is the probability of completing the exam in one hour or less?
  - What is the probability that a student will complete the exam in more than 60 minutes but less than 75 minutes?
  - Assume that the class has 60 students and that the examination period is 90 minutes in length. How many students do you expect will be unable to complete the exam in the allotted time?

$X =$  Time needed.  $X = N(80, 10)$

- $P(X < 60) = P(Z < (60 - 80) / 10)$   
 $= P(Z < -2) = P(Z > 2)$   
 $= 1 - 0.9772 = 0.0228$
- $P(60 < X < 75) = P(-2 < Z < -0.5)$   
 $= P(0.5 < Z < 2)$   
 $= 0.9772 - 0.6915 = 0.2857$
- $P(x > 90) = P(Z > 1) = 1 - 0.8413 = 0.1587$

Expected number of students who will be unable to complete the exam is  $(0.1587)(60) = 10$  students