

1 Boas, problem p.564, 12.1-1

Solve the following differential equations by series and by another elementary method and check that the results agree:

$$xy' = xy + y \quad (1)$$

- by series: substituting the power series $y(x) = \sum_{n=0}^{\infty} a_n x^n$ in (1) we have

$$xy' - xy - y = 0 \iff x \left(\sum_{n=1}^{\infty} n a_n x^{n-1} \right) - x \left(\sum_{n=0}^{\infty} a_n x^n \right) - \sum_{n=0}^{\infty} a_n x^n = 0 \quad (2)$$

calling $n = m + 1$ and substituting,

$$\sum_{m=0}^{\infty} (m+1) a_{m+1} x^{m+1} - \sum_{n=0}^{\infty} a_n x^{n+1} - \sum_{n=0}^{\infty} a_n x^n = 0 \quad (3)$$

$$\sum_{m=0}^{\infty} [(m+1) a_{m+1} - a_m] x^{m+1} - a_0 - \sum_{m=0}^{\infty} a_{m+1} x^{m+1} = 0 \quad (4)$$

The only term with a 0-th power of x is a_0 , which tells us $a_0 = 0$. Asking for the coefficient of the $(m+1)$ -th power of x in (4) to be zero we get

$$a_{m+1} = \frac{1}{m} a_m = \frac{1}{m} \frac{1}{m-1} a_{m-1} = \dots = \frac{1}{m!} a_1 \quad (5)$$

So the solution for (1) is

$$y(x) = \sum_{m=0}^{\infty} a_{m+1} x^{m+1} = a_1 \sum_{m=0}^{\infty} \frac{1}{m!} x^{m+1} \quad (6)$$

Factoring out one power of x , one recognizes the power series of the exponential e^x and writes the solution as

$$y(x) = a_1 x e^x \quad (7)$$

- by separation of variables:

$$\frac{dy}{y} = \left(1 + \frac{1}{x}\right) dx \implies \ln y = x + \ln x + \ln c \implies y(x) = c x e^x \quad (8)$$

which is the same as in (7).

2 Boas, problem p.564, 12.1-10

Solve the following differential equations by series and by another elementary method and check that the results agree:

$$y'' - 4xy' + (4x^2 - 2)y = 0 \quad (9)$$

- by series: substituting the power series $y(x) = \sum_{n=0}^{\infty} a_n x^n$ in (9) we have

$$\begin{aligned} \sum_{n=2}^{\infty} n(n-1)a_n x^{n-2} - 4x \sum_{n=1}^{\infty} n a_n x^{n-1} + (4x^2 - 2) \sum_{n=0}^{\infty} a_n x^n &= 0 \\ 2a_2 + 6a_3x + \sum_{n=2}^{\infty} (n+2)(n+1)a_{n+2}x^n - 4a_1x - 4 \sum_{n=2}^{\infty} n a_n x^n + 4 \sum_{n=0}^{\infty} a_n x^{n+2} - 2a_0 - 2a_1x - 2 \sum_{n=2}^{\infty} a_n x^n &= 0 \\ 2a_2 + 6a_3x - 4a_1x - 2a_0 - 2a_1x + \sum_{n=2}^{\infty} \left[(n+2)(n+1)a_{n+2} - 2(1+2n)a_n + 4a_{n-2} \right] x^n &= 0 \end{aligned} \quad (10)$$

the first terms give

$$a_2 = a_0, \quad a_3 = a_1 \quad (11)$$

while the recursion formula is

$$(n+2)(n+1)a_{n+2} - 2(1+2n)a_n + 4a_{n-2} = 0, \quad \text{for } n \geq 2. \quad (12)$$

Consider first the case of $n = 2p$ even. Using (11), we can use the recursion formula to obtain a_4 . By repeated use of the recursion formula, we can obtain a_6, a_8, \dots . After computing a few values, it appears that the general form is

$$a_{2p} = \frac{a_0}{p!}. \quad (13)$$

Note that (13) is also valid for $p = 0$ and $p = 1$. To test the validity of (13), we insert this equation into (12):

$$\frac{2(p+1)(2p+1)}{(p+1)!} - \frac{2(1+4p)}{p!} + \frac{4}{(p-1)!} \stackrel{?}{=} 0. \quad (14)$$

Simple algebra verifies the validity of the equation above. Next, consider the case of $n = 2p+1$ odd. Using (11), we can use the recursion formula to obtain a_5 . By repeated use of the recursion formula, we can obtain a_7, a_9, \dots . After computing a few values, it appears that the general form is

$$a_{2p+1} = \frac{a_1}{p!}. \quad (15)$$

Note that (15) is also valid for $p = 0$ and $p = 1$. To test the validity of (15), we insert this equation into (12):

$$\frac{2(p+1)(2p+3)}{(p+1)!} - \frac{2(3+4p)}{p!} + \frac{4}{(p-1)!} \stackrel{?}{=} 0. \quad (16)$$

Simple algebra verifies the validity of the equation above. Hence, we conclude that

$$\begin{aligned} y(x) = \sum_{n=0}^{\infty} a_n x^n &= \sum_{p=0}^{\infty} a_{2p} x^{2p} + \sum_{p=0}^{\infty} a_{2p+1} x^{2p+1} \\ &= a_0 \sum_{p=0}^{\infty} \frac{x^{2p}}{p!} + a_1 \sum_{p=0}^{\infty} \frac{x^{2p+1}}{p!} \\ &= a_0 \sum_{p=0}^{\infty} \frac{[x^2]^p}{p!} + a_1 x \sum_{p=0}^{\infty} \frac{[x^2]^p}{p!} \\ &= (a_0 + a_1 x) e^{x^2}. \end{aligned} \quad (17)$$

- reduction of the order: one checks that the equation is solved by $y_0(x) = e^{x^2}$; then we look for another solution of the form $y(x) = u(x)y_0(x)$:

$$y' = u'y_0 + uy_0', \quad y'' = u''y_0 + 2u'y_0' + uy_0'' \quad (18)$$

$$y'' - 4xy' + (4x^2 - 2)y = u[y_0'' - 4xy_0' + (4x^2 - 2)y_0] + 2u'y_0' + u''y_0 - 4xu'y_0 = 0 \quad (19)$$

$$e^{x^2}[u'' - 4xu' + 4xu'] = 0 \implies u'' = 0 \implies u = A + Bx \quad (20)$$

$$y = (A + Bx)e^{x^2} \quad (21)$$

3 Boas, problem p.586, 12.11-5

Solve the following differential equations by the method of Frobenius:

$$2xy'' + y' + 2y = 0 \quad (22)$$

Substituting the generalized power series $y(x) = \sum_{n=0}^{\infty} a_n x^{n+s}$ in (22) we have

$$2x \sum_{n=0}^{\infty} (n+s)(n+s-1)a_n x^{n+s-2} + \sum_{n=0}^{\infty} (n+s)a_n x^{n+s-1} + 2 \sum_{n=0}^{\infty} a_n x^{n+s} = 0 \quad (23)$$

$$\text{the } n=0 \text{ term gives } [2s(s-1)a_0 + sa_0]x^{n+s-1} = 0 \implies 2s^2 - s = 0 \implies s = 0, \frac{1}{2} \quad (24)$$

$$\text{while the other terms are } \sum_{n=1}^{\infty} \left[2(n+s)(n+s-1)a_n + (n+s)a_n + 2a_{n-1} \right] x^{n+s-1} = 0 \quad (25)$$

$$(26)$$

For $s = 0$ we have

$$\sum_{n=1}^{\infty} \left[n(2n-1)a_n + 2a_{n-1} \right] x^{n-1} = 0 \quad (27)$$

$$a_n = \frac{-2}{n(2n-1)} a_{n-1} = \dots = \frac{(-2)^n}{n!(2n-1)!!} a_0 \quad (28)$$

Where the double factorial is $m!! = m(m-2)(m-4)\dots$. Also, we note that $(2n)! = 2n(2n-1)(2n-2)(2n-3)\dots = 2^n(2n-1)!!n!$. Inserting these coefficients back in the series gives

$$y(x) = \sum_{n=0}^{\infty} a_n x^n = \sum_{n=0}^{\infty} \frac{(-2x)^n}{n!(2n-1)!!} a_0 = \sum_{n=0}^{\infty} \frac{(-4x)^n}{(2n)!} a_0 = a_0 \sum_{n=0}^{\infty} \frac{(-1)^n (2\sqrt{x})^{2n}}{(2n)!} = a_0 \cos(2x^{1/2}). \quad (29)$$

For $s = \frac{1}{2}$ we have instead

$$\sum_{n=1}^{\infty} \left[n(2n+1)a_n + 2a_{n-1} \right] x^{n-\frac{1}{2}} = 0 \implies a_n = \frac{-2}{n(2n+1)} a_{n-1} = \dots = \frac{(-2)^n}{n!(2n+1)!!} a_0 \quad (30)$$

$$\implies y(x) = a_0 \sum \frac{(-1)^n (2\sqrt{x})^{2n+1}}{(2n+1)!} = a_0 \sin(2x^{1/2}) \quad (31)$$

The general solution is then given by the linear combination of (29), (31):

$$y = A \cos(2x^{1/2}) + B \sin(2x^{1/2}) \quad (32)$$

4 Boas, problem p.587, 12.11-14

Solve $y'' = -y$ by the Frobenius method.

We take the generalized power series $y(x) = \sum_{n=0}^{\infty} a_n x^{n+s}$ so that

$$y'' + y = \sum_{n=0}^{\infty} \left[(n+s)(n+s-1)a_n x^{n+s-2} + a_n x^{n+s} \right] = 0 \quad (33)$$

Taking the $n = 0$ term gives $s(s-1)a_0 = 0$, that is, $s = 0, 1$. For $s = 0$ we have

$$n(n-1)a_n + a_{n-2} = 0 \implies a_n = \frac{-a_{n-2}}{n(n-1)} \implies \begin{cases} a_{2n+1} = \frac{(-1)^n}{(2n+1)!} a_1 \\ a_{2n} = \frac{(-1)^n}{(2n)!} a_0 \end{cases} \quad (34)$$

These two series are those defining the sine and the cosine, so that we have found the well known result $y = a_0 \cos x + a_1 \sin x$.

If we now take $s = 1$ we have

$$(n+1)nb_n + b_{n-2} = 0 \implies b_n = -\frac{b_{n-2}}{n(n+1)} \quad (35)$$

The solution for $b_0 \neq 0$ is then

$$y(x) = \sum_{n=0}^{\infty} b_n x^{n+1} = b_1 \sin x + b_0 \sum_{n=1}^{\infty} \frac{(-1)^n}{(2n)!} x^{2n} \quad (36)$$

In this expression we are missing the x^0 term that would give the expansion of the cosine, and one easily check that this is not a solution of the differential equation. That happens because in (33) for $s = 1$ the first term coefficient reads $(n+1)(n+0)b_n$; then, b_0 coefficient has a 0 in front, which cancels it out from the rest of the problem, so we are calculating the solution *modulo* the constant b_0 .

5 Boas, problem p.567, 12.2-2

Show that $P_l(-1) = (-1)^l$.

Using eq. (2.6) on p. 565 of Boas, the general solution to the Legendre differential equation is:

$$y(x) = a_0 \left[1 - \frac{l(l+1)}{2!}x^2 + \frac{l(l+1)(l-2)(l+3)}{4!}x^4 - \dots \right] + a_1 \left[x - \frac{(l-1)(l+2)}{3!}x^3 + \frac{(l-1)(l+2)(l-3)(l+4)}{5!}x^5 - \dots \right]. \quad (37)$$

If l is even, then the Legendre polynomial is defined to be the polynomial proportional to a_0 (up to an overall normalization determined by convention). If l is odd, then the Legendre polynomial is defined to be the polynomial proportional to a_1 (up to an overall normalization determined by convention). It immediately follows that if l is even, then $P_l(x)$ is an even function of x , whereas if l is odd, then $P_l(x)$ is an odd function of x . This means that

$$P_l(-x) = (-1)^l P_l(x). \quad (38)$$

The normalization convention for the Legendre polynomials defines $P_l(1) = 1$. Hence, inserting $x = 1$ into (38) yields

$$P_l(-1) = (-1)^l \quad (39)$$

Note that eq. (38) is also an immediate consequence of the Rodrigues' formula,

$$P_l(x) = \frac{1}{2^l l!} \frac{d^l}{dx^l} (x^2 - 1)^l, \quad (40)$$

and provides another way of deriving (39).

6 Boas, problem p.567, 12.2-4

We will solve Legendre equation

$$(1 - x^2)y'' - 2xy' + l(l + 1)y = 0 \quad (41)$$

using the method of reduction of order: given the known solution $P_l(x)$, we look for an independent solution of the form $y(x) = P_l(x)v(x)$ and then solve for $v(x)$ in (41):

$$(1 - x^2)(v''P_l + 2v'P_l' + vP_l'') - 2x(v'P_l + vP_l') + l(l + 1)P_l = 0 \quad (42)$$

$$(1 - x^2)(v''P_l(x) + 2v'P_l'(x)) - 2xP_l(x)v' = 0 \implies (1 - x^2)P_l(x)v'' + 2((1 - x^2)P_l'(x) - xP_l(x))v' = 0$$

$$\implies \frac{v''}{v'} = 2 \frac{xP_l(x) - (1 - x^2)P_l'}{(1 - x^2)P_l(x)} = 2 \frac{x}{1 - x^2} - 2 \frac{P_l'}{P_l} = \frac{1}{1 - x} - \frac{1}{1 + x} - 2 \frac{P_l'}{P_l} \quad (43)$$

which is solved by

$$\ln v' = -\ln(1 - x) - \ln(1 + x) - 2 \ln P_l = \ln \frac{1}{(1 - x)(1 + x)P_l^2}, \text{ that is,} \quad (44)$$

$$v(x) = \int \frac{dx}{(1 - x)(1 + x)P_l^2} \quad (45)$$

The second solution of the Legendre equation is then

$$Q_l(x) = P_l(x)v(x) \quad (46)$$

We evaluate this expression for the two cases $l = 0, 1$:

- $l = 0$: $P_0(x) = 1$, so the other solution is

$$Q_0(x) = \int dx \frac{1}{(1 - x)(1 + x)} = \frac{1}{2} \int dx \left(\frac{1}{1 - x} + \frac{1}{1 + x} \right) = \frac{1}{2} \ln \frac{1 + x}{1 - x} \quad (47)$$

- $l = 1$: $P_1(x) = x$, so the other solution is

$$Q_1(x) = x \int dx \frac{1}{(1 - x)(1 + x)x^2} = x \int dx \left(\frac{1}{2} \frac{1}{1 - x} + \frac{1}{2} \frac{1}{1 + x} + \frac{1}{x^2} \right) = \quad (48)$$

$$= \frac{x}{2} \ln \frac{1 + x}{1 - x} - 1 \quad (49)$$

7 Boas, problem p.568, 12.3-1

We will use the hint in problem 12.3-6: if we write

$$\frac{d}{dx}uv = D(uv) = (D_u + D_v)uv \quad (50)$$

where D_u, D_v are operators that act only separately on u, v , we have

$$\frac{d^n}{dx^n}uv = (D_u + D_v)^n uv = \sum_{k=0}^n \binom{n}{k} D_u^k D_v^{n-k} uv = \sum_{k=0}^n \binom{n}{k} D_u^k u D_v^{n-k} v = \sum_{k=0}^n \binom{n}{k} \frac{d^k}{dx^k} u \frac{d^{n-k}}{dx^{n-k}} v \quad (51)$$

where we have used the expansion of the n -th power of a binomial formed by the two operators D_u, D_v (which commute with each other). $\binom{n}{k} = \frac{n(n-1)\dots(n-k+1)}{k!}$ is the binomial coefficient.

8 Boas, problem p.569, 12.4-2

By Rodrigues' formula

$$P_l(x) = \frac{1}{2^l l!} \frac{d^l}{dx^l} (x^2 - 1)^l \quad (52)$$

we have, after applying Leibniz' rule (51)

$$P_l(x) = \frac{1}{2^l l!} \sum_{k=0}^l \binom{l}{k} \frac{d^k}{dx^k} (x+1)^l \frac{d^{l-k}}{dx^{l-k}} (x-1)^l \quad (53)$$

Now, every time we differentiate $(x-1)^l$ we lower the exponent by one; in particular, when we differentiate l times, we are left with a constant; when we calculate $P_l(1)$ any factor of $(x-1)$ will become zero, so that the only non zero contribution comes from the 0-th term in the sum: this gives

$$P_l(1) = \frac{2^l}{l!} \cdot 2^l \cdot l! = 1 \quad (54)$$

where the term 2^l comes from $(x+1)^l$ for $x=1$ and $\frac{d^l}{dx^l} (x-1)^l = l \frac{d^{l-1}}{dx^{l-1}} (x-1)^{l-1} = l(l-1) \frac{d^{l-2}}{dx^{l-2}} (x-1)^{l-2} = \dots = l!$.

9 Boas, problem p.569, 12.4-4

We want to prove that

$$\int_{-1}^1 x^m P_l(x) dx = 0, \quad \text{for } m < l \quad (55)$$

Substituting Rodrigues' formula (52) we have

$$\int_{-1}^1 x^m P_l(x) dx = \int_{-1}^1 \frac{x^m}{2^l l!} \frac{d^l}{dx^l} (x^2 - 1)^l dx \propto \quad (56)$$

$$\propto x^m \frac{d^{l-1}}{dx^{l-1}} (x^2 - 1)^l \Big|_{-1}^1 - \int_{-1}^1 m x^{m-1} \frac{d^{l-1}}{dx^{l-1}} (x^2 - 1)^l dx = \quad (57)$$

$$= 0 - m x^{m-1} \frac{d^{l-2}}{dx^{l-2}} (x^2 - 1)^l \Big|_{-1}^1 + \int_{-1}^1 m(m-1) x^{m-2} \frac{d^{l-2}}{dx^{l-2}} (x^2 - 1)^l dx = \dots \quad (58)$$

in the first passage we have neglected the constant $\frac{1}{2^l l!}$ and integrated by parts; in the second passage, we see that the first term is null because after we have differentiated $(l - 1)$ times we will still have at least one factor of $(x - 1)$ and one of $(x + 1)$ (as you can quickly check by using Leibniz' rule), which are zero when evaluated at ± 1 . The same happens for all the other terms evaluated at ± 1 , so that, after we have integrated by parts m times (assuming $m < l$), we are left with

$$m! \int_{-1}^1 \frac{d^{l-m}}{dx^{l-m}} (x^2 - 1)^l dx = m! \frac{d^{l-m-1}}{dx^{l-m-1}} (x^2 - 1)^l \Big|_{-1}^1 = 0 \quad (59)$$

for the same argument we used above. Note that this does not hold for $m > l$, because in that case between (58) and (59) we reach a step in which $l - k = 0$ and we have $\int x^{m-k} (x^2 - 1)^l \neq 0$

10 Boas, problem p.574, 12.5-10

Express the following polynomial as a linear combination of the Legendre polynomials:

$$f(x) = x^4 \quad (60)$$

The first five Legendre polynomials are:

$$P_0 = 1, \quad P_1 = x, \quad P_2 = \frac{1}{2}(3x^2 - 1), \quad P_3 = \frac{1}{2}(5x^3 - 3x), \quad P_4 = \frac{1}{8}(35x^4 - 30x^2 + 3) \quad (61)$$

We are going to expand x^4 as a linear combination of the Legendre polynomials, with unknown coefficients; these will be found imposing that the factors for the different powers of x coincide. Because we have x^4 , $f(x) = \sum_0^4 c_n P_n$ must contain P_4 ; in particular, $c_4 = \frac{8}{35}$, so that the coefficient of x^4 is 1. Then we must put to zero the coefficient of x^3 : x^3 only appears in P_3 so we can put $c_3 = 0$. Right now, our function is written as

$$f(x) = \sum_0^2 c_n P_n + \frac{8}{35} P_4(x) \quad (62)$$

Now we fix to zero the coefficient of x^2 : it appears in P_4 and P_2 and it is

$$\frac{3}{2}c_2 + \frac{-30}{35} = 0 \implies c_2 = \frac{4}{7}$$

A term linear in x appears only in P_1 , so we can set $c_1 = 0$. Finally, the constant term is given by

$$c_0 - \frac{1}{2}c_2 + \frac{3}{8}c_4 = 0 \implies c_0 = \frac{1}{5} \quad (63)$$

Then we have found

$$x^4 = \frac{1}{5}P_0(x) + \frac{4}{7}P_2(x) + \frac{8}{35}P_4(x) \quad (64)$$

11 Boas, problem p.577, 12.6-6

We want to show that P_l and P'_l are orthogonal on $[-1,1]$ in two ways:

- we can use the fact that the Legendre polynomials are either even or odd functions of x (depending on whether l is even or odd, respectively), as shown in problem 5. Then, if P_l is odd, its derivative P'_l is even, and vice versa. In general, if $f(x)$ is an even function of x and $g(x)$ is an odd function of x , then

$$\int_{-a}^a f(x)g(x) = 0. \quad (65)$$

This is easily proven by changing the integration variable to $y = -x$, in which case

$$\int_{-a}^a f(x)g(x) = - \int_{+a}^{-a} f(-y)g(-y)dy = \int_{-a}^a f(-y)g(-y)dy = - \int_{-a}^a f(y)g(y)dy, \quad (66)$$

As the integral is equal to minus itself, it must be equal to zero. Hence, we conclude that

$$\int_{-1}^1 P_l(x)P'_l(x) = 0. \quad (67)$$

- we can also use the result of problem 9: remember that P_l is a polynomial of order l and P'_l is a polynomial of order $(l-1)$. Then

$$\int_{-1}^1 P_l(x)P'_l(x) \quad (68)$$

is given by a sum of terms which have the form $c_n \int_{-1}^1 x^m P_l(x)dx$, where $m = 0, 1, \dots, l-1$, that is, $m < l$, so they are all zero and the two functions are orthogonal.

12 Boas, problem p.615, 12.23-2

The generating functional of the Legendre polynomials is

$$\Phi(x, h) = \frac{1}{\sqrt{1-2xh+h^2}} = \sum_{l=0}^{\infty} h^l P_l(x); \quad (69)$$

for $x = 0$ this gives

$$\Phi(0, h) = \sum_{l=0}^{\infty} h^l P_l(0) = \frac{1}{\sqrt{1+h^2}} = \sum c_l h^l, \quad (70)$$

But the function $\Phi(0, h)$ looks exactly like the power of a binomial:

$$\Phi(0, h) = (1+h^2)^{-1/2} = \sum_{n=0}^{\infty} \binom{-1/2}{n} h^{2n} \quad (71)$$

Here we can read the Legendre polynomials in zero as

$$P_{2n+1}(0) = 0; \quad (72)$$

$$P_{2n}(0) = \binom{-1/2}{n} = \frac{-\frac{1}{2}(-\frac{1}{2}-1)\dots(-\frac{1}{2}-n+1)}{(n)!} = \frac{(-1)^n(2n-1)!!}{2^n n!}. \quad (73)$$