

45

HW #3

Artem Mavrin

MATH 342B ASSIGNMENT 5

PROBLEMS, SECTION 12.1

Solve the following differential equations by series and also by an elementary method and verify that your solutions agree. Check your results by computer.

1.
 $xy' = xy + y.$

Separation of variables:

From the given equation we get

$$(1) \quad xy' = xy + y \implies xy' = y(x + 1) \implies \frac{1}{y} \frac{dy}{dx} = \frac{x + 1}{x} \implies \int \frac{1}{y} dy = \int \frac{x + 1}{x} dx.$$

So, the left hand side evaluates to

$$(2) \quad \ln y + C_1$$

for some constant $C_1 \in \mathbb{R}$ and the right hand side is simplified as follows:

$$(3) \quad \int \frac{x + 1}{x} dx = \int (1 + x^{-1}) dx = x + \ln x + C_2$$

for some $C_2 \in \mathbb{R}$. Let $C = C_2 - C_1$. Then we have from (1), (2), and (3) that

$$(4) \quad \ln y = x + \ln x + C,$$

so

$$(5) \quad \begin{aligned} y &= \exp(x + \ln x + C) \\ &= \exp(x) \exp(\ln x) \exp(C) \\ &= y_0 x e^x, \end{aligned}$$

where $y_0 = \exp(C)$. So, $y = y_0 x e^x$. Wolfram Alpha confirms this.

Series method:

Assume a solution

$$(6) \quad y = \sum_{n=0}^{\infty} a_n x^n$$

for $a_n \in \mathbb{R}$. Then

$$(7) \quad y' = \sum_{n=0}^{\infty} a_n n x^{n-1}, \quad y'' = \sum_{n=0}^{\infty} a_n n(n-1) x^{n-2}.$$

Plugging (6) and (7) into the differential equation, we get

$$(8) \quad xy' = xy + y \implies x \sum_{n=0}^{\infty} a_n n x^{n-1} = (x + 1) \sum_{n=0}^{\infty} a_n x^n.$$

So,

$$(9) \quad \sum_{n=0}^{\infty} a_n n x^n = \sum_{n=0}^{\infty} a_n x^n + \sum_{n=0}^{\infty} a_n x^{n+1}$$

$$\implies \sum_{n=0}^{\infty} a_{n+1} (n+1) x^{n+1} = a_0 + \sum_{n=0}^{\infty} (a_{n+1} + a_n) x^{n+1}.$$

So, $a_0 = 0$ and

$$(10) \quad a_{n+1} (n+1) = a_{n+1} + a_n \implies a_{n+1} = \frac{1}{n} a_n.$$

So, $a_2 = a_1$, $a_3 = \frac{1}{2} a_2 = \frac{1}{2} a_1$, $a_4 = \frac{1}{3} a_3 = \frac{1}{6} a_1$. In general, for $n \geq 1$, we have

$$(11) \quad a_n = \frac{1}{(n-1)!} a_1.$$

Therefore, our solution becomes

$$(12) \quad y = a_1 \sum_{n=1}^{\infty} \frac{1}{(n-1)!} x^n.$$

This is equivalent to $y = y_0 x e^x$ if $y_0 = a_1$:

$$(13) \quad a_1 \sum_{n=1}^{\infty} \frac{1}{(n-1)!} x^n = a_1 x \sum_{n=1}^{\infty} \frac{1}{(n-1)!} x^{n-1} = a_1 x \sum_{n=0}^{\infty} \frac{1}{n!} x^n = a_1 x e^x.$$

Thus, the solution to the differential equation is

$$(14) \quad y = a_1 \sum_{n=1}^{\infty} \frac{1}{(n-1)!} x^n = a_1 x e^x$$

for $a_1 \in \mathbb{R}$.

2.

$$y' = 3x^2 y.$$

Separation of variables:

From the given equation we get

$$(15) \quad y' = 3x^2 y \implies \frac{1}{y} \frac{dy}{dx} = 3x^2 \implies \int \frac{1}{y} dy = 3 \int x^2 dx \implies \ln y = x^3 + C$$

where $C \in \mathbb{R}$ is a constant. Let $y_0 = \exp(C)$. Then we get

$$(16) \quad y = \exp(x^3 + C) = \exp(x^3) \exp(C) = y_0 e^{x^3}.$$

So, $y = y_0 e^{x^3}$. Wolfram Alpha agrees.

Series method:

Assume a solution as in (6), so that (7) follows. Then we plug (6) and (7) into the differential equation to get

$$(17) \quad \sum_{n=0}^{\infty} a_n n x^{n-1} = 3x^2 \sum_{n=0}^{\infty} a_n x^n = 3 \sum_{n=0}^{\infty} a_n x^{n+2}.$$

So, we have

$$(18) \quad a_1 + 2a_2 + \sum_{n=0}^{\infty} a_{n+3}(n+3)x^{n+2} = 3 \sum_{n=0}^{\infty} a_n x^{n+2}.$$

So, $a_1 = a_2 = 0$, a_0 is arbitrary, and

$$(19) \quad a_{n+3}(n+3) = 3a_n \implies a_{n+3} = \frac{3}{n+3}a_n.$$

So, $a_3 = \frac{3}{3}a_0 = a_0$, $a_6 = \frac{3}{6}a_3 = \frac{1}{2}a_3 = \frac{1}{2}a_0$, $a_9 = \frac{3}{9}a_6 = \frac{1}{3}a_6 = \frac{1}{6}a_0$. In general,

$$(20) \quad a_{3n} = \frac{1}{n!}a_0.$$

Thus, our solution is

$$(21) \quad y = a_0 \left(1 + x^3 + \frac{1}{2}x^6 + \frac{1}{9}x^9 + \dots \right) = a_0 \sum_{n=0}^{\infty} \frac{1}{n!}x^{3n}.$$

This is exactly the Taylor expansion of $a_0 e^{x^3}$. So, if $y_0 = a_0$, we have the general solution of the differential equation:

$$(22) \quad y = a_0 \sum_{n=0}^{\infty} \frac{1}{n!}x^{3n} = a_0 e^{x^3}$$

for $a_0 \in \mathbb{R}$.

4.

$$y'' = -4y.$$

Linear homogeneous second order DE:

Since the differential equation is linear, assume a solution $y = e^{nx}$. Then $y' = ne^{nx}$ and $y'' = n^2 e^{nx}$. So, we get

$$(23) \quad y'' = -4y \implies n^2 e^{nx} = -4e^{nx} \implies n^2 = -4 \implies n = \pm 2i.$$

Thus,

$$(24) \quad y = Ae^{i2x} + Be^{-i2x}$$

for some $A, B \in \mathbb{R}$. So, using Euler's formula, (24) becomes

$$(25) \quad \begin{aligned} y &= A[\cos(2x) + i \sin(2x)] + B[\cos(-2x) + i \sin(-2x)] \\ &= A[\cos(2x) + i \sin(2x)] + B[\cos(2x) - i \sin(2x)] \\ &= (A + B) \cos(2x) + i(A - B) \sin(2x). \end{aligned}$$

Let $C_0 = A + B$ and let $C_1 = (A - B)i$. Then $y = C_0 \cos(2x) + C_1 \sin(2x)$. Wolfram Alpha confirms this.

Series method:

Assume a solution as in (6), so that (7) follows. Then we plug (6) and (7) into the differential equation to get

$$(26) \quad \sum_{n=0}^{\infty} a_n n(n-1)x^{n-2} = -4 \sum_{n=0}^{\infty} a_n x^n.$$

So, we have

$$(27) \quad \sum_{n=0}^{\infty} a_{n+2}(n+2)(n+1)x^n = -4 \sum_{n=0}^{\infty} a_n x^n.$$

Thus, a_0 and a_1 are arbitrary and

$$(28) \quad a_{n+2}(n+2)(n+1) = -4a_n \implies a_{n+2} = -\frac{4}{(n+2)(n+1)}a_n.$$

So, $a_2 = -\frac{4}{2 \cdot 1}a_0$, $a_3 = -\frac{4}{3 \cdot 2}a_1$, $a_4 = -\frac{4}{4 \cdot 3}a_2 = \frac{16}{4 \cdot 3 \cdot 2 \cdot 1}a_0$, $a_5 = -\frac{4}{5 \cdot 4}a_3 = \frac{16}{5 \cdot 4 \cdot 3 \cdot 2}a_1$. Thus, in general we have

$$(29) \quad a_{2n} = \frac{(-1)^n 4^n}{(2n)!} a_0, \quad n \geq 0,$$

and

$$(30) \quad a_{2n+1} = \frac{(-1)^n 4^n}{(2n+1)!} a_1, \quad n \geq 0.$$

So, the solution becomes

$$(31) \quad \begin{aligned} y &= a_0 \sum_{n=0}^{\infty} \frac{(-1)^n 4^n}{(2n)!} x^{2n} + a_1 \sum_{n=0}^{\infty} \frac{(-1)^n 4^n}{(2n+1)!} x^{2n+1} \\ &= a_0 \sum_{n=0}^{\infty} \frac{(-1)^n}{(2n)!} (2x)^{2n} + \frac{a_1}{2} \sum_{n=0}^{\infty} \frac{(-1)^n}{(2n+1)!} (2x)^{2n+1} \\ &= a_0 \cos(2x) + \frac{a_1}{2} \sin(2x). \end{aligned}$$

If we let $a_0 = C_0$ and $\frac{a_1}{2} = C_1$ then (31) is identical to the original answer of $y = C_0 \cos(2x) + C_1 \sin(2x)$. Thus, the general solution of the differential equation is

$$(32) \quad y = C_0 \sum_{n=0}^{\infty} \frac{(-1)^n}{(2n)!} (2x)^{2n} + C_1 \sum_{n=0}^{\infty} \frac{(-1)^n}{(2n+1)!} (2x)^{2n+1} = C_0 \cos(2x) + C_1 \sin(2x).$$

9.

$$(x^2 + 1)y'' - 2xy' + 2y = 0.$$

Reduction of order:

Given the differential equation $(x^2 + 1)y'' - 2xy' + 2y = 0$, assume a solution $y_1 = x^m$ for some $m \in \mathbb{R}$. Then $y_1' = mx^{m-1}$ and $y_1'' = m(m-1)x^{m-2}$, so we get

$$(33) \quad \begin{aligned} (x^2 + 1)y_1'' - 2xy_1' + 2y_1 &= 0 \implies m(m-1)(x^m + x^{m-2}) - 2mx^m + 2x^m = 0 \\ \implies (m^2 - 3m + 2)x^m + (m^2 - m)x^{m-2} &= 0 \implies m^2 - 3m + 2 = m^2 - m = 0 \\ \implies m &= 1. \end{aligned}$$

So $y_1 = x$ is a solution of the differential equation. Suppose $y_2 = y_1(x)v(x) = xv(x)$ is another solution to the differential equation for some function $v : \mathbb{R} \rightarrow \mathbb{R}$. Then

$y_2' = v + xv'$ and $y_2'' = 2v' + xv''$. Plugging these into the differential equation, we get

$$\begin{aligned}
 (34) \quad & (x^2 + 1)y_2'' - 2xy_2' + 2y_2 = 0 \\
 \implies & (x^2 + 1)(2v' + xv'') - 2x(v + xv') + 2xv = 0 \\
 \implies & 2x^2v' + 2v' + x^3v'' + xv'' - 2xv - 2x^2v' + 2xv = 0 \\
 \implies & 2v' + x^3v'' + xv'' = 0 \\
 \implies & (x^3 + x)v'' + 2v' = 0 \\
 \implies & \frac{dv'}{dx} = -\frac{2}{x^3 + x}v' \\
 \implies & \int \frac{1}{v'} dv' = -2 \int \frac{1}{x^3 + x} dx = -2 \int \left(\frac{1}{x} - \frac{x}{x^2 + 1} \right) dx \\
 \implies & \ln v' = -2 \ln(x) + \ln(x^2 + 1) + C \quad [C \in \mathbb{R}] \\
 \implies & v' = e^C \left(\frac{x^2 + 1}{x^2} \right) = e^C \left(1 + \frac{1}{x^2} \right) \\
 \implies & v = e^C \left(x - \frac{1}{x} \right).
 \end{aligned}$$

So,

$$(35) \quad y_2 = xv = e^C x \left(x - \frac{1}{x} \right) = e^C (x^2 - 1).$$

Let $B = e^C$ and let $A \in \mathbb{R}$ be an arbitrary constant. Since $y_2 = B(x^2 - 1)$ is a solution and $y_1 = x$ is a solution (hence $y_1 = Bx$ is a solution), $y = y_1 + y_2$ is the general solution. So, $y = A(x^2 - 1) + Bx$ is the solution to the differential equation. Wolfram Alpha confirms this.

Series method:

Assume a solution as in (6), so that (7) follows. Then we plug (6) and (7) into the differential equation to get

$$(36) \quad (x^2 + 1) \sum_{n=0}^{\infty} a_n n(n-1)x^{n-2} - 2x \sum_{n=0}^{\infty} a_n n x^{n-1} + 2 \sum_{n=0}^{\infty} a_n x^n = 0.$$

Thus,

$$(37) \quad \sum_{n=0}^{\infty} a_n n(n-1)x^{n-2} + \sum_{n=0}^{\infty} (n(n-1) - 2n + 2)a_n x^n = 0.$$

So,

$$(38) \quad \sum_{n=0}^{\infty} (n(n-1) - 2n + 2)a_n x^n = - \sum_{n=0}^{\infty} a_{n+2}(n+2)(n+1)x^n,$$

and a_0 and a_1 are arbitrary constants. Therefore,

$$(39) \quad (n(n-1) - 2n + 2)a_n = -a_{n+2}(n+2)(n+1),$$

so

$$(40) \quad a_{n+2} = -\frac{n(n-1) - 2n + 2}{(n+2)(n+1)} a_n = -\frac{n^2 - 3n + 2}{n^2 + 3n + 2} a_n.$$

Then $a_2 = -a_0$, $a_3 = -\frac{1-3+2}{1+2+3}a_1 = 0$, $a_4 = -\frac{4-6+2}{4+6+2}a_2 = 0$. Thus, for $n \geq 2$ we have $a_n = 0$. So, the solution is

$$(41) \quad y = \sum_{n=0}^2 a_n x^n = a_0 + a_1 x + -a_2 x^2 = a_0(1 - x^2) + a_1 x.$$

If we set $A = -a_0$ and $B = a_1$ then we get the same solution we found earlier, namely

$$(42) \quad y = A(x^2 - 1) + Bx.$$

PROBLEMS, SECTION 12.2

2.

Show that $P_l(-1) = (-1)^l$.

The Legendre differential equation is

$$(43) \quad (1 - x^2) \frac{d^2 y}{dx^2} - 2x \frac{dy}{dx} + l(l+1)y = 0.$$

A general solution of (43) is

$$(44) \quad y(x) = a_0 \left[1 - \frac{l(l+1)}{2!} x^2 + \frac{l(l+1)(l-2)(l+3)}{4!} x^4 - \dots \right] \\ + a_1 \left[x - \frac{(l-1)(l+2)}{3!} x^3 + \frac{(l-1)(l+2)(l-3)(l+4)}{4!} x^5 - \dots \right].$$

Put $-x$ in place of x in the above equation. Then we get

$$(45) \quad y(-x) = a_0 \left[1 - \frac{l(l+1)}{2!} x^2 + \frac{l(l+1)(l-2)(l+3)}{4!} x^4 - \dots \right] \\ - a_1 \left[x - \frac{(l-1)(l+2)}{3!} x^3 + \frac{(l-1)(l+2)(l-3)(l+4)}{4!} x^5 - \dots \right].$$

The Legendre polynomials $P_l(x)$ are just special cases of (44) where l is a positive integer and we have the restriction $P_l(1) = 1$. For odd l we choose $a_0 = 0$ and for even l we choose $a_1 = 0$. Thus, for odd l we have

$$(46) \quad P_l(x) = a_1 \left[x - \frac{(l-1)(l+2)}{3!} x^3 + \frac{(l-1)(l+2)(l-3)(l+4)}{4!} x^5 - \dots \right],$$

so

$$(47) \quad P_l(-x) = a_1 \left[-x - \frac{(l-1)(l+2)}{3!} (-x)^3 + \frac{(l-1)(l+2)(l-3)(l+4)}{4!} (-x)^5 - \dots \right] \\ = -a_1 \left[x - \frac{(l-1)(l+2)}{3!} x^3 + \frac{(l-1)(l+2)(l-3)(l+4)}{4!} x^5 - \dots \right] \\ = -P_l(x) = (-1)^l P_l(x).$$

Similarly, for even l we have

$$(48) \quad P_l(x) = a_0 \left[1 - \frac{l(l+1)}{2!} x^2 + \frac{l(l+1)(l-2)(l+3)}{4!} x^4 - \dots \right],$$


so

$$\begin{aligned} (49) \quad P_l(-x) &= a_0 \left[1 - \frac{l(l+1)}{2!}(-x)^2 + \frac{l(l+1)(l-2)(l+3)}{4!}(-x)^4 - \dots \right] \\ &= a_0 \left[1 - \frac{l(l+1)}{2!}x^2 + \frac{l(l+1)(l-2)(l+3)}{4!}x^4 - \dots \right] \\ &= P_l(x) = (-1)^l P_l(x). \end{aligned}$$


Thus, for any $l \in \mathbb{N}$, we have $P_l(-x) = (-1)^l P_l(x)$. Since we construct P_l such that $P_l(1) = 1$, it follows that $P_l(-1) = (-1)^l$, which was to be shown. \square

[Graphs of Legendre Polynomials


```
> plot(LegendreP(0,x),x=-1..1);
```



```
> plot(LegendreP(1,x),x=-1..1);
```



```
> plot(LegendreP(2,x),x=-1..1);
```

[>

MATH 342B ASSIGNMENT 6

PROBLEMS, SECTION 12.5

4.

Show from (5.1) that

$$(x - h) \frac{\partial \Phi}{\partial x} = h \frac{\partial \Phi}{\partial h}.$$

Substitute the series (5.2) for Φ , and so prove the recursion relation (5.8b).

In the textbook, equation (5.1) reads,

$$(1) \quad \Phi(x, h) = (1 - 2xh + h^2)^{-1/2}, \quad |h| < 1,$$

and Φ is the generating function for Legendre polynomials. We take partial derivatives of Φ :

$$(2) \quad \begin{aligned} \frac{\partial \Phi}{\partial x} &= -\frac{1}{2}(1 - 2xh + h^2)^{-3/2}(-2h) \\ &= h(1 - 2xh + h^2)^{-3/2}, \quad \text{and} \end{aligned}$$

$$(3) \quad \begin{aligned} \frac{\partial \Phi}{\partial h} &= -\frac{1}{2}(1 - 2xh + h^2)^{-3/2}(-2x + 2h) \\ &= (x - h)(1 - 2xh + h^2)^{-3/2}. \end{aligned}$$

Multiplying (2) by $x - h$ and (3) by h , we get

$$(4) \quad (x - h) \frac{\partial \Phi}{\partial x} = h(x - h)(1 - 2xh + h^2)^{-3/2}, \quad \text{and}$$

$$(5) \quad h \frac{\partial \Phi}{\partial h} = h(x - h)(1 - 2xh + h^2)^{-3/2},$$

so that

$$(6) \quad \boxed{(x - h) \frac{\partial \Phi}{\partial x} = h \frac{\partial \Phi}{\partial h}},$$

as desired. Furthermore, equation (5.2) in the textbook reads

$$(7) \quad \Phi(x, h) = \sum_{l=0}^{\infty} h^l P_l(x).$$

Substituting (7) into (6), we get

$$(8) \quad (x - h) \frac{\partial}{\partial x} \sum_{l=0}^{\infty} h^l P_l(x) = h \frac{\partial}{\partial h} \sum_{l=0}^{\infty} h^l P_l(x).$$

The left hand side of (8) becomes

$$\begin{aligned}
(9) \quad (x-h) \frac{\partial}{\partial x} \sum_{l=0}^{\infty} h^l P_l(x) &= (x-h) \left[\frac{\partial}{\partial x} P_0(x) + \frac{\partial}{\partial x} \sum_{l=1}^{\infty} h^l P_l(x) \right] \\
&= (x-h) \left[\frac{\partial}{\partial x} (1) + \frac{\partial}{\partial x} \sum_{l=1}^{\infty} h^l P_l(x) \right] \\
&= (x-h) \sum_{l=1}^{\infty} h^l P'_l(x) \\
&= \sum_{l=1}^{\infty} h^l x P'_l(x) - \sum_{l=1}^{\infty} h^{l+1} P'_l(x) \\
&= \sum_{l=1}^{\infty} h^l x P'_l(x) - \sum_{l=1}^{\infty} h^l P'_{l-1}(x) \\
&= \sum_{l=1}^{\infty} h^l [x P'_l(x) - P'_{l-1}(x)].
\end{aligned}$$

The second to last step is possible since at $l = 1$, $P'_{l-1}(x) = \frac{d}{dx} 1 = 0$, so

$$\begin{aligned}
(10) \quad \sum_{l=1}^{\infty} h^{l+1} P'_l(x) &= \sum_{l=1}^{\infty} h^{l+1} P'_l(x) + 0 \\
&= \sum_{l=1}^{\infty} h^{l+1} P'_l(x) + h^1 P'_0(x) \\
&= \sum_{l=0}^{\infty} h^{l+1} P'_l(x) \\
&= \sum_{l=1}^{\infty} h^l P'_{l-1}(x).
\end{aligned}$$

Next, the right hand side of (8) becomes

$$(11) \quad h \frac{\partial}{\partial h} \sum_{l=0}^{\infty} h^l P_l(x) = \sum_{l=0}^{\infty} l h^l P_l(x).$$

Since (9) and (11) are equal, we have

$$(12) \quad \sum_{l=1}^{\infty} h^l [x P'_l(x) - P'_{l-1}(x)] = \sum_{l=0}^{\infty} l h^l P_l(x).$$

Thus, for each $l \in \mathbb{N}$, the summands must be equal. Therefore, we have

$$(13) \quad h^l [x P'_l(x) - P'_{l-1}(x)] = l h^l P_l(x).$$

Dividing both sides of (13) by h^l , we get

$$(14) \quad \boxed{x P'_l(x) - P'_{l-1}(x) = l P_l(x)},$$

the desired recursion relation.

5.

Differentiate the recursion relation (5.8a) and use the recursion relation (5.8b) with l replaced by $l - 1$ to prove the recursion relation (5.8c).

In the textbook, equation (5.8a) is the recursion relation

$$(15) \quad lP_l(x) = (2l - 1)xP_{l-1}(x) - (l - 1)P_{l-2}(x).$$

Differentiating (15) with respect to x gives

$$(16) \quad \begin{aligned} lP'_l(x) &= (2l - 1)P_{l-1}(x) + (2l - 1)xP'_{l-1}(x) - (l - 1)P'_{l-2}(x) \\ &= 2lP_{l-1}(x) - P_{l-1}(x) + 2lxP'_{l-1}(x) - xP'_{l-1}(x) - lP'_{l-2}(x) + P'_{l-2}(x). \end{aligned}$$

Recursion relation (5.8b) in the textbook is (once again)

$$(17) \quad xP'_l(x) - P'_{l-1}(x) = lP_l(x).$$

If we make the substitution $l \rightarrow l - 1$, (17) becomes

$$(18) \quad xP'_{l-1}(x) - P'_{l-2}(x) = (l - 1)P_{l-1}(x),$$

so that

$$(19) \quad \begin{aligned} xP'_{l-1}(x) &= (l - 1)P_{l-1}(x) + P'_{l-2}(x) \\ &= lP_{l-1}(x) - P_{l-1}(x) + P'_{l-2}(x). \end{aligned}$$

Using (19) in (16) gives

$$(20) \quad \begin{aligned} lP'_l(x) &= 2lP_{l-1}(x) - P_{l-1}(x) + 2l^2P_{l-1}(x) - 2lP_{l-1}(x) + 2lP'_{l-2}(x) \\ &\quad - lP_{l-1}(x) + P_{l-1}(x) - P'_{l-2}(x) - lP'_{l-2}(x) + P'_{l-2}(x) \\ &= (2l - 1 + 2l^2 - 2l - l + 1)P_{l-1}(x) + (2l - 1 - l + 1)P'_{l-2}(x) \\ &= (2l^2 - l)P_{l-1}(x) + lP'_{l-2}(x). \end{aligned}$$

Dividing by l on both sides of (20) gives

$$(21) \quad P'_l(x) = (2l - 1)P_{l-1}(x) + P'_{l-2}(x).$$

From (19) and (21) we get

$$(22) \quad \begin{aligned} P'_l(x) - xP'_{l-1}(x) &= (2l - 1)P_{l-1}(x) + P'_{l-2}(x) - lP_{l-1}(x) + P_{l-1}(x) - P'_{l-2}(x) \\ &= (2l - 1 - l + 1)P_{l-1}(x) + (1 - 1)P'_{l-2}(x) \\ &= lP_{l-1}(x). \end{aligned}$$

Thus, we get the desired recursion relation:

$$(23) \quad \boxed{P'_l(x) - xP'_{l-1}(x) = lP_{l-1}(x).}$$

12.

Express the polynomial $7x^4 - 3x + 1$ as a linear combination of Legendre polynomials. *Hint:* Start with the highest power of x and work down in finding the correct combination.

The first five Legendre polynomials are

$$(24) \quad P_0(x) = 1,$$

$$(25) \quad P_1(x) = x,$$

$$(26) \quad P_2(x) = \frac{1}{2}(3x^2 - 1),$$

$$(27) \quad P_3(x) = \frac{1}{2}(5x^3 - 3x), \quad \text{and}$$

$$(28) \quad P_4(x) = \frac{1}{8}(35x^4 - 30x^2 + 3).$$

Consider the field $\mathbb{R}[x]$ of polynomials in x with real coefficients and the ideal $\langle x^5 \rangle \subseteq \mathbb{R}[x]$. Then the quotient field $\mathbb{R}[x]/\langle x^5 \rangle$ consists of all cosets $p(x) + \langle x^5 \rangle$ with $p(x) \in \mathbb{R}[x]$. This can be thought of as a vector space over $\mathbb{R}/\langle x^5 \rangle$ since $\mathbb{R}/\langle x^5 \rangle$ is a subfield of $\mathbb{R}[x]/\langle x^5 \rangle$ (and since any field is a vector space over one of its subfields). Specifically, if we abandon the formalism of cosets, $\mathbb{R}[x]/\langle x^5 \rangle$ becomes a vector space (call it V) over \mathbb{R} whose vectors are polynomials in x with degree $n \leq 4$. The usual standard basis of V is

$$(29) \quad B = \{1, x, x^2, x^3, x^4\},$$

So that any polynomial $p(x) = \alpha_0 + \alpha_1x + \alpha_2x^2 + \alpha_3x^3 + \alpha_4x^4 \in V$ can be written as a coordinate vector relative to B :

$$(30) \quad [p(x)]_B = [\alpha_0 \ \alpha_1 \ \alpha_2 \ \alpha_3 \ \alpha_4]^T.$$

Following from (24), (25), (26), (27), and (28), the coordinate vector representations of the Legendre polynomials of degree 4 or less are:

$$(31) \quad [P_0(x)]_B = [1 \ 0 \ 0 \ 0 \ 0]^T,$$

$$(32) \quad [P_1(x)]_B = [0 \ 1 \ 0 \ 0 \ 0]^T,$$

$$(33) \quad [P_2(x)]_B = [-1/2 \ 0 \ 3/2 \ 0 \ 0]^T,$$

$$(34) \quad [P_3(x)]_B = [0 \ -3/2 \ 0 \ 5/2 \ 0]^T, \quad \text{and}$$

$$(35) \quad [P_4(x)]_B = [3/8 \ 0 \ -15/4 \ 0 \ 35/8]^T.$$

Let $p(x) = 7x^4 - 3x + 1$, the polynomial we are trying to express as a linear combination of Legendre polynomials. The coordinate vector of $p(x)$ is

$$(36) \quad [p(x)]_B = [1 \ -3 \ 0 \ 0 \ 7]^T.$$

Our problem now becomes finding coefficients $a_0, a_1, a_2, a_3, a_4 \in \mathbb{R}$ such that

$$(37) \quad p(x) = a_0P_0(x) + a_1P_1(x) + a_2P_2(x) + a_3P_3(x) + a_4P_4(x).$$

Thus, we get the following matrix equation that is equivalent to (37):

$$(38) \quad \begin{bmatrix} 1 & 0 & -1/2 & 0 & 3/8 \\ 0 & 1 & 0 & -3/2 & 0 \\ 0 & 0 & 3/2 & 0 & -15/4 \\ 0 & 0 & 0 & 5/2 & 0 \\ 0 & 0 & 0 & 0 & 35/8 \end{bmatrix} \begin{bmatrix} a_0 \\ a_1 \\ a_2 \\ a_3 \\ a_4 \end{bmatrix} = \begin{bmatrix} 1 \\ -3 \\ 0 \\ 0 \\ 7 \end{bmatrix}.$$

For the 5×5 sparse upper triangular matrix in (38), it is easy to find its inverse using elementary row reduction:

$$(39) \quad \begin{bmatrix} 1 & 0 & -1/2 & 0 & 3/8 & 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & -3/2 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 3/2 & 0 & -15/4 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 5/2 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 35/8 & 0 & 0 & 0 & 0 & 1 \end{bmatrix} \\ \rightarrow \begin{bmatrix} 1 & 0 & -1/2 & 0 & 3/8 & 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & -3/2 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & -5/2 & 0 & 0 & 2/3 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 2/5 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 8/35 \end{bmatrix} \\ \rightarrow \begin{bmatrix} 1 & 0 & -1/2 & 0 & 0 & 1 & 0 & 0 & -3/35 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 1 & 0 & 3/5 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 & 2/3 & 0 & 4/7 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 2/5 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 8/35 \end{bmatrix} \\ \rightarrow \begin{bmatrix} 1 & 0 & 0 & 0 & 0 & 1 & 0 & 1/3 & 0 & 7/35 \\ 0 & 1 & 0 & 0 & 0 & 0 & 1 & 0 & 3/5 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 & 2/3 & 0 & 4/7 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 2/5 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 8/35 \end{bmatrix}$$

Thus, the solution to the system in (38) is

$$(40) \quad \begin{bmatrix} a_0 \\ a_1 \\ a_2 \\ a_3 \\ a_4 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 1/3 & 0 & 7/35 \\ 0 & 1 & 0 & 3/5 & 0 \\ 0 & 0 & 2/3 & 0 & 4/7 \\ 0 & 0 & 0 & 2/5 & 0 \\ 0 & 0 & 0 & 0 & 8/35 \end{bmatrix} \begin{bmatrix} 1 \\ -3 \\ 0 \\ 0 \\ 7 \end{bmatrix} = \begin{bmatrix} 12/5 \\ -3 \\ 4 \\ 0 \\ 8/5 \end{bmatrix}.$$

Therefore, the Legendre polynomial expansion of $p(x) = 7x^4 - 3x + 1$ is

$$(41) \quad \boxed{7x^4 - 3x + 1 = \frac{12}{5}P_0(x) - 3P_1(x) + 4P_2(x) + \frac{8}{5}P_4(x)}.$$