

P3.2 at  $\theta_i = \theta_t = 0$

(1)

$$r_s = \frac{\sin(\theta_t - \theta_i)}{\sin(\theta_t + \theta_i)} = \frac{n_i \cos \theta_i - n_t \cos \theta_t}{n_i \cos \theta_i + n_t \cos \theta_t}$$

$$\lim_{\substack{\theta_i \rightarrow 0 \\ \theta_t \rightarrow 0}} r_s = \frac{n_i - n_t}{n_i + n_t} = - \left( \frac{n_t - n_i}{n_t + n_i} \right)$$

$$\lim_{\substack{\theta_i \rightarrow 0 \\ \theta_t \rightarrow 0}} r_p = \lim_{\substack{\theta_i \rightarrow 0 \\ \theta_t \rightarrow 0}} \frac{n_i \cos \theta_t - n_t \cos \theta_i}{n_i \cos \theta_t + n_t \cos \theta_i} = \frac{n_i - n_t}{n_i + n_t} \\ = - \left( \frac{n_t - n_i}{n_t + n_i} \right)$$

$$\lim_{\substack{\theta_i \rightarrow 0 \\ \theta_t \rightarrow 0}} t_s = \lim_{\theta_i, \theta_t \rightarrow 0} \frac{2n_i \cos \theta_i}{n_i \cos \theta_i + n_t \cos \theta_t} = \frac{2n_i}{n_i + n_t}$$

$$\lim_{\substack{\theta_i \rightarrow 0 \\ \theta_t \rightarrow 0}} t_p = \lim_{\theta_i, \theta_t \rightarrow 0} \frac{2n_i \cos \theta_i}{n_i \cos \theta_t + n_t \cos \theta_i} = \frac{2n_i}{n_i + n_t}$$

Why is there no difference between (P) & (S) for normal incidence?

for

$n_i$	$n_t$
air 1	glass 1.5

(2)

$$r_s = r_p = - \left( \frac{1.5 - 1}{1.5 + 1} \right) = - \left( \frac{0.5}{2.5} \right) = -0.2$$

$$t_s = t_p = \frac{2(1)}{1.5 + 1} = \frac{2}{2.5} = \frac{4}{5}$$

in general.

$$R_s = \frac{(n_t - n_i)^2}{(n_t + n_i)^2} \quad T_s = \frac{4n_i^2}{(n_i + n_t)^2}$$


$$R_s + T_s = ?$$

$$= \frac{(n_t - n_i)^2 + 4n_i^2}{(n_t + n_i)^2} = \frac{n_t^2 + 2n_i n_t + n_i^2}{(n_t + n_i)^2}$$

$$= 1 \quad \checkmark$$

$$R_p + T_p = ?$$

(3)



$$r_s = \frac{\sin(\theta_t - \theta_i)}{\sin(\theta_t + \theta_i)} ;$$

We get  $\theta_t$  from Snell's law

$$n_i \sin \theta_i = n_t \sin \theta_t$$


---

# ch4 Multiple Parallel Interfaces

(19)

$n_i$     $n_t$

$r_s, t_s, r_p, t_p$   
 $R_s, T_s, R_p, T_p$




two interfaces

The goal is to find:  $r^{tot}, t^{tot}$   
 $R^{tot}, T^{tot}$

# 4.1 Double Interface / Fresnel Coeffs

(2a)



Assumptions:  $n_0, n_1, n_2$ ; Real (no absorption)

Note:  $E_{0\leftarrow}^{(s)}, E_{0\leftarrow}^{(p)}$   
 $E_{2\rightarrow}^{(s)}, E_{2\rightarrow}^{(p)}$  } include light from middle layer.

$$r_s = \frac{\sin(\theta_t - \theta_i)}{\sin(\theta_t + \theta_i)} ; t_s = \frac{2 \sin \theta_t \cos \theta_i}{\sin(\theta_t + \theta_i)} \quad (3)$$

$$\left\{ \begin{aligned} r_p &= \frac{\tan(\theta_t - \theta_i)}{\tan(\theta_t + \theta_i)} ; t_p = \frac{2 \sin \theta_t \cos \theta_i}{\sin(\theta_t + \theta_i) \cos(\theta_t - \theta_i)} \\ r_p &= \frac{\sin \theta_t \cos \theta_t - \sin \theta_i \cos \theta_i}{\sin \theta_t \cos \theta_t + \sin \theta_i \cos \theta_i} \end{aligned} \right.$$

$$r_s^{0 \rightarrow 1} = \frac{\sin \theta_1 \cos \theta_0 - \sin \theta_0 \cos \theta_1}{\sin \theta_1 \cos \theta_0 + \sin \theta_0 \cos \theta_1}$$

$$r_p^{0 \rightarrow 1} = \frac{\sin \theta_1 \cos \theta_1 - \sin \theta_0 \cos \theta_0}{\sin \theta_1 \cos \theta_1 + \sin \theta_0 \cos \theta_0}$$


$$t_s^{0 \rightarrow 1} = \frac{2 \sin \theta_1 \cos \theta_0}{\sin \theta_1 \cos \theta_0 + \sin \theta_0 \cos \theta_1}$$

$$t_p^{0 \rightarrow 1} = \frac{2 \sin \theta_1 \cos \theta_0}{\sin \theta_1 \cos \theta_1 + \sin \theta_0 \cos \theta_0}$$

0 → 1  
From left  
to  
middle  
layer.

From Middle layer to first interface:

$$\left\{ \begin{aligned} r_s^{0 \leftarrow 1} &= -r_s^{0 \rightarrow 1} ? \\ r_p^{0 \leftarrow 1} &= -r_p^{0 \rightarrow 1} ? \\ t_s^{0 \leftarrow 1} &= \frac{2 \sin \theta_0 \cos \theta_1}{\sin \theta_0 \cos \theta_1 + \sin \theta_1 \cos \theta_0} \end{aligned} \right. \quad \left. \begin{aligned} t_p^{0 \leftarrow 1} &= \frac{2 \sin \theta_0 \cos \theta_1}{\sin \theta_0 \cos \theta_0 + \sin \theta_1 \cos \theta_1} \end{aligned} \right.$$



At second interface:

(4)


$$r_s^{1 \rightarrow 2} = \frac{\sin \theta_2 \cos \theta_1 - \sin \theta_1 \cos \theta_2}{\sin \theta_2 \cos \theta_1 + \sin \theta_1 \cos \theta_2}$$

$$r_p^{1 \rightarrow 2} = \frac{\sin \theta_2 \cos \theta_2 - \sin \theta_1 \cos \theta_1}{\sin \theta_2 \cos \theta_2 + \sin \theta_1 \cos \theta_1}$$

$$t_s^{1 \rightarrow 2} = \frac{2 \sin \theta_2 \cos \theta_1}{\sin \theta_2 \cos \theta_1 + \sin \theta_1 \cos \theta_2}$$

$$t_p^{1 \rightarrow 2} = \frac{2 \sin \theta_2 \cos \theta_1}{\sin \theta_2 \cos \theta_2 + \sin \theta_1 \cos \theta_1}$$

For the s-polarized case:




Using Fresnel coefficients.

$$E_{1 \rightarrow}^{(s)} = t_s^{0 \rightarrow 1} E_{0 \rightarrow}^{(s)} + r_s^{0 \leftarrow 1} E_{1 \leftarrow}^{(s)} \quad (4.4)$$

Similarly  $E_{0 \leftarrow}^{(s)} = r_s^{0 \rightarrow 1} E_{0 \rightarrow}^{(s)} + t_s^{0 \leftarrow 1} E_{1 \leftarrow}^{(s)}$  (4.5)

Due to thickness of middle region, we need to worry about phase shifts.



at  $(y, z) = (0, d)$  the adjusted field is  $E_{1 \rightarrow}^{(s)} e^{i \vec{k}_1 \cdot \vec{r}} = E_{1 \rightarrow}^{(s)} e^{i k_1 d \cos \theta_1}$

$$E_{2 \rightarrow}^{(s)} = t_s^{1 \rightarrow 2} E_{1 \rightarrow}^{(s)} e^{i k_1 d \cos \theta_1} \quad (4.6)$$



Similarly

$$E_{1\leftarrow}^{(s)} = E_{1\rightarrow}^{(s)} e^{ik_1 d \cos \theta_1} r_s^{1\rightarrow 2} e^{ik_1 d \cos \theta_1}$$



a transmission through middle layer  
 + reflection from second interface  
 + a transmission through middle layer back to first interface.

Example 4.1 We have 4 unknowns for s-polar.

$$E_{0\leftarrow}^s ; E_{1\rightarrow}^s ; E_{1\leftarrow}^s ; E_{2\rightarrow}^s$$

Find  $E_s^{\text{tot}} = \frac{E_{2\rightarrow}^s}{E_{0\rightarrow}^s} ?$

(4.8) 1)  $E_{1\rightarrow}^s = \frac{E_{2\rightarrow}^s}{t_s^{1\rightarrow 2}} e^{-ik_1 d \cos \theta_1}$  from (4.6)

(4.9) 2) sub into 4.7

$$\begin{aligned} E_{1\leftarrow}^s &= E_{1\rightarrow}^s e^{ik_1 d \cos \theta_1} r_s^{1\rightarrow 2} e^{ik_1 d \cos \theta_1} \\ &= \frac{E_{2\rightarrow}^s}{t_s^{1\rightarrow 2}} e^{-ik_1 d \cos \theta_1} e^{ik_1 d \cos \theta_1} r_s^{1\rightarrow 2} e^{ik_1 d \cos \theta_1} \end{aligned}$$

sub 4.8 + 4.9 into 4.4

(7)

$$E_{1 \rightarrow}^s = t_s^{0 \rightarrow 1} E_{0 \rightarrow}^s + r_s^{0 \leftarrow 1} E_{1 \leftarrow}^s$$



$$\frac{E_{2 \rightarrow}^s}{t_s^{1 \rightarrow 2}} e^{-ik_1 d \cos \theta_1} = t_s^{0 \rightarrow 1} E_{0 \rightarrow}^s + r_s^{0 \leftarrow 1} E_{2 \rightarrow}^{(s)} \frac{r_s^{1 \rightarrow 2} e^{ik_1 d}}{t_s^{1 \rightarrow 2}}$$

$$t_s^{1 \rightarrow 2} = \frac{E_{2 \rightarrow}^s}{E_{0 \rightarrow}^s} = \frac{t_s^{0 \rightarrow 1} e^{ik_1 d \cos \theta_1}}{1 - r_s^{0 \leftarrow 1} r_s^{1 \rightarrow 2} e^{ik_1 d \cos \theta_1}}$$