

# Chapter 1

# Chemical Measurements

---

# Overview

---

1-1 SI Units

1-2 Chemical Concentrations

1-3 Preparing Solutions

1-4 Stoichiometry Calculations for Gravimetric Analysis

# SI Units

---

SI base units include the following:

- meter (m)
- kilogram (kg)
- second (s)
- ampere (A)
- kelvin (K)
- mole (mol)

# Derived SI Units

---

Other quantities can be derived in terms of base SI units. See Table 1-2.

- force (newton, N),  $\text{Kg}\cdot\text{m} / \text{s}^2$
- pressure (pascal, Pa),  $\text{N} / \text{m}^2$
- energy (joule, J),  $\text{N}\cdot\text{m}$

# Prefixes

**TABLE 1-3** Prefixes

Prefix	Symbol	Factor	Prefix	Symbol	Factor
yotta	Y	$10^{24}$	deci	d	$10^{-1}$
zetta	Z	$10^{21}$	centi	c	$10^{-2}$
exa	E	$10^{18}$	milli	m	$10^{-3}$
peta	P	$10^{15}$	micro	$\mu$	$10^{-6}$
tera	T	$10^{12}$	nano	n	$10^{-9}$
giga	G	$10^9$	pico	p	$10^{-12}$
mega	M	$10^6$	femto	f	$10^{-15}$
kilo	k	$10^3$	atto	a	$10^{-18}$
hecto	h	$10^2$	zepto	z	$10^{-21}$
deca	da	$10^1$	yocto	y	$10^{-24}$

Table 1.3

Memorize these prefixes with their abbreviations and powers of ten.

# Using Prefixes in Calculations

---

- In calculations, units should be carried along with the numbers.
- Prefixes such as kilo- and milli- are used to denote multiples of units.
- Prefixes can be used to simplify conversions between units.

## Example:

Express 19.3 mPa in terms of atm

19.3 mPa can be written as  $19.3 \times 10^{-3}$  Pa

$$19.3 \times 10^{-3} \text{ Pa} \times \frac{1.00 \text{ atm}}{101325 \text{ Pa}} = 1.90 \times 10^{-7} \text{ atm}$$

# Concentrations

---

- **Molarity** (moles of solute per liter of solution), M
- **Molality** (moles of solute per kilogram of solvent), m
- **Formal concentration** (formula units per liter), F
- **Percent composition** (w/w, v/v or w/v), %
- **Parts per million**, ppm
- **Parts per billion**, ppb

# % Composition (w/w or v/v)

---

$$\text{Weight percent} = \frac{\text{mass of solute}}{\text{mass of total solution or mixture}} \times 100$$

$$\text{Volume percent} = \frac{\text{volume of solute}}{\text{volume of total solution}} \times 100$$

95% (w/w) ethanol contains 95 g of ethanol for every 100 g of solution.



# ppm, Parts per Million

---

$$\text{ppm} = \frac{\text{mass of substance}}{\text{mass of sample}} \times 10^6$$

$$\text{ppb} = \frac{\text{mass of substance}}{\text{mass of sample}} \times 10^9$$

- **Shortcut:** ppm is also mg/L or  $\mu\text{g/mL}$  if the density of the solution is 1.0 g/mL.
- An aqueous solution that is 1 000 ppm in Cu contains 1 000 mg Cu per liter of solution.
- It also contains 1000  $\mu\text{g}$  per mL of solution.

# ppm, Parts per Million

## EXAMPLE Converting Parts per Billion into Molarity

Normal alkanes are hydrocarbons with the formula  $C_nH_{2n+2}$ . Plants selectively synthesize alkanes with an odd number of carbon atoms. The concentration of  $C_{29}H_{60}$  in summer rainwater collected in Hannover, Germany, is 34 ppb. Find the molarity of  $C_{29}H_{60}$  and express the answer with a prefix from Table 1-3.

**Solution** A concentration of 34 ppb means there are 34 ng of  $C_{29}H_{60}$  per gram of rainwater, which is nearly the same as 34 ng/mL because the density of rainwater is close to 1.00 g/mL. To find the molarity, we need to know how many grams of  $C_{29}H_{60}$  are contained in a liter. Multiplying nanograms and milliliters by 1 000 gives 34  $\mu\text{g}$  of  $C_{29}H_{60}$  per liter of rainwater:

$$\frac{34 \text{ ng } C_{29}H_{60}}{\text{mL}} \left( \frac{1\,000 \text{ mL/L}}{1\,000 \text{ ng}/\mu\text{g}} \right) = \frac{34 \mu\text{g } C_{29}H_{60}}{\text{L}}$$

The molecular mass of  $C_{29}H_{60}$  is  $29 \times 12.011 + 60 \times 1.008 = 408.8 \text{ g/mol}$ , so the molarity is

$$\text{Molarity of } C_{29}H_{60} \text{ in rainwater} = \frac{34 \times 10^{-6} \text{ g/L}}{408.8 \text{ g/mol}} = 8.3 \times 10^{-8} \text{ M}$$

An appropriate prefix from Table 1-3 would be nano (n), which is a multiple of  $10^{-9}$ :

$$8.3 \times 10^{-8} \text{ M} \left( \frac{1 \text{ nM}}{10^{-9} \text{ M}} \right) = 83 \text{ nM}$$

# ppm, Parts per Million

## EXAMPLE Converting Parts per Billion into Molarity

Normal alkanes are hydrocarbons with the formula  $C_nH_{2n+2}$ . Plants selectively synthesize alkanes with an odd number of carbon atoms. The concentration of  $C_{29}H_{60}$  in summer rainwater collected in Hannover, Germany, is 34 ppb. Find the molarity of  $C_{29}H_{60}$  and express the answer with a prefix from Table 1-3.

**Solution** A concentration of 34 ppb means there are 34 ng of  $C_{29}H_{60}$  per gram of rainwater, which is nearly the same as 34 ng/mL because the density of rainwater is close to 1.00 g/mL. To find the molarity, we need to know how many grams of  $C_{29}H_{60}$  are contained in a liter. Multiplying nanograms and milliliters by 1 000 gives 34  $\mu\text{g}$  of  $C_{29}H_{60}$  per liter of rainwater:

$$\frac{34 \text{ ng } C_{29}H_{60}}{\text{mL}} \left( \frac{1\,000 \text{ mL/L}}{1\,000 \text{ ng}/\mu\text{g}} \right) = \frac{34 \mu\text{g } C_{29}H_{60}}{\text{L}}$$

# ppm, Parts per Million

---

The molecular mass of  $C_{29}H_{60}$  is  $29 \times 12.011 + 60 \times 1.008 = 408.8 \text{ g/mol}$ , so the molarity is

$$\text{Molarity of } C_{29}H_{60} \text{ in rainwater} = \frac{34 \times 10^{-6} \text{ g/L}}{408.8 \text{ g/mol}} = 8.3 \times 10^{-8} \text{ M}$$

An appropriate prefix from Table 1-3 would be nano (n), which is a multiple of  $10^{-9}$ :

$$8.3 \times 10^{-8} \text{ M} \left( \frac{1 \text{ nM}}{10^{-9} \text{ M}} \right) = 83 \text{ nM}$$

# Making Solutions

---

- To calculate quantities of reagents needed to prepare solutions, use the relation

$$(M_{\text{conc}})(V_{\text{conc}}) = (M_{\text{dil}})(V_{\text{dil}})$$

- Equates moles of reagent removed from a stock solution to moles delivered into a new solution.

# Making Solutions

---

- How many **moles** of  $\text{CuCl}_2 \cdot 5\text{H}_2\text{O}$  are needed to make 500 mL of a solution that is 1 000.0 ppm in Cu?
- **grams** of  $\text{CuCl}_2 \cdot 5\text{H}_2\text{O}$ ?
- **milliliters** of 10 000 ppm  $\text{CuCl}_2 \cdot 5\text{H}_2\text{O}$  solution?



Figure 1-4

# Making Solutions

- How many **grams** of  $\text{CuCl}_2 \cdot 5\text{H}_2\text{O}$  are needed to make 500 mL of a solution that is 1 000 ppm in Cu? (MM of  $\text{CuCl}_2 \cdot 5\text{H}_2\text{O}$  is 224.53 g/mol)

- First calculate the grams of Cu required
  - $(500\text{-mL}) \times (1000 \mu\text{g/mL}) = 5.0 \times 10^5 \mu\text{g Cu}$
  - $5.0 \times 10^5 \mu\text{g Cu} = (5.0 \times 10^5)(10^{-6})\text{g Cu}$
  - $5.0 \times 10^{-1}\text{g Cu}$



$$\left(5.0 \times 10^{-1} \text{g Cu}\right) \times \left(\frac{224.53 \text{ g CuCl}_2 \cdot 5\text{H}_2\text{O} / \text{mol}}{63.546 \text{ g Cu} / \text{mol}}\right) = \mathbf{1.76_7 \text{ g CuCl}_2 \cdot 5\text{H}_2\text{O}}$$

# Making Solutions

- How many **milliliters** of  $\text{CuCl}_2 \cdot 5\text{H}_2\text{O}$  are needed to make 500 mL of a solution that is 1 000.0 ppm in Cu?
- **moles** of  $\text{CuCl}_2 \cdot 5\text{H}_2\text{O}$  solution?

$$(M_{\text{conc}})(V_{\text{conc}}) = (M_{\text{dil}})(V_{\text{dil}})$$

$$(10\,000\text{ ppm})(V_{\text{conc}}) = (1\,000\text{ ppm})(500\text{ mL})$$

$$V_{\text{conc}} = (1\,000\text{ ppm})(500\text{ mL}) / (10\,000\text{ ppm})$$

$$V_{\text{conc}} = \mathbf{50\text{ mL}}$$



Figure 1-4

Place 50 mL of 10 000 ppm  $\text{CuCl}_2$  stock solution in the 500 mL flask and fill to the 500 mL mark!



# Making Solutions

- How many milliliters of concentrated HCl are needed to make 500 mL of a solution that is 0.250 M in HCl?

$$(M_{\text{conc}})(V_{\text{conc}}) = (M_{\text{dil}})(V_{\text{dil}})$$

- Need to know molarity of concentrated HCl.



Figure 1-4

Concentrated HCl is  
37.2% (w/w) HCl and has  
a density of 1.188 g/cm<sup>3</sup>

# Making Solutions

- How many milliliters of concentrated HCl are needed to make 500 mL of a solution that is 0.250 M in HCl?

$$M_{\text{conc. HCl}} = ?$$

$$(M_{\text{conc}})(V_{\text{conc}}) = (M_{\text{dil}})(V_{\text{dil}})$$



Figure 1-4

# Molarity of the Conc. HCl Solution

---

$$c = 37.2\% = 37.2 \text{ gHCl} / 100 \text{ g}_{\text{soln}}$$

$$\text{MM} = 36.46 \text{ g/mol}$$

$$\rho = 1.188 \text{ g/cm}^3$$

} desired units  
 $\frac{\text{mol HCl}}{\text{L soln}}$

$$\frac{37.2 \text{ g HCl}}{100 \text{ g}_{\text{soln}}} \times \frac{1 \text{ mol HCl}}{36.46 \text{ g HCl}} \times \frac{1.188 \text{ g}_{\text{soln}}}{1.000 \text{ cm}^3_{\text{soln}}} \times \frac{1000 \text{ cm}^3_{\text{soln}}}{1 \text{ L}}$$
$$= 12.1 \text{ mol/L}$$

# Making Solutions

- How many **milliliters** of concentrated HCl are needed to make 500 mL of a solution that is 0.250 M in HCl?

$$M_{\text{conc. HCl}} = 12.1 \text{ M}$$

$$(M_{\text{conc}})(V_{\text{conc}}) = (M_{\text{dil}})(V_{\text{dil}})$$

$$(12.1 \text{ M})(V_{\text{conc}}) = (0.250 \text{ M})(500 \text{ mL})$$

$$V_{\text{conc}} = (0.250 \text{ M})(500 \text{ mL}) / (12.1 \text{ M})$$

$$V_{\text{conc}} = \mathbf{10.3_3 \text{ mL}}$$



Figure 1-4

# Stoichiometry

---

- Use stoichiometry relationships to calculate required masses or volumes of reagents for chemical reactions.
- From the mass of product of a reaction, you should be able to compute how much reactant was consumed.


# Stoichiometry

Iron from a dietary supplement tablet can be measured gravimetrically by dissolving the tablet and then converting the dissolved iron into solid  $\text{Fe}_2\text{O}_3$ . The mass of  $\text{Fe}_2\text{O}_3$  tells us the mass of iron in the original tablet.


Here are the steps in the procedure:

**Step 1** Tablets containing iron(II) fumarate ( $\text{Fe}^{2+}\text{C}_4\text{H}_2\text{O}_4^{2-}$ ) and inert binder are mixed with 150 mL of 0.100 M HCl to dissolve the  $\text{Fe}^{2+}$ . The mixture is filtered to remove insoluble binder.

**Step 2** Iron(II) in the clear liquid is oxidized to iron(III) with excess hydrogen peroxide:



**Step 3** Ammonium hydroxide is added to precipitate hydrous iron(III) oxide, which is a gel. The gel is filtered and heated in a furnace to convert it to pure solid  $\text{Fe}_2\text{O}_3$ .



# Stoichiometry

## EXAMPLE How Many Tablets Should We Analyze?

In a gravimetric analysis, we need enough product to weigh accurately. Each tablet provides ~15 mg of iron. How many tablets should we analyze to provide 0.25 g of  $\text{Fe}_2\text{O}_3$  product?

**Solution** We can answer the question if we know how many grams of iron are in 0.25 g of  $\text{Fe}_2\text{O}_3$ . The formula mass of  $\text{Fe}_2\text{O}_3$  is 159.69 g/mol, so 0.25 g is equal to

$$\text{mol Fe}_2\text{O}_3 = \frac{0.25 \text{ g}}{159.69 \text{ g/mol}} = 1.6 \times 10^{-3} \text{ mol}$$

Each mol of  $\text{Fe}_2\text{O}_3$  has 2 mol of Fe, so 0.25 g of  $\text{Fe}_2\text{O}_3$  contains

$$1.6 \times 10^{-3} \text{ mol Fe}_2\text{O}_3 \times \frac{2 \text{ mol Fe}}{1 \text{ mol Fe}_2\text{O}_3} = 3.2 \times 10^{-3} \text{ mol Fe}$$

The mass of Fe is

$$3.2 \times 10^{-3} \text{ mol Fe} \times \frac{55.845 \text{ g Fe}}{\text{mol Fe}} = 0.18 \text{ g Fe}$$

If each tablet contains 15 mg Fe, the number of tablets required is

$$\text{Number of tablets} = \frac{0.18 \text{ g Fe}}{0.015 \text{ g Fe/tablet}} = 12 \text{ tablets}$$

# Limiting Reagent

---

- The **limiting reagent** in a chemical reaction is the one that is consumed first.
- Once the limiting reagent is gone, the reaction ceases.

**Example:** For the reaction:  $A + 2B \rightarrow P$

0.751 moles of A are mixed with 1.43 moles of B.


- What is the limiting reagent and how much excess reagent remains unreacted?



# Limiting Reagent

---

Solution:



If A is the L.R., how much B is required to completely consume A?

$$0.751 \text{ mol A} \times \left( \frac{2 \text{ mol B}}{1 \text{ mol A}} \right) = \mathbf{1.502 \text{ mol B required}}$$

However, we only have 1.43 mol B, so B must be the L.R.

# Limiting Reagent

---

**Solution:**

	A	+	2B	→	P
Initial	0.751		1.43		0
Final	?		0		

If B is the L.R., how much is required to completely consume A?


$$0.1.43 \text{ mol B} \times \left( \frac{1 \text{ mol A}}{2 \text{ mol B}} \right) = \mathbf{0.715 \text{ mol A}}$$
 react with B.

So,  $0.751 \text{ mol A} - 0.715 \text{ mol A} = \mathbf{0.036 \text{ mol A}}$  remain after the reaction is complete. All of B is consumed.

# Limiting Reagent

## EXAMPLE Limiting Reagent

Reaction 1-9 requires one mole of oxalate for each mole of calcium.



If you mix 1.00 g of  $\text{CaCl}_2$  (FM 110.98) with 1.15 g of  $\text{Na}_2\text{C}_2\text{O}_4$  (FM 134.00) in water, which is the limiting reagent? What fraction of the nonlimiting reagent is left over?

**Solution** The available moles of each reagent are

$$\frac{1.00 \text{ g CaCl}_2}{110.98 \text{ g/mol}} = 9.01 \text{ mmol Ca}^{2+} \qquad \frac{1.15 \text{ g Na}_2\text{C}_2\text{O}_4}{134.00 \text{ g/mol}} = 8.58 \text{ mmol C}_2\text{O}_4^{2-}$$

The reaction requires 1 mol  $\text{Ca}^{2+}$  for 1 mol  $\text{C}_2\text{O}_4^{2-}$ , so oxalate will be used up first. The  $\text{Ca}^{2+}$  remaining is  $9.01 - 8.58 = 0.43$  mmol. The fraction of unreacted  $\text{Ca}^{2+}$  is  $(0.43 \text{ mmol}/9.01 \text{ mmol}) = 4.8\%$

**TEST YOURSELF** The reaction  $5\text{H}_2\text{C}_2\text{O}_4 + 2\text{MnO}_4^- + 6\text{H}^+ \rightarrow 10\text{CO}_2 + 2\text{Mn}^{2+} + 8\text{H}_2\text{O}$  requires 5 mol  $\text{H}_2\text{C}_2\text{O}_4$  for 2 mol  $\text{MnO}_4^-$ . If you mix 1.15 g  $\text{Na}_2\text{C}_2\text{O}_4$  (FM 134.00) with 0.60 g  $\text{KMnO}_4$  (FM 158.03) and excess aqueous acid, which reactant is limiting? How much  $\text{CO}_2$  is produced?