

Organic Chemistry, *Fourth Edition*

Janice Gorzynski Smith
University of Hawai'i

Chapter 2 **Acids and Bases**

Prepared by Layne A. Morsch
The University of Illinois - Springfield

Copyright © 2014 The McGraw-Hill Companies, Inc. Permission required for reproduction or display.

1

Chapter 2 Acids and Bases

- 2.1 Brønsted-Lowry Acids and Bases
- 2.2 Reactions of Brønsted-Lowry Acids and Bases
- 2.3 Acid Strength and pK_a
- 2.4 Predicting the Outcome of Acid-Base Reactions
- 2.5 Factors That Determine Acid Strength
- 2.6 Common Acids and Bases
- 2.7 Aspirin
- 2.8 Lewis Acids and Bases

Copyright © 2014 The McGraw-Hill Companies, Inc. Permission required for reproduction or display.

2

2.1 Brønsted-Lowry Acids and Bases

- A hydrogen atom without its electron is a proton.
 H^+ = proton
- A **Brønsted-Lowry acid** is a proton donor.
 • It must have a proton.
- A **Brønsted-Lowry base** is a proton acceptor.
 • It must be able to form a bond to a proton.

3

Brønsted-Lowry Acids and Bases

Figure 2.1

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display

Brønsted-Lowry acids [H – A]

Inorganic	Organic
HCl	CH ₃ CO ₂ H
H ₂ SO ₄	acetic acid
HSO ₄ ⁻	
H ₂ O	OH
H ₃ O ⁺	HO ₂ CCH ₂ –C–CH ₂ CO ₂ H
	COOH
	citric acid

- All Brønsted-Lowry acids contain a proton.
- The net charge may be zero, (+), or (-).

Brønsted-Lowry bases [B:]


Inorganic	Organic
H ₂ O:	CH ₃ NH ₂
:NH ₃	methylamine
	CH ₃ O ⁻
	methoxide
	CH ₃
	C=O
	CH ₃
	acetone
	CH ₂ =CH ₂
	ethylene

- All Brønsted-Lowry bases contain a lone pair of electrons or a π bond.
- The net charge may be zero or (-).

4

Acidic and Basic Sites in Morphine

Some molecules contain both hydrogen atoms and lone pairs and thus, can act either as acids or bases, depending on the particular reaction.




5

2.1 Reactions of Brønsted-Lowry Acids and Bases

- A Brønsted-Lowry acid base reaction results in the transfer of a proton from an acid to a base.
- The electron pair of the base $B:$ forms a new bond to the proton of the acid forming the **conjugate acid** of the base.
- The acid $H-A$ loses a proton, leaving the electron pair in the $H-A$ bond on A . This forms the **conjugate base** of the acid.


Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display



6

Reactions of Brønsted-Lowry Acids and Bases


- The movement of electrons in reactions can be illustrated using **curved arrow** notation.
- Because two electron pairs are involved in this reaction, two curved arrows are needed.



7

Examples of Brønsted-Lowry Acid-Base Reactions

- A double reaction arrow (indicating equilibrium) is used between starting materials and products to indicate that the reaction can proceed in the forward and reverse directions.



8

General Reactivity

These reactions illustrate a general pattern of reactivity:


- **Electron-rich** species react with **electron-deficient** species
 - Bases are electron rich
 - While the H of an acid has a partial positive charge making it electron deficient
- Therefore, acids and bases react with one another

9

2.3 Acid Strength and pK_a

- Acid strength is the tendency of an acid to donate a proton.
- The more readily a compound donates a proton, the stronger an acid it is.

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display



10

Acid-Base Equilibrium

- Acidity is measured by an **equilibrium constant**.
- When a Brønsted-Lowry acid H-A is dissolved in water, an acid-base reaction occurs, and an **equilibrium constant** can be written for the reaction.

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display

$$\text{Equilibrium constant } K_{\text{eq}} = \frac{[\text{products}]}{[\text{starting materials}]} = \frac{[\text{H}_3\text{O}^+][\text{A}^-]}{[\text{H-A}][\text{H}_2\text{O}]}$$

11

Acidity Constant

- Because the concentration of the solvent H₂O is essentially constant, the equation can be rearranged and a new equilibrium constant, called the **acidity constant**, K_a , can be defined.

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display

$$\text{Acidity constant } = K_a = [\text{H}_2\text{O}]K_{\text{eq}} = \frac{[\text{H}_3\text{O}^+][\text{A}^-]}{[\text{H-A}]}$$

12


K_a and pK_a

- It is generally more convenient when describing acid strength to use " pK_a " values than K_a values.

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display

Definition: $pK_a = -\log K_a$

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display



13

Acidity of Some Common Compounds

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display


Table 2.1 Selected pK_a Values

	Acid	p <i>K</i> _a	Conjugate base	
↑ Increasing acidity	H-Cl	-7	Cl ⁻	↓ Increasing basicity
	CH ₃ COO-H	4.8	CH ₃ COO ⁻	
	HO-H	15.7	HO ⁻	
	CH ₃ CH ₂ O-H	16	CH ₃ CH ₂ O ⁻	
	HC≡CH	25	HC≡C ⁻	
	H-H	35	H ⁻	
	H ₂ N-H	38	H ₂ N ⁻	
	CH ₂ =CH ₂	44	CH ₂ =C ⁻ H	
	CH ₃ -H	50	CH ₃ ⁻	

14

2.4 Outcome of Acid-Base Reactions


- The position of the equilibrium depends on the relative strengths of the acids and bases.
- Equilibrium always favors formation of the weaker acid and base.



- Because the $\text{p}K_a$ of the starting acid (25) is lower than that of the conjugate acid (38), equilibrium favors the products.

15

Steps in Solving Acid-Base Reaction Equilibria



- Step [1]** Identify the acid and base in the starting materials. Assume :NH_2^- is the base because it bears a net negative charge. That makes $\text{HC}\equiv\text{CH}$ the acid.
- Step [2]** Draw the products of proton transfer and identify the conjugate acid and base in the products. Acetylene gives up its proton to :NH_2^- .
- Step [3]** Compare the $\text{p}K_a$ values of the acid and the conjugate acid. Equilibrium favors formation of the weaker acid with the higher $\text{p}K_a$. The $\text{p}K_a$ of NH_3 is higher; therefore products are favored.

16

2.5 Factors that Determine Acid Strength

- Anything that stabilizes a conjugate base $A:^-$ makes the starting acid H-A more acidic.
- Four factors affect the acidity of H-A. These are:
 - Element effects
 - Inductive effects
 - Resonance effects
 - Hybridization effects

17


Comparing the Acidity of Any Two Acids

- Always draw the conjugate bases.
- Determine which conjugate base is more stable.
- The **more stable** the conjugate base, the **more acidic** the acid.

18

Element Effects—Trends in the Periodic Table

Why does H₂O have a much lower pK_a than CH₄?




Since oxygen is much more electronegative than carbon, it more readily accepts a negative charge.


19

Element Effects—Trends in the Periodic Table

Across a row of the periodic table, the acidity of H-A increases as the electronegativity of A increases.



Positive or negative charge is stabilized when it is spread over a larger volume.




20

Element Effects Down a Column in the Periodic Table

- Down a column of the periodic table, size, and not electronegativity, determines acidity.
- The acidity of H-A increases as the size of A increases.

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display



21

Inductive Effects


- An **inductive effect** is the pull of electron density through σ bonds caused by electronegativity differences of atoms.
- More electronegative atoms stabilize regions of high electron density by an electron withdrawing inductive effect.
- The more electronegative the atom and the closer it is to the site of the negative charge, the greater the effect.
- The acidity of H-A increases with the presence of electron withdrawing groups in A.

22

Inductive Effects in Trifluoroethanol

- In the example below, note that 2,2,2-trifluoroethanol is more acidic than ethanol.

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display




- The reason for the increased acidity of 2,2,2-trifluoroethanol is that the three electronegative fluorine atoms stabilize the negatively charged conjugate base.


23

Rationale for Inductive Effects

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display



No additional electronegative atoms stabilize the conjugate base.




CF_3 withdraws electron density, stabilizing the conjugate base.



Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display



Figure 2.2



The dark red of the O atom indicates a region of high electron density.




The O atom is yellow, indicating it is less electron rich.

24

Resonance Effects

- Delocalization of charge through resonance influences acidity.
- Acetic acid is more acidic than ethanol, even though both conjugate bases have the negative charge on the same element, O.

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display




25

Comparison of Ethoxide and Acetate ions


- The conjugate base of ethanol has a localized charge.

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display



- The conjugate base of acetic acid is resonance delocalized.

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display




26


Electrostatic Potential Plots of Ethoxide and Acetate

Figure 2.3

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display



The negative charge is concentrated on the single oxygen atom, making this anion *less stable*.




The negative charge is delocalized over both oxygen atoms, making this anion *more stable*.

27


Hybridization Effects

Consider the relative acidities of three different compounds containing C-H bonds.

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display



Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display




28

Stability of Conjugate Bases

The higher the percent of *s*-character of the hybrid orbital, the more stable the conjugate base.

Figure 2.4



- As the lone pair of electrons is pulled closer to the nucleus, the negatively charged carbon appears less intensely red.

29

Summary of Factors that Determine Acid Strength

Figure 2.5

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display

Factor	Example																	
1. Element effects: The acidity of H–A increases both left-to-right across a row and down a column of the periodic table.	<div style="text-align: center;"> <p>Increasing acidity →</p> <table style="margin: auto;"> <tr> <td style="text-align: center;"> —C—H </td> <td style="text-align: center;">—N—H </td> <td style="text-align: center;">—O—H</td> <td style="text-align: center;">H—F</td> </tr> <tr> <td></td> <td></td> <td style="text-align: center;">—S—H</td> <td style="text-align: center;">H—Cl</td> </tr> <tr> <td></td> <td></td> <td></td> <td style="text-align: center;">H—Br</td> </tr> <tr> <td></td> <td></td> <td></td> <td style="text-align: center;">H—I</td> </tr> </table> <p style="text-align: right;">↑ Increasing acidity</p> </div>		 —C—H 	—N—H 	—O—H	H—F			—S—H	H—Cl				H—Br				H—I
 —C—H 	—N—H 	—O—H	H—F															
		—S—H	H—Cl															
			H—Br															
			H—I															
2. Inductive effects: The acidity of H–A increases with the presence of electron-withdrawing groups in A.	CH ₃ CH ₂ O–H	CF ₃ CH ₂ O–H more acidic																
3. Resonance effects: The acidity of H–A increases when the conjugate base A: [–] is resonance stabilized.	CH ₃ CH ₂ O–H	CH ₃ COO–H more acidic																
4. Hybridization effects: The acidity of H–A increases as the percent <i>s</i> -character of A: [–] increases.	CH ₃ CH ₃	CH ₂ =CH ₂	H–C≡C–H															
	<p>→ Increasing acidity</p>																	

30

How To: Determine the Relative Acidity of Protons

Step [1] Identify the atoms bonded to hydrogen, and use periodic trends to assign relative acidity.

- The most common H-A bonds in organic compounds are C-H, N-H and O-H.
- Acidity increases left to right across a row, the relative acidity of these bonds is C-H < N-H < O-H.
- H atoms bonded to C atoms are usually *less acidic* than H atoms bonded to any heteroatom.

Step [2] If the two H atoms in question are bonded to the same element, draw the conjugate bases and look for other points of difference.

- Do electron-withdrawing groups stabilize the conjugate base?
- Is the conjugate base resonance stabilized?
- How is the conjugate base hybridized?

31


2.6 Commonly Used Acids in Organic Chemistry

- The familiar acids HCl and H₂SO₄ are often used in organic reactions.
- Various organic acids are also commonly used (e.g., acetic acid and *p*-toluenesulfonic acid (TsOH)).

Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display



acetic acid
pK_a = 4.8




32

Commonly Used Bases in Organic Chemistry

- Common strong bases used in organic reactions are more varied in structure.

Figure 2.6


Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display



33

Characteristics of Strong Organic Bases

- Strong bases have weak conjugate acids with high pK_a values, usually > 12 .
- Strong bases have a net negative charge, but not all negatively charged species are strong bases. For example, none of the halides F^- , Cl^- , Br^- , or I^- , is a strong base.
- Carbanions, negatively charged carbon atoms, are especially strong bases. A common example is butyllithium:





34

Other Common Bases in Organic Chemistry

- Amines (e.g., triethylamine and pyridine) are organic bases.
- They are basic due to having a lone pair on N.
- They are weaker bases since they are neutral, not negatively charged.


Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display



35

2.7 Example: Aspirin

- **Aspirin (acetylsalicylic acid)** is one of the most well known drugs. It functions as an analgesic مسكن, anti-pyretic خافض الحرارة and anti-inflammatory agent مضاد للالتهابات.




- In **ionic form**, it is soluble in the aqueous blood stream and transported to tissues where it is needed.
- When **protonated**, it becomes neutral and can pass through cell membranes to inhibit certain chemical reactions.

36

2.8 Lewis Acids and Bases


- A **Lewis base** is an electron pair donor.
- Lewis bases are structurally the same as Brønsted-Lowry bases. Both have an available electron pair—a lone pair or an electron pair in a π bond.
- A Brønsted-Lowry base always donates this electron pair to a proton, but a Lewis base donates this electron pair to anything that is electron deficient.



37


Lewis Acids

- A **Lewis acid** is an electron pair acceptor.
- Any species that is electron deficient and capable of accepting an electron pair is also a Lewis acid.
- All Brønsted-Lowry acids are also Lewis acids, but the reverse is not necessarily true.
- Common Lewis acids (which are not Brønsted-Lowry acids) contain elements in group 3A that can accept an electron pair because they do not have filled valence shells.



Lewis Acid-Base Reactions

- In a Lewis acid-base reaction, a Lewis base donates an electron pair to a Lewis acid.
- This is illustrated in the reaction of BF_3 with H_2O . H_2O donates an electron pair to BF_3 to form a new bond.



39


Electrophiles and Nucleophiles

- Lewis acid-base reactions illustrate a general pattern in organic chemistry.
- Electron-rich species react with electron-poor species.
- A Lewis acid is also called an **electrophile**.
- When a Lewis base reacts with an electrophile other than a proton, the Lewis base is also called a **nucleophile**.
- In this example, BF_3 is the electrophile and H_2O is the nucleophile.

40

Lewis Acid-Base Reactions that Form One New Covalent Bond

- Note that in each reaction below, the electron pair is not removed from the Lewis base.
- Instead, it is donated to an atom of the Lewis acid and one new covalent bond is formed.



41


Drawing Products of Lewis Acid-Base Reactions

- In other Lewis acid-base reactions, one bond is formed and one bond is broken.
- To draw the products of these reactions, keep in mind the following steps:
 - Always identify the Lewis acid and base first.
 - Draw a curved arrow from the electron pair of the base to the electron-deficient atom of the acid.
 - Count electron pairs and break a bond when needed to keep the correct number of valence electrons.

42

Alkene-HCl Reaction


- The reaction between cyclohexene and HCl can be treated as a Lewis acid-base interaction.
- HCl acts as the Lewis acid, and cyclohexene, having a π bond, is the Lewis base.



43

Drawing the Product of the Reaction of HCl with Cyclohexene

- The electron pair in the π bond of the Lewis base forms a new bond to the proton of the Lewis acid, generating a carbocation.
- The H-Cl bond must break, giving its two electrons to Cl, forming Cl^- .
- Because two electron pairs are involved, two curved arrows are needed.



44