



BIRZEIT UNIVERSITY

Faculty of Engineering and Technology
Department of Electrical and Computer Engineering
Engineering Probability and Statistics ENEE 2307

Dr. Wael. Hashlamoun, Mr. Nofal.Nofal, Dr. Mohammed. Jubran, Dr. Abdalkarim Awad

Final Exam

Date: Wednesday 25/1/2017

Time: 120 minutes

Name:

Student #:

Opening Remarks:

- This is a 120-minute exam. Calculators are allowed. Mobile phones, books, notes, formula sheets, and other aids are not allowed.
- You are required to show all your work and provide the necessary explanations everywhere to get full credit.
- There are two extra problems for those who were absent in the midterm exam.

Problem 1(16 points) (ABET criterion a)

The voltage V across a $1\text{-}\Omega$ resistor R is a uniform random variable over the interval $(0, 1)$. The instantaneous power is $P = V^2/R$.

- Find the expected value of the power P .
- Find the pdf of the instantaneous power P .

Problem 2 (18 points)(ABET criterion a)

Let X and Y denote the lengths of life, in years, of two components in an electronic system. If the joint density function of these variables is

$$f_{X,Y}(x,y) = \begin{cases} k e^{-(x+y)} & 0 < x, 0 < y \\ 0 & \text{otherwise} \end{cases}$$

- Find k so that $f_{X,Y}(x,y)$ is a valid joint probability density function.
- Are X and Y statistically independent? Explain.
- Are X and Y correlated? Explain.
- Let $W=2X+3Y$, determine the standard deviation of W .

Problem 3(18 points)(ABET criterion e)

The time, in hours, it takes for computer programmer A to complete his program is a uniform random variable X , which is uniformly distributed over the interval $(0, 2)$. The time it takes for programmer B is also a random variable Y (independent of X), which is uniformly distributed over the interval $(0, 2)$.

- Write down the marginal probability density functions $f_X(x)$ and $f_Y(y)$.
- Find the joint probability density functions $f_{X,Y}(x,y)$ and the region over which it is defined.
- Find the probability that B needs at least twice the time needed by A to complete his program.

Problem 4(18 points) (ABET criterion e)

The weights of cement bags are normally distributed with a mean of (50) kg and a standard deviation of 2 kg.

- a. What is the probability that one randomly selected cement bag will weigh more than 52 kg?
- b. What is the probability that 5 randomly selected cement bags will have a mean weight of more than 52 kg
- c. Find n, such that the probability that the mean weight of n randomly selected cement bags be larger than 51 kg is less than 0.01.

Problem 5 (14 points) (ABET criterion a)

Given a random sample X_1, X_2, \dots, X_n of size n drawn from a distribution with pdf

$$f(x) = \theta e^{-\theta x}, x > 0.$$

Find a maximum likelihood estimator for the unknown parameter θ in terms of the observations.

Problem 6 (16 points) (ABET criterion e)

An electrical firm manufactures light bulbs that have a length of life that is approximately normally distributed with a standard deviation of 40 hours.

- a. If a sample of 30 bulbs has an average life of 780 hours, find a 96% confidence interval for the population mean of all bulbs produced by this firm.
- b. How large a sample is needed if we wish to be 96% confident that our sample mean will be within 10 hours of the true mean?

End of Exam

Good Luck

Problem 1 :


$$P = V^2/R \quad ; \quad R = 1 \, \Omega \Rightarrow$$

$$P = V^2$$

a. $E(P) = E(V^2)$

$$= \int_0^1 V^2 f_V(V) dV$$

$$= \int_0^1 V^2 (1) dV = \frac{V^3}{3} \Big|_0^1 = \frac{1}{3}$$



b. $f_P(P) = \frac{f_V(V)}{|dP/dV|} = \frac{(1)}{2V}$ "one-to-one mapping)

$$= \frac{(1)}{2\sqrt{P}}$$

when $v=0 \Rightarrow P=0$

$v=1 \Rightarrow P=1$

$$\Rightarrow f_P(P) = \begin{cases} \frac{1}{2\sqrt{P}} & 0 < P \leq 1 \\ 0 & \text{o.w.} \end{cases}$$

Problem 2: $f_{x,y}(x,y) = \begin{cases} k e^{-(x+y)} & x > 0, y > 0 \quad (2) \\ 0 & \text{o.w.} \end{cases}$

a. $\int_0^{\infty} \int_0^{\infty} k e^{-(x+y)} dx dy = 1 \quad \Rightarrow \quad k = 1$

b. $f_x(x) = \int_0^{\infty} k e^{-(x+y)} dy = e^{-x} \quad x > 0$

$f_y(y) = \int_0^{\infty} k e^{-(x+y)} dx = e^{-y} \quad y > 0$

since $f_{x,y}(x,y) = f_x(x) \cdot f_y(y) \Rightarrow x$ & y are indep.

c. Independence \Rightarrow uncorrelated $\Rightarrow \rho = 0$

d. $w = 2x + 3y$

$\sigma_w^2 = 4\sigma_x^2 + 9\sigma_y^2$

$f_x(x) = e^{-x} u(x) \Rightarrow \sigma_x^2 = \frac{1}{(\lambda^2)} = \frac{1}{(1)} = 1$

$\sigma_w^2 = 4(1) + 9(1)$


$= 13$

$\sigma_w = \sqrt{13}$

(3)

$$a. f_x(x) = \begin{cases} \frac{1}{2} & 0 < x \leq 2 \\ 0 & \text{o.w.} \end{cases}$$

$$f_y(y) = \begin{cases} \frac{1}{2} & 0 < y \leq 2 \\ 0 & \text{o.w.} \end{cases}$$



$$b. f_{x,y}(x,y) = \begin{cases} \frac{1}{4} & 0 < x \leq 2 \cap \\ & 0 < y \leq 2 \\ 0 & \text{o.w.} \end{cases}$$

$$c. P(Y \geq 2X) = \int_0^1 \int_{2x}^2 \left(\frac{1}{4}\right) dy dx$$

$$= \frac{1}{4} \int_0^1 (2 - 2x) dx$$

$$= \frac{1}{4} \left[2x - x^2 \right]_0^1 = \frac{1}{4} [2 - 1] = \frac{1}{4}$$

Problem 4

(4)

$$\mu_x = 50, \sigma_x = 2$$

$$\begin{aligned} \text{a. } P(X > 52) &= 1 - \Phi\left(\frac{52 - \mu_x}{\sigma_x}\right) = 1 - \Phi\left(\frac{52 - 50}{2}\right) \\ &= 1 - \Phi(1) = 1 - 0.8413 = 0.1587 \text{ From } \Phi \text{ table} \\ &= \boxed{Q(1) = 0.15866} \text{ from } Q \text{ functions} \end{aligned}$$

$$\text{b. } Y_1 = \frac{\sum x_i}{5} \Rightarrow \mu_y = \mu_x = 50$$

$$\sigma_y^2 = \frac{\sigma_x^2}{n} = \frac{4}{5} \Rightarrow \sigma_y = \frac{2}{\sqrt{5}} = \sqrt{0.8}$$

$$\begin{aligned} P(Y > 52) &= 1 - \Phi\left(\frac{52 - \mu_y}{\sigma_y}\right) = 1 - \Phi\left(\frac{52 - 50}{\sqrt{0.8}}\right) \\ &= 1 - \Phi(2.236) = 1 - 0.9871 = \boxed{0.0129} \\ &= Q(2.236) \approx \frac{Q(2.0) + Q(2.25)}{2} \approx \boxed{0.013} \end{aligned}$$

$$\text{c. } Y_2 = \frac{\sum x_i}{n} \Rightarrow E(Y_2) = \mu_x = 50$$

$$\sigma_{Y_2} = \frac{\sigma_x}{\sqrt{n}} = \frac{2}{\sqrt{n}}$$

$$P(Y_2 > 51) < 0.01$$

$$1 - \Phi\left(\frac{51 - 50}{2/\sqrt{n}}\right) \leq 0.01 \Rightarrow \Phi\left(\frac{51 - 50}{2/\sqrt{n}}\right) = 0.99$$

$$\Phi\left(\frac{\sqrt{n}}{2}\right) = 0.99 \Rightarrow \frac{\sqrt{n}}{2} = 2.33$$

$$\Rightarrow n = (2 \times 2.33)^2 = 21.7196$$

$$\Rightarrow \boxed{n = 22}$$

$$Q\left(\frac{\sqrt{n}}{2}\right) = 0.01 \Rightarrow \frac{\sqrt{n}}{2} \approx 2.3 \Rightarrow n = 21.16$$

$$\Rightarrow \boxed{n = 22}$$

Problem 5

(5)

$$f_x(x; \theta) = \theta e^{-\theta x}$$

$$L(x_1, x_2, \dots, \theta) = \theta e^{-\theta x_1} \cdot \theta e^{-\theta x_2} \dots \theta e^{-\theta x_n}$$

$$L = \theta^n e^{-\theta \sum x_i}$$

$$\ln L(\theta) = n \ln \theta - \theta \sum x_i$$

$$\frac{\partial \ln L(\theta)}{\partial \theta} = \frac{n}{\theta} - \sum x_i = 0$$

$$\frac{n}{\hat{\theta}} = \sum x_i \Rightarrow \boxed{\hat{\theta} = \frac{n}{\sum x_i}}$$

Problem 6

a.

$$\sigma_x = 40$$

$$n = 30$$

$$P\left(\hat{\mu} - z_{\alpha/2} \frac{\sigma}{\sqrt{n}} \leq \mu_x \leq \hat{\mu} + z_{\alpha/2} \frac{\sigma}{\sqrt{n}}\right) \geq 1 - \alpha$$

$$\alpha = 0.04$$

$$\frac{\alpha}{2} = 0.02$$

$$z_{\alpha/2} \approx 2.05$$

$$\Rightarrow P\left(780 - 2.05 \frac{40}{\sqrt{30}} \leq \mu_x \leq 780 + 2.05 \frac{40}{\sqrt{30}}\right) \geq 0.96$$

$$P(780 - 14.971 \leq \mu_x \leq 780 + 14.971) \geq 0.96$$

$$P(765.02 \leq \mu_x \leq 794.97) \geq 0.96$$

b. need $z_{\alpha/2} \frac{\sigma}{\sqrt{n}} = 10$

$$2.05 \frac{40}{\sqrt{n}} = 10$$

$$\Rightarrow \sqrt{n} = \frac{2.05 * 40}{10} \Rightarrow 8.2$$

$$\Rightarrow n = 67.24$$

$$\Rightarrow \boxed{n \geq 68}$$