


CHAPTER SEVEN

Membrane Structure and Function




Overview: Life at the Edge

- The plasma membrane is the boundary that separates the living cell from its surroundings
 - The plasma membrane exhibits **selective permeability**, allowing some substances to cross it more easily than others
-


Fig. 6-8

Surface area increases while total volume remains constant



<p>Total surface area [Sum of the surface areas (height × width) of all boxes sides × number of boxes]</p>	6	150	750
<p>Total volume [height × width × length × number of boxes]</p>	1	125	125
<p>Surface-to-volume (S-to-V) ratio [surface area ÷ volume]</p>	6	<u>1.2</u>	6

Figure 7.1




Concept: Cellular membranes are fluid mosaics of lipids and proteins

- **Phospholipids** are the most abundant lipid in the plasma membrane
 - Phospholipids are **amphipathic molecules**, containing **hydrophobic** and **hydrophilic** regions
 - The **fluid mosaic model** *states that a membrane is a fluid structure with a “mosaic” of various proteins embedded in it*
-

Membrane Models: *Scientific Inquiry*


- Membranes have been chemically analyzed and found to be **made of proteins and lipids**
 - Scientists studying the plasma membrane reasoned that **it must be a phospholipid bilayer**
-

Figure 7.2




-
- In 1972, **J. Singer** and **G. Nicolson** proposed the **fluid mosaic model** which states that the membrane is a *mosaic of proteins dispersed within the bilayer, with only the hydrophilic regions exposed to water*
-

Figure 7.3



-
- **Freeze-fracture studies** of the plasma membrane supported the fluid mosaic model
 - Freeze-fracture is a specialized preparation technique that **splits a membrane along the middle of the phospholipid bilayer**
-

TECHNIQUE



RESULTS



Inside of extracellular layer



Inside of cytoplasmic layer

Figure 7.4a



Inside of extracellular layer

© 2011 Pearson Education, Inc.

Figure 7.4b



Inside of cytoplasmic layer

© 2011 Pearson Education, Inc.

The **Fluidity** of Membranes

- ***Phospholipids*** in the plasma membrane **can move within the bilayer**
 - **Most of the lipids, and some proteins, drift laterally**
 - **Rarely does a molecule **flip-flop transversely** across the membrane**
-

Figure 7.5






Figure 7.6





RESULTS



-
- As **temperatures cool**, membranes *switch from a fluid state to a solid state*
 - The temperature at which a membrane solidifies depends on the types of lipids
 - Membranes rich in unsaturated fatty acids are more fluid than those rich in **saturated fatty acids**
 - Membranes **must be fluid to work properly**; they are usually about as fluid as salad oil
-


-
- The **steroid cholesterol** has different effects on membrane fluidity at different temperatures
 - **At warm temperatures** (such as 37°C), cholesterol restrains movement of phospholipids
 - **At cool temperatures**, it maintains fluidity by preventing tight packing
-

Figure 7.8



(a) Unsaturated versus saturated hydrocarbon tails

(b) Cholesterol within the animal cell membrane




Membrane Proteins and Their Functions


- A membrane is a collage of different proteins embedded in the fluid matrix of the lipid bilayer
 - Proteins **determine most of the membrane's specific functions**
-

-
- **Peripheral proteins** are bound to the surface of the membrane
 - **Integral proteins** penetrate the hydrophobic core
 - Integral proteins that span the membrane are called **transmembrane proteins**
 - The hydrophobic regions of an integral protein consist of one or more stretches of nonpolar amino acids, *often coiled into alpha helices*
-


Figure 7.9




-
- **Six major functions of membrane proteins:**
 - **Transport**
 - **Enzymatic activity**
 - **Signal transduction**
 - **Cell-cell recognition**
 - **Intercellular joining**
 - **Attachment to the cytoskeleton and extracellular matrix (ECM)**
-




(a) Transport




(b) Enzymatic activity




(c) Signal transduction



(d) Cell-cell recognition



(e) Intercellular joining



(f) Attachment to the cytoskeleton and extracellular matrix (ECM)


The Role of Membrane Carbohydrates in Cell-Cell Recognition

- Cells recognize each other by binding to **surface molecules**, often carbohydrates, on the plasma membrane
 - Membrane carbohydrates may be **covalently bonded to lipids (forming glycolipids)** or more commonly to **proteins (forming glycoproteins)**
-

Synthesis and Sidedness of Membranes

- Membranes have **distinct inside and outside faces**
 - The **asymmetrical distribution** of proteins, lipids, and associated carbohydrates in the plasma membrane is determined when the membrane is built by the ER and Golgi apparatus
-

Figure 7.12



Concept: Membrane structure results in selective permeability

- A cell must exchange materials with its surroundings, a process controlled by the plasma membrane
 - Plasma membranes are **selectively permeable**, regulating the cell's molecular traffic
-

The Permeability of the Lipid Bilayer

- **Hydrophobic (nonpolar) molecules, such as hydrocarbons, can dissolve in the lipid bilayer and pass through the membrane rapidly**
 - **Polar molecules, such as sugars, do not cross the membrane easily**
-

Transport Proteins


- **Transport proteins** allow passage of hydrophilic substances across the membrane
 - Some transport proteins, called 1) **channel proteins**, have a hydrophilic channel that certain molecules or ions can use as a tunnel
 - Channel proteins called **aquaporins** facilitate the passage of water
-

-
- Other transport proteins, called **carrier proteins**, *bind to molecules and change shape to shuttle them across the membrane*
 - A transport protein is **specific** for the substance it moves
-

Concept: Passive transport is diffusion of a substance across a membrane with **no energy investment**

- **Diffusion:** the tendency for molecules to spread out evenly into available space
 - Although each molecule moves randomly, diffusion of a population of molecules may exhibit a net movement in one direction
 - At dynamic equilibrium, as many molecules cross one way as cross in the other direction
-

Extracellular
fluid



Cytoplasm






Figure 7.13



(a) Diffusion of one solute




(b) Diffusion of two solutes

-
- **Substances diffuse down their concentration gradient, the difference in concentration of a substance from one area to another**
 - **No work must be done to move substances down the concentration gradient**
-

Effects of Osmosis on Water Balance


- **Osmosis is the diffusion of water across a selectively permeable membrane**
 - **Water diffuses across a membrane from the region of lower solute concentration to the region of higher solute concentration**
-

Figure 7.14




Water Balance of Cells Without Walls

- **Tonicity** *is the ability of a solution to cause a cell to gain or lose water*
 - **Isotonic solution:** Solute concentration is the same as that inside the cell; no net water movement across membrane
 - **Hypertonic solution:** Solute concentration is greater than that inside the cell; cell loses water
 - **Hypotonic solution:** Solute concentration is less than that inside the cell; cell gains water
-



-
- **Hypertonic** or **hypotonic** environments create osmotic problems for organisms
 - **Osmoregulation**, the control of water balance, is a necessary adaptation for life in such environments
 - The protist *Paramecium*, which is hypertonic to its pond water environment, has a **contractile vacuole** that **acts as a pump**
-

Figure 7.16



Water Balance of Cells with Walls


- Cell walls help maintain water balance
 - A plant cell in a **hypotonic solution** swells until the wall opposes uptake; the cell is now **turgid (firm)**
 - If a plant cell and its surroundings are isotonic, there is no net movement of water into the cell; the cell becomes **flaccid (limp)**, and the plant **may wilt**
-

-
- In a **hypertonic environment**, plant **cells lose water**; eventually, the membrane pulls away from the wall, a usually lethal effect called **plasmolysis**
-


Facilitated Diffusion: Passive Transport Aided by Proteins

- In **facilitated diffusion**, transport proteins speed the passive movement of molecules across the plasma membrane
 - **Channel proteins** provide corridors that allow a specific molecule or ion to cross the membrane
 - Channel proteins include
 - **Aquaporins**, for facilitated diffusion of water
 - **Ion channels** that open or close in response to a stimulus (**gated channels**)
-

Figure 7.17



(a) A channel protein




(b) A carrier protein

-
- **Carrier proteins** undergo a subtle change in shape that translocates the solute-binding site across the membrane
-

Concept: **Active transport uses energy to move solutes against their gradients**

- **Facilitated diffusion is still passive because the solute moves down its concentration gradient**
 - **Some transport proteins, however, can move solutes against their concentration gradients**
 - **Active transport requires energy, usually in the form of ATP**
 - **Active transport is performed by specific proteins embedded in the membranes**
-



Animation: Active Transport

Right-click slide / select "Play"

-
- **Active transport** allows cells to maintain concentration gradients that differ from their surroundings
 - The **sodium-potassium pump** is one type of active transport system
-

Figure 7.18-1




Figure 7.18-2




Figure 7.18-3




Figure 7.18-4




Figure 7.18-5




Figure 7.18-6






Figure 7.19

Passive transport

Active transport




How Ion Pumps Maintain Membrane Potential?

- **Membrane potential** is the voltage difference across a membrane
 - Voltage is created by differences in the distribution of **positive and negative ions** across a membrane.
-

-
- **Two combined forces, collectively called the **electrochemical gradient**, drive the diffusion of ions across a membrane:**
 - **A chemical force (the ion's concentration gradient)**
 - **An electrical force (the effect of the membrane potential on the ion's movement)**
-

-
- An **electrogenic pump** is a transport protein that **generates voltage across a membrane**
 - The **sodium-potassium pump** is the major electrogenic pump of **animal cells**
 - The main electrogenic pump of **plants, fungi, and bacteria** is a **proton pump**
-


Figure 7.20



Cotransport: Coupled Transport by a Membrane Protein

- **Cotransport** *occurs when active transport of a solute indirectly drives transport of another solute*
 - **Plants commonly use the gradient of hydrogen ions generated by proton pumps to drive active transport of nutrients into the cell**
-

Figure 7.21



Concept: **Bulk transport** across the plasma membrane occurs by **exocytosis** and **endocytosis**

- **Small molecules and water enter or leave the cell through the lipid bilayer or by transport proteins**
 - **Large molecules**, such as polysaccharides and proteins, **cross the membrane in bulk via vesicles**
 - **Bulk transport requires energy**
-

Exocytosis



- In **exocytosis**, transport vesicles migrate to the membrane, fuse with it, and release their contents
 - Many **secretory cells** use exocytosis to export their products
-

Endocytosis


- In **endocytosis**, the cell takes in macromolecules by forming vesicles from the plasma membrane
 - Endocytosis is a reversal of exocytosis, involving different proteins
 - There are **three types of endocytosis**:
 - **Phagocytosis** (“cellular eating”)
 - **Pinocytosis** (“cellular drinking”)
 - **Receptor-mediated endocytosis**
-

-
- In **phagocytosis** a cell engulfs a particle in a vacuole
 - The vacuole **fuses with a lysosome** to digest the particle
-

PHAGOCYTOSIS



An amoeba engulfing a bacterium via phagocytosis (TEM)





Animation: Phagocytosis


Right-click slide / select "Play"

-
- In **pinocytosis**, molecules are taken up when extracellular fluid is “gulped” into tiny vesicles
-

PINOCYTOSIS



Pinocytosis vesicles forming (arrows) in a cell lining a small blood vessel (TEM)



-
- **In receptor-mediated endocytosis, binding of ligands to receptors triggers vesicle formation**
 - **A ligand is any molecule that binds specifically to a receptor site of another molecule**
-







Figure 7.22



Familial *hypercholesterolemia* (FH)

- **Cholesterol is transported in blood in particles called LDL**
- **LDL binds to receptors on plasma membranes & enter cells by receptor-mediated endocytosis**
- **In FH, LDL receptors are defective or missing >> cholesterol accumulates in blood vessels and lead to atherosclerosis**