

Prescott's

MICROBIOLOGY

ELEVENTH EDITION


JOANNE WILLEY

KATHLEEN SANDMAN

DOROTHY WOOD

Chapter 35

Infection and Pathogenicity



Infection

Host.

- Larger organism that supports the survival and growth of a smaller organism. ✓

Infection.

- A microbe growing and multiplying on or within a host.
- May or may not result in overt infectious disease.

Infectious disease.

- Any change from a state of health. ✓
- Part or all of the host is incapable of carrying on normal functions due to presence of a pathogen or its products.

Pathogenicity

Pathogen.

- Any organism that causes disease.
- Opportunistic pathogen—may be part of normal flora and causes disease when it gains access to other tissue sites or when the host is immunocompromised.

Pathogenicity.

- Ability of a pathogen to cause disease.

Virulence.

- Degree of harm (pathogenicity) inflicted on the host.

Extracellular vs. Intracellular Pathogens

A pathogen must contact a host AND survive within it to cause a disease.

Extracellular pathogens.

- Grow outside host cells in tissues and fluids.

Intracellular pathogens.

- Grow and multiply within host cells.
- **Facultative intracellular pathogens**—reside within the cells of the host or in the environment but can also be grown in pure culture without host cell support.
- **Obligate intracellular pathogens**—only grow when inside host cells.

Course of Infectious Disease

1 Incubation period. ✓

- Period after pathogen entry, before signs and symptoms.

2 Prodromal stage. ✓

- Onset of signs and symptoms.
- Not clear enough for diagnosis.


3 Illness period. ✓

- Disease is most severe, signs and symptoms.

4 Convalescence. ✓

- Signs and symptoms begin to disappear.

Copyright © McGraw-Hill Education. All rights reserved. No reproduction or distribution without the prior written consent of McGraw-Hill Education.



[Access the text alternative for these images](#)

Course of Infectious Disease—Important Terms

Signs.

- Objective changes in the body that can be directly observed.
- For example, fever, rash.

Symptoms.

- Subjective changes experienced by patient.
- For example, pain, loss of appetite.

Disease syndrome.

- Set of characteristic signs and symptoms for a disease.

Importance of Resources

The host provides the pathogen:

- Protection. ✓
- Nutrients. ✓
- Energy to use. ✓

Infectious agents develop mechanisms to access and exploit hosts.

- To survive, must also devise methods to move on to a new location or host when necessary.

→ Some microorganism can infect.

Events in Infection and Disease

Transmission from previous host or reservoir to new host.

- Virulence of organism, number of invading organisms, and presence of adhesion and invasion factors will affect success.

Organism outcompetes the resident microbiota for resources and survives host defense mechanisms.

Disease occurs when organism produces molecules that directly damage host cells OR stimulates host immune cells to destroy infected tissue OR alters the host cell genome affecting normal function.

Sources of Pathogens

Can be **animate** (other **humans** or **animals**).

- Infections passed from animal to human are termed **zoonoses**.

Can be **inanimate** (**water**, **food**).

Reservoir.

- Natural environmental location in which the pathogen normally resides.

Vector.

- Organism that spreads disease from one host to another.
- For example, mosquitoes, ticks, fleas, mites, or biting flies.

Pathogen Transmission

Occurs either directly or indirectly.

Four main routes:


- Airborne.
- Contact.
- Vehicle.
- Vector-borne.

Pregnant women can also pass a pathogen to their unborn child via vertical transmission.

Placentes.

Transmission of Infectious Diseases

Copyright © McGraw-Hill Education. All rights reserved. No reproduction or distribution without the prior written consent of McGraw-Hill Education.



[Access the text alternative for these images](#)

Transmission and Virulence

Pathogen's virulence may be strongly influenced by its mode of transmission and ability to live outside its host.

Transmission alone is not enough for infection to occur.

- Tropism—pathogen must make contact with appropriate host tissue; determined by specific cell surface receptors.

Airborne Transmission

Droplets—direct transmission.

- Up to 2 mm in diameter.
- Produced when liquids are placed under force (For example, sneezing, coughing).
- Can travel <1 m.

Droplet nuclei—indirect transmission.

- 1 to 5 μm in diameter.
- Result from evaporation of the larger droplets.
- May remain airborne for hours or days and travel long distances.

Dust particles—indirect transmission.

- Microorganisms adhere to dust particles.
- Can survive long periods outside host and can lead to hospital-acquired infections.

Contact Transmission

Coming together or touching of source/reservoir and host.

Direct contact (person-to-person).

- Physical interaction between source/reservoir and host.
- For example, kissing, touching, and sexual contact.

Indirect contact.

- Involves an inanimate object (fomite).
- For example, eating utensils, bedding.

Vehicle Transmission

Copyright © McGraw-Hill Education. All rights reserved. No reproduction or distribution without the prior written consent of McGraw-Hill Education.



Vehicles.

- **Inanimate materials that transmit pathogens.**
- For example, **food, water, biological materials (fluids and tissues), air.**

Vehicle transmission.

- **Single vehicle spreads pathogen to multiple hosts.**

Vector-Borne Transmission

Vector—direct living transmitter of a pathogen.

- Most are arthropods (For example, insects, ticks, mites, fleas) or vertebrates (For example, dogs, cats, skunks, bats).

Pathogens transmitted by arthropods are often highly virulent.

- Cause diseases such as malaria, typhus, and sleeping sickness.

Important that pathogens do not harm their vectors.

Vertical Transmission

Pregnant women

Occurs when the unborn child acquires a pathogen from an infected mother.

Kiss - / Sex

Not as common as horizontal transmission.

Babies born with an infectious disease are said to have a congenital infection.

Examples include:

- Gonorrhea.
- Syphilis.
- Herpes.
- German measles.
- Toxoplasmosis.


Infectious Dose

Infectious dose 50 (ID₅₀).

- Number of pathogens that will infect 50% of inoculated hosts.
- Varies with pathogen.

Hand washing reduces number of pathogens.

Copyright © McGraw-Hill Education. All rights reserved. No reproduction or distribution without the prior written consent of McGraw-Hill Education.



[Access the text alternative for these images](#)

Lethal Dose

Lethal dose 50 (LD50).

- Dose that kills 50% of experimental animals within a specified period.

Cytopathology—cellular changes.

- Can be used to observe cells in tissue culture for death rates rather than entire organisms.

Examining virulence factors and their release.

Adherence and Colonization

First steps in disease are entry and attachment

Portal of entry.

- Skin, respiratory, gastrointestinal, urogenital systems, or conjunctiva of eye.

Adherence.

- Mediated by special molecules called adhesins.

Colonization.


- A site of microbial replication on or within host.
- Does not necessarily result in tissue invasion or damage.

Microbial Adherence Mechanisms


Adherence structures bind complementary receptor sites on host cell surface; classified as virulence factors.

- Pili.
- Fimbriae.
- Membrane and capsular materials.
- Specialized adhesion molecules on bacterium's cell surface.


Copyright © McGraw-Hill Education. All rights reserved. No reproduction or distribution without the prior written consent of McGraw-Hill Education.



(a) Fimbriae (F)



(b) Capsules (C)



(c) Spikes

Invasion Disseminates Pathogens

Infectivity—ability to create a discrete point of infection

Invasiveness—ability to spread to adjacent tissues.

Penetration can be active or passive:

- Active occurs through lytic substances that alter host tissue by (1) **attacking** the extracellular matrix and basement membranes of **integuments and intestinal linings**, (2) **degrading carbohydrate-protein complexes** between cells or on the **cell surface**, or (3) **disrupting the host cell surface**.
- Passive—not related to the pathogen itself (For example, **skin lesions, insect bites, wounds**).

Invasion—Specific Examples

Once in circulatory system, bacteria have access to all organs and systems.

- Bacteremia—presence of viable bacteria in the blood.
- Septicemia—bacterial or fungal toxins in the blood.

Varies among pathogens.

- For example, *Clostridium tetani* (tetanus) produces a number of virulence factors but is noninvasive.
- For example, *Bacillus anthracis* (anthrax) and *Yersinia pestis* (plague) also produce many virulence factors and are highly invasive.
- For example, *Streptococcus* spp. span the spectrum of virulence factors and invasiveness.

Overcoming Host Defenses

Most microbes eliminated before they can cause disease due to immune system.

Successful pathogens overcome the competition and elude initial host responses as well as the adaptive immune system:

- Produce Type VI secretion system.
- Find shelter to avoid recognition by defense cells.
- Survive and replicate inside host cells.
- Squeeze between host cells.
- Make capsules to avoid phagocytosis.
- Burrow under mucus.
- Secrete exopolysaccharides to form communal shelters within biofilms.
- Produce enzymes that inactivate innate resistance mechanisms.
- Excrete specialized proteins to selectively kill host cells.

Strategies to Evade Host Immune Response

Production of decoy proteins to bind available neutralizing antibodies.

Change cell surface proteins by mutation or recombination or downregulate the level of expression of cell surface proteins.

Produce capsules that resemble host tissue components.

- Some bacterial capsules prevent deposition of host complement, protecting the pathogen from host defenses.

Phage variation to alter pili protein sequence and expression.

Produce proteases that degrade host proteins.

Produce special proteins that interfere with the host's ability to detect and remove them.

Cause host cell fusion.


Suppressing Host Immune Response

Infect cells of the immune system and diminish their function while ensuring their own survival.

Bacteria such as *Streptococcus pneumoniae*, *Neisseria meningitidis*, and *Haemophilus influenzae* produce a slippery mucoid capsule that prevents the host immune cells from effectively capturing the bacterium.

Eliminate O-antigen on lipopolysaccharide to diminish immune response and clearance.

Biofilms



Biofilm bacteria are protected from nutrient deprivation, predators, environmental shifts, antimicrobial agents, and host immune cells.

Some pathogenic bacteria within biofilms exchange plasmids, nutrients, and quorum-sensing molecules, altering their behavior.

- Make them less sensitive to antibiotics and more resistant to host defense mechanisms.

[Access the text alternative for these images](#)

Pathogenicity Islands

Large segments (10 to 200 kilobases) of bacterial chromosomal and plasmid DNA found to encode virulence factors.

- Increase bacterial virulence.
- Absent in nonpathogenic members of same genus or species.
- Can be spread through horizontal transfer of virulence genes to bacteria.

Toxigenicity

Some microbes possess toxigenicity.

- Ability to produce toxins.

Toxin.

- Specific substance that damages host.

Intoxications.

- Diseases that result from entry of a specific preformed toxin into host.

Exotoxins

Soluble, heat-labile proteins.

Secreted into surroundings as pathogen metabolizes.

Often travel from site of infection to other tissues or cells where they exert their effects.

Usually synthesized by specific bacteria that have toxin genes in their plasmids or prophage DNA.

Among the most lethal substances known.

Types of Exotoxins


AB toxins.

- Composed of two subunits: A subunit (responsible for toxic effect) and B subunit (binds to specific target cell).


Specific host site exotoxins.

Membrane-disrupting exotoxins.

Superantigens.




Copyright © McGraw-Hill Education. All rights reserved. No reproduction or distribution without the prior written consent of McGraw-Hill Education.



[Access the text alternative for these images](#)

Superantigens



Stimulate about 30% of T cells of the immune system.

- Causes the T cells to overexpress and release pro-inflammatory cytokines.
- Results in failure of multiple host organs allowing time for the microbe to disseminate.

[Access the text alternative for these images](#)

Endotoxin—Lipopolysaccharide

Lipopolysaccharide (LPS) in Gram-negative cell wall can be toxic to specific hosts.

- Called endotoxin because it is an endogenous , bound to the bacterium and released when the microorganism lyses; some is also released during multiplication.
- Toxic component is the lipid portion, lipid A.

Endotoxins—General Features

Heat stable.

Toxic (nanogram amounts).

Weakly immunogenic.


Generally similar, despite source.

Cause general system effects:

- Fever, shock, damage to the lining of blood vessels, weakness, diarrhea, inflammation, intestinal hemorrhage, and fibrinolysis.

The Septic Shock Cascade

Copyright © McGraw-Hill Education. All rights reserved. No reproduction or distribution without the prior written consent of McGraw-Hill Education.



[Access the text alternative for these images](#)

Mycotoxins

Toxins produced by fungi.

Common contaminants of food crops.

- For example, *Aspergillus flavus* and *A. parasiticus* produce aflatoxins, cause chronic and acute liver disease and liver cancer.

Contaminate water-damaged buildings.

- For example, *Stachybotrys* produce satratoxins, potent inhibitors of DNA, RNA, and protein synthesis; induce inflammation, disrupt surfactant phospholipids in the lungs, and may lead to pathological changes in tissues.

Claviceps purpurea (ergot) produce hallucinogen lysergic acid (LSD).